



Full wwPDB X-ray Structure Validation Report ⓘ

Aug 5, 2026 – 05:08 AM JST

PDB ID : 5B46 / pdb_00005b46
Title : 2-Oxoacid:Ferredoxin Oxidoreductase 2 from Sulfolobus tokodai - ligand free form
Authors : Yan, Z.; Maruyama, A.; Arakawa, T.; Fushinobu, S.; Wakagi, T.
Deposited on : 2016-04-01
Resolution : 2.10 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4-5-2 with Phenix2.0
Mogul	:	1.8.5 (274361), CSD as541be (2020)
Xtriage (Phenix)	:	2.0
EDS	:	3.0
Buster-report	:	wwPDB partial adaption of 1.1.7 (2018)
Percentile statistics	:	20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4	:	9.0.015 (Gargrove)
Density-Fitness	:	1.0.12
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.50


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.10 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	6658 (2.10-2.10)
Clashscore	190562	7164 (2.10-2.10)
Ramachandran outliers	187476	7099 (2.10-2.10)
Sidechain outliers	187428	7100 (2.10-2.10)
RSRZ outliers	180081	6662 (2.10-2.10)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	628	4% 90% 7% .
2	B	304	10% 87% 11% ..

2 Entry composition [i](#)

There are 6 unique types of molecules in this entry. The entry contains 7592 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.


- Molecule 1 is a protein called 2-oxoacid--ferredoxin oxidoreductase alpha subunit.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	627	Total	C	N	O	S	0	0	0
			4937	3159	822	937	19			

- Molecule 2 is a protein called 2-oxoacid--ferredoxin oxidoreductase beta subunit.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
2	B	301	Total	C	N	O	S	0	0	0
			2357	1513	409	425	10			

- Molecule 3 is IRON/SULFUR CLUSTER (CCD ID: SF4) (formula: Fe₄S₄).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
3	B	1	Total	Fe	S	0	0
			8	4	4		

- Molecule 4 is THIAMINE DIPHOSPHATE (CCD ID: TPP) (formula: C₁₂H₁₉N₄O₇P₂S).



- Molecule 5 is MAGNESIUM ION (CCD ID: MG) (formula: Mg).


- Molecule 6 is water.

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
6	A	192	Total O 192 192	0	0
6	B	71	Total O 71 71	0	0


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: 2-oxoacid--ferredoxin oxidoreductase alpha subunit



- Molecule 2: 2-oxoacid--ferredoxin oxidoreductase beta subunit



4 Data and refinement statistics

Property	Value	Source
Space group	C 2 2 21	Depositor
Cell constants a, b, c, α , β , γ	101.89Å 205.05Å 126.99Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	33.93 – 2.10 33.93 – 2.10	Depositor EDS
% Data completeness (in resolution range)	98.7 (33.93-2.10) 98.7 (33.93-2.10)	Depositor EDS
R_{merge}	0.07	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	3.34 (at 2.10Å)	Xtriage
Refinement program	REFMAC 5.8.0135	Depositor
R, R_{free}	0.175 , 0.219 0.182 , 0.224	Depositor DCC
R_{free} test set	3866 reflections (4.98%)	wwPDB-VP
Wilson B-factor (Å ²)	35.4	Xtriage
Anisotropy	0.081	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.35 , 26.2	EDS
L-test for twinning ²	$\langle L \rangle = 0.49$, $\langle L^2 \rangle = 0.33$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.96	EDS
Total number of atoms	7592	wwPDB-VP
Average B, all atoms (Å ²)	42.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 4.21% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: TPP, MG, SF4

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	1.33	10/5031 (0.2%)	1.17	8/6797 (0.1%)
2	B	1.41	6/2412 (0.2%)	1.24	11/3267 (0.3%)
All	All	1.35	16/7443 (0.2%)	1.19	19/10064 (0.2%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	A	0	1

All (16) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	B	160	GLY	N-CA	-8.04	1.32	1.45
1	A	424	ASN	C-O	-6.49	1.15	1.24
2	B	146	ILE	CA-C	6.21	1.59	1.52
1	A	333	ASN	N-CA	-5.72	1.38	1.46
1	A	455	ARG	CA-C	5.55	1.59	1.52
2	B	5	LYS	CA-C	5.38	1.58	1.52
2	B	50	ILE	CA-CB	5.35	1.56	1.54
2	B	275	TYR	N-CA	5.28	1.52	1.46
2	B	191	VAL	C-O	-5.23	1.18	1.24
1	A	543	MET	SD-CE	-5.20	1.66	1.79
1	A	288	VAL	CA-CB	-5.16	1.47	1.54
1	A	569	ALA	C-O	-5.15	1.17	1.24
1	A	310	ARG	C-O	-5.14	1.17	1.24
1	A	64	SER	CB-OG	5.14	1.52	1.42
1	A	326	GLY	C-O	-5.01	1.17	1.23
1	A	8	ILE	N-CA	5.00	1.51	1.46

All (19) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	B	159	ALA	N-CA-C	11.96	126.97	112.38
2	B	63	THR	N-CA-C	9.22	116.34	108.78
2	B	159	ALA	O-C-N	-8.23	111.69	122.39
2	B	102	VAL	N-CA-C	7.51	118.31	110.72
2	B	159	ALA	CA-C-N	7.13	134.53	121.70
2	B	159	ALA	C-N-CA	7.13	134.53	121.70
2	B	252	ILE	CA-CB-CG2	6.78	122.02	110.50
1	A	560	ILE	CB-CA-C	-6.44	101.83	112.26
1	A	628	ALA	CB-CA-C	-6.32	101.02	110.50
1	A	431	ARG	NE-CZ-NH2	-5.89	113.90	119.20
1	A	500	GLU	N-CA-C	5.79	119.53	112.23
1	A	408	LYS	CB-CG-CD	5.55	124.06	111.30
1	A	455	ARG	NE-CZ-NH1	5.49	126.99	121.50
2	B	282	LYS	N-CA-C	-5.32	106.78	113.28
2	B	292	THR	N-CA-CB	-5.12	102.91	110.49
1	A	122	ARG	NE-CZ-NH2	-5.11	114.61	119.20
2	B	50	ILE	N-CA-C	5.11	119.05	112.67
1	A	425	ARG	N-CA-C	5.04	118.15	110.50
2	B	63	THR	CA-C-O	5.02	120.85	117.94

There are no chirality outliers.

All (1) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	A	627	GLY	Peptide

5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	4937	0	4964	30	0
2	B	2357	0	2391	12	0
3	B	8	0	0	0	0
4	B	26	0	16	1	0
5	B	1	0	0	0	0
6	A	192	0	0	1	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
6	B	71	0	0	0	0
All	All	7592	0	7371	41	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 3.

All (41) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
2:B:284:GLN:O	2:B:292:THR:HG21	1.86	0.75
1:A:50:TYR:OH	1:A:628:ALA:HB2	1.90	0.71
1:A:258:PRO:HG3	1:A:408:LYS:HB2	1.74	0.69
1:A:556:LEU:O	1:A:560:ILE:HG13	1.94	0.67
2:B:197:CYS:O	2:B:200:TYR:O	2.16	0.63
2:B:174:HIS:CE1	2:B:252:ILE:HG23	2.34	0.62
1:A:600:ARG:HE	1:A:628:ALA:HB3	1.65	0.61
1:A:197:ASN:OD1	1:A:197:ASN:C	2.44	0.60
1:A:527:ILE:HG22	1:A:543:MET:HE3	1.85	0.58
1:A:396:GLU:OE1	1:A:431:ARG:HD2	2.04	0.57
1:A:2:THR:N	1:A:57:GLU:OE1	2.39	0.55
1:A:80:GLN:HA	1:A:110:MET:HE2	1.89	0.54
1:A:455:ARG:HD2	6:A:864:HOH:O	2.06	0.54
1:A:527:ILE:CG2	1:A:543:MET:HE3	2.37	0.54
2:B:94:LEU:O	2:B:98:MET:HE2	2.08	0.54
1:A:338:VAL:HG13	1:A:392:LEU:HD23	1.93	0.51
1:A:600:ARG:HE	1:A:628:ALA:CB	2.24	0.51
1:A:524:LYS:O	1:A:543:MET:HE1	2.11	0.50
2:B:43:GLY:O	2:B:48:GLY:HA3	2.12	0.49
1:A:167:ALA:HB1	1:A:207:ALA:HB2	1.93	0.49
1:A:527:ILE:HB	1:A:543:MET:CE	2.42	0.49
1:A:516:VAL:HG23	1:A:564:LYS:HG2	1.96	0.48
1:A:342:TYR:HA	1:A:406:ILE:O	2.15	0.46
1:A:431:ARG:HB3	1:A:434:ILE:HD12	1.97	0.46
2:B:21:LEU:HA	2:B:50:ILE:HD11	1.98	0.46
1:A:516:VAL:CG2	1:A:564:LYS:HG2	2.47	0.45
1:A:600:ARG:NE	1:A:628:ALA:HB3	2.29	0.44
2:B:58:ILE:C	2:B:58:ILE:HD12	2.43	0.44
1:A:527:ILE:CG2	1:A:543:MET:CE	2.97	0.43
1:A:355:SER:HB2	1:A:408:LYS:HG3	2.00	0.43
1:A:233:VAL:HB	1:A:238:ILE:HD11	2.01	0.43
1:A:346:ALA:HB1	1:A:347:PRO:HA	2.00	0.42

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:44:ILE:HG21	2:B:199:THR:HG21	2.00	0.42
1:A:469:GLU:HG2	1:A:483:ARG:HD3	2.01	0.42
2:B:50:ILE:HG23	2:B:51:PRO:HD3	2.02	0.42
2:B:109:ILE:HG22	2:B:110:ASP:N	2.35	0.42
2:B:121:TYR:HB3	4:B:402:TPP:H62	2.01	0.41
1:A:197:ASN:OD1	1:A:199:LEU:N	2.53	0.41
1:A:217:ASN:OD1	1:A:217:ASN:C	2.63	0.41
1:A:396:GLU:OE2	1:A:455:ARG:NH2	2.54	0.40
2:B:194:LEU:HD12	2:B:252:ILE:HD11	2.02	0.40

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	625/628 (100%)	608 (97%)	16 (3%)	1 (0%)	43	44
2	B	299/304 (98%)	289 (97%)	8 (3%)	2 (1%)	18	15
All	All	924/932 (99%)	897 (97%)	24 (3%)	3 (0%)	36	36

All (3) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
2	B	247	GLY
2	B	125	LYS
1	A	277	GLU

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	527/528 (100%)	509 (97%)	18 (3%)	32	35
2	B	254/257 (99%)	242 (95%)	12 (5%)	23	24
All	All	781/785 (100%)	751 (96%)	30 (4%)	29	32

All (30) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	2	THR
1	A	29	LYS
1	A	64	SER
1	A	183	LYS
1	A	193	SER
1	A	197	ASN
1	A	199	LEU
1	A	233	VAL
1	A	276	VAL
1	A	282	ARG
1	A	338	VAL
1	A	400	THR
1	A	408	LYS
1	A	425	ARG
1	A	436	LYS
1	A	456	VAL
1	A	537	ASP
1	A	565	SER
2	B	5	LYS
2	B	25	GLU
2	B	102	VAL
2	B	207	GLU
2	B	211	LYS
2	B	217	ASP
2	B	234	LYS
2	B	242	LYS
2	B	252	ILE
2	B	280	PRO
2	B	282	LYS
2	B	298	LEU

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (7) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	204	ASN
1	A	229	GLN
1	A	270	GLN
1	A	279	ASN
1	A	299	ASN
2	B	238	GLN
2	B	257	GLN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

Of 3 ligands modelled in this entry, 1 is monoatomic - leaving 2 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
4	TPP	B	402	5	25,27,27	1.75	4 (16%)	37,40,40	2.17	13 (35%)
3	SF4	B	401	2	0,12,12	-	-	-		

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
4	TPP	B	402	5	-	4/17/17/17	0/2/2/2
3	SF4	B	401	2	-	-	0/6/5/5

All (4) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
4	B	402	TPP	C5-C4	5.38	1.46	1.35
4	B	402	TPP	C5'-C4'	5.11	1.51	1.42
4	B	402	TPP	C2-S1	2.24	1.76	1.69
4	B	402	TPP	PB-O1B	2.14	1.57	1.50

All (13) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
4	B	402	TPP	C2-S1-C5	-6.20	86.93	91.21
4	B	402	TPP	S1-C2-N3	4.60	118.34	112.29
4	B	402	TPP	CM4-C4-C5	-3.55	118.27	127.72
4	B	402	TPP	CM4-C4-N3	3.34	128.50	120.52
4	B	402	TPP	C6'-N1'-C2'	3.14	121.31	115.96
4	B	402	TPP	PA-O3A-PB	-2.98	122.61	132.83
4	B	402	TPP	C2'-N3'-C4'	2.96	122.69	118.08
4	B	402	TPP	N1'-C2'-N3'	-2.92	120.51	125.54
4	B	402	TPP	C2-N3-C4	-2.73	110.22	114.05
4	B	402	TPP	O3B-PB-O3A	2.67	113.58	104.64
4	B	402	TPP	C5'-C4'-N3'	-2.45	117.39	121.24
4	B	402	TPP	CM2-C2'-N3'	2.25	120.66	117.15
4	B	402	TPP	O2A-PA-O1A	2.07	122.47	112.24

There are no chirality outliers.

All (4) torsion outliers are listed below:


Mol	Chain	Res	Type	Atoms
4	B	402	TPP	PA-O3A-PB-O2B
4	B	402	TPP	PA-O3A-PB-O3B
4	B	402	TPP	S1-C5-C6-C7
4	B	402	TPP	C4-C5-C6-C7

There are no ring outliers.

1 monomer is involved in 1 short contact:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
4	B	402	TPP	1	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.



5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	627/628 (99%)	-0.00	22 (3%) 47 49	23, 38, 63, 101	0
2	B	301/304 (99%)	0.47	30 (9%) 12 13	25, 44, 70, 120	0
All	All	928/932 (99%)	0.15	52 (5%) 30 32	23, 39, 66, 120	0

All (52) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
2	B	4	ARG	6.0
2	B	6	PRO	4.1
1	A	281	LEU	4.1
1	A	628	ALA	3.9
2	B	5	LYS	3.8
1	A	278	GLY	3.5
1	A	104	LYS	3.3
2	B	7	VAL	3.3
2	B	12	CYS	3.2
1	A	436	LYS	3.0
2	B	146	ILE	3.0
2	B	50	ILE	3.0
1	A	280	GLU	3.0
2	B	247	GLY	3.0
1	A	347	PRO	2.9
2	B	143	LYS	2.9
2	B	198	PRO	2.8
2	B	249	LYS	2.8
1	A	196	LYS	2.8
2	B	125	LYS	2.7
2	B	209	TYR	2.7
1	A	434	ILE	2.6
2	B	252	ILE	2.6
1	A	625	TYR	2.5

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	A	348	ALA	2.5
1	A	441	TYR	2.5
2	B	51	PRO	2.5
1	A	424	ASN	2.3
1	A	472	GLU	2.3
2	B	46	CYS	2.3
2	B	139	LYS	2.3
1	A	438	THR	2.3
2	B	200	TYR	2.3
1	A	277	GLU	2.3
2	B	120	VAL	2.3
1	A	151	LYS	2.3
2	B	124	THR	2.2
2	B	14	GLY	2.2
1	A	627	GLY	2.2
1	A	617	ARG	2.2
2	B	44	ILE	2.2
2	B	304	ILE	2.2
1	A	227	GLU	2.1
1	A	442	PHE	2.1
2	B	123	LEU	2.1
2	B	121	TYR	2.1
2	B	11	TRP	2.1
2	B	234	LYS	2.1
2	B	244	TYR	2.1
2	B	241	MET	2.0
1	A	276	VAL	2.0
2	B	170	TYR	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.


6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum,

median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
5	MG	B	403	1/1	0.85	0.10	52,52,52,52	0
4	TPP	B	402	26/26	0.95	0.12	40,50,56,59	0
3	SF4	B	401	8/8	0.96	0.09	35,38,41,45	8

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.



6.5 Other polymers ⓘ

There are no such residues in this entry.