



# wwPDB NMR Structure Validation Summary Report ⓘ

Mar 5, 2026 – 05:16 PM UTC

PDB ID : 2B95 / pdb\_00002b95  
Title : Solution NMR structure of protein dynein light chain 2A, cytoplasmic; North-east structural genomics consortium TARGET HR2106  
Authors : Liu, G.; Atreya, H.S.; Xiao, R.; Acton, T.; Montelione, G.T.; Szyperski, T.; Northeast Structural Genomics Consortium (NESG)  
Deposited on : 2005-10-10

This is a wwPDB NMR Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<https://www.wwpdb.org/validation/2017/NMRValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4-5-2 with Phenix2.0  
Percentile statistics : 20250101.v01 (using entries in the PDB archive January 1st 2025)  
wwPDB-RCI : v\_1n\_11\_5\_13\_A (Berjanski et al., 2005)  
PANAV : Wang et al. (2010)  
wwPDB-ShiftChecker : v1.2  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : 2.49


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

*SOLUTION NMR*

The overall completeness of chemical shifts assignment was not calculated.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	NMR archive (#Entries)
Clashscore	229148	14424
Ramachandran outliers	224038	12848
Sidechain outliers	223484	12823

The table below summarises the geometric issues observed across the polymeric chains and their fit to the experimental data. The red, orange, yellow and green segments indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria. A cyan segment indicates the fraction of residues that are not part of the well-defined cores, and a grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$

Mol	Chain	Length	Quality of chain
1	A	106	
1	B	106	

## 2 Ensemble composition and analysis i

This entry contains 20 models. Model 4 is the overall representative, medoid model (most similar to other models). The authors have identified model 1 as representative, based on the following criterion: *fewest violations*.

The following residues are included in the computation of the global validation metrics.

Well-defined (core) protein residues			
Well-defined core	Residue range (total)	Backbone RMSD (Å)	Medoid model
1	A:11-A:106, B:11-B:106 (192)	0.76	4

Ill-defined regions of proteins are excluded from the global statistics.

Ligands and non-protein polymers are included in the analysis.

The models can be grouped into 6 clusters and 2 single-model clusters were found.

Cluster number	Models
1	8, 9, 11, 13
2	1, 10, 14, 20
3	7, 12, 18
4	3, 6, 17
5	4, 15
6	2, 5
Single-model clusters	16; 19

### 3 Entry composition

There is only 1 type of molecule in this entry. The entry contains 3106 atoms, of which 1578 are hydrogens and 0 are deuteriums.

- Molecule 1 is a protein called Dynein light chain 2A.


Mol	Chain	Residues	Atoms						Trace
			Total	C	H	N	O	S	
1	A	96	1553	481	789	131	148	4	0
1	B	96	1553	481	789	131	148	4	0

There are 20 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	1	MET	-	cloning artifact	UNP Q9NP97
A	2	GLY	-	cloning artifact	UNP Q9NP97
A	3	HIS	-	expression tag	UNP Q9NP97
A	4	HIS	-	expression tag	UNP Q9NP97
A	5	HIS	-	expression tag	UNP Q9NP97
A	6	HIS	-	expression tag	UNP Q9NP97
A	7	HIS	-	expression tag	UNP Q9NP97
A	8	HIS	-	expression tag	UNP Q9NP97
A	9	SER	-	cloning artifact	UNP Q9NP97
A	10	HIS	-	cloning artifact	UNP Q9NP97
B	1	MET	-	cloning artifact	UNP Q9NP97
B	2	GLY	-	cloning artifact	UNP Q9NP97
B	3	HIS	-	expression tag	UNP Q9NP97
B	4	HIS	-	expression tag	UNP Q9NP97
B	5	HIS	-	expression tag	UNP Q9NP97
B	6	HIS	-	expression tag	UNP Q9NP97
B	7	HIS	-	expression tag	UNP Q9NP97
B	8	HIS	-	expression tag	UNP Q9NP97
B	9	SER	-	cloning artifact	UNP Q9NP97
B	10	HIS	-	cloning artifact	UNP Q9NP97



## ● Molecule 1: Dynein light chain 2A



## 5 Refinement protocol and experimental data overview

The models were refined using the following method: *distance geometry torsion angle dynamics simulated annealing*.

Of the 100 calculated structures, 20 were deposited, based on the following criterion: *structures with the least restraint violations*.

The following table shows the software used for structure solution, optimisation and refinement.

Software name	Classification	Version
DYANA	structure solution	1.5
CYANA	refinement	2.1

No chemical shift data was provided.

## 6 Model quality [i](#)

### 6.1 Standard geometry [i](#)

There are no covalent bond-length or bond-angle outliers.

There are no bond-length outliers.

There are no bond-angle outliers.

There are no chirality outliers.

There are no planarity outliers.

### 6.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in each chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes averaged over the ensemble.

Mol	Chain	Non-H	H(model)	H(added)	Clashes
1	A	764	789	788	29±4
1	B	764	789	788	27±4
All	All	30560	31560	31520	946

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 15.

5 of 252 unique clashes are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Clash(Å)	Distance(Å)	Models	
				Worst	Total
1:B:56:MET:HE2	1:B:90:VAL:HG21	1.01	1.31	14	7
1:A:67:VAL:HG21	1:A:76:LEU:HD21	0.91	1.42	19	5
1:B:67:VAL:HG21	1:B:76:LEU:HD23	0.86	1.47	15	11
1:B:67:VAL:HG21	1:B:76:LEU:HD21	0.85	1.48	12	5
1:A:44:MET:HE1	1:A:101:ILE:HD13	0.83	1.48	7	7

### 6.3 Torsion angles [i](#)

#### 6.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all PDB entries followed by that with respect to all NMR

entries. The Analysed column shows the number of residues for which the backbone conformation was analysed and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	94/106 (89%)	81±2 (86±2%)	13±2 (14±2%)	0±0 (0±0%)	44	80
1	B	94/106 (89%)	80±2 (86±2%)	13±2 (14±2%)	0±0 (0±1%)	31	76
All	All	3760/4240 (89%)	3223 (86%)	525 (14%)	12 (0%)	37	78

All 5 unique Ramachandran outliers are listed below. They are sorted by the frequency of occurrence in the ensemble.

Mol	Chain	Res	Type	Models (Total)
1	B	44	MET	4
1	A	44	MET	3
1	B	74	ASN	3
1	A	37	GLY	1
1	B	85	LYS	1

### 6.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all PDB entries followed by that with respect to all NMR entries. The Analysed column shows the number of residues for which the sidechain conformation was analysed and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	89/98 (91%)	62±4 (70±4%)	27±4 (30±4%)	1	16
1	B	89/98 (91%)	62±2 (69±2%)	27±2 (31±2%)	1	15
All	All	3560/3920 (91%)	2474 (69%)	1086 (31%)	1	16

5 of 108 unique residues with a non-rotameric sidechain are listed below. They are sorted by the frequency of occurrence in the ensemble.

Mol	Chain	Res	Type	Models (Total)
1	A	18	LEU	20
1	A	46	ASN	20
1	A	83	SER	20
1	B	18	LEU	20
1	B	22	GLN	20

### 6.3.3 RNA [i](#)

There are no RNA molecules in this entry.

### 6.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

### 6.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

### 6.6 Ligand geometry [i](#)

There are no ligands in this entry.

### 6.7 Other polymers [i](#)

There are no such molecules in this entry.

### 6.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

## 7 Chemical shift validation

No chemical shift data were provided