



wwPDB X-ray Structure Validation Summary Report ⓘ

Mar 12, 2026 – 12:32 PM UTC

PDB ID : 2BCN / pdb_00002bcn
Title : Solvent isotope effects on interfacial protein electron transfer between cytochrome c and cytochrome c peroxidase
Authors : Kang, S.A.; Crane, B.R.
Deposited on : 2005-10-19
Resolution : 1.70 Å (reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:


MolProbity : 4-5-2 with Phenix2.0
Mogul : 2022.3.0, CSD as543be (2022)
Xtrriage (Phenix) : 2.0
EDS : 3.0
Buster-report : wwPDB partial adaption of 1.1.7 (2018)
Percentile statistics : 20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4 : 9.0.010 (Gargrove)
Density-Fitness : 1.0.12
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.49

1 Overall quality at a glance

The following experimental techniques were used to determine the structure:
X-RAY DIFFRACTION

The reported resolution of this entry is 1.70 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	5551 (1.70-1.70)
Clashscore	190562	5924 (1.70-1.70)
Ramachandran outliers	187476	5846 (1.70-1.70)
Sidechain outliers	187428	5846 (1.70-1.70)
RSRZ outliers	180081	5554 (1.70-1.70)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	296	 14% (red), 65% (green), 22% (yellow), 9% (orange), .. (grey)
1	C	296	 18% (red), 66% (green), 22% (yellow), 9% (orange), .. (grey)
2	B	108	 81% (red), 34% (green), 27% (yellow), 33% (orange), 6% (grey)

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
3	ZNH	A	295	X	-	-	-
3	ZNH	C	295	X	-	-	-

2 Entry composition i

There are 5 unique types of molecules in this entry. The entry contains 6265 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called cytochrome c peroxidase.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	294	2371	1514	395	456	6	0	0	0
1	C	294	2371	1514	395	456	6	0	0	0

There are 4 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	-1	MET	-	cloning artifact	GB 171177
A	0	ILE	-	cloning artifact	GB 171177
C	-1	MET	-	cloning artifact	GB 171177
C	0	ILE	-	cloning artifact	GB 171177

- Molecule 2 is a protein called Cytochrome c iso-1.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
2	B	108	847	534	151	158	4	0	0	0

There is a discrepancy between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
B	107	SER	CYS	engineered mutation	UNP P00044


- Molecule 3 is PROTOPORPHYRIN IX CONTAINING ZN (CCD ID: ZNH) (formula: $C_{34}H_{32}N_4O_4Zn$).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
5	A	286	Total 286	O 286	0	0
5	B	27	Total 27	O 27	0	0
5	C	234	Total 234	O 234	0	0


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: cytochrome c peroxidase



- Molecule 1: cytochrome c peroxidase



- Molecule 2: Cytochrome c iso-1



V62	L63	W64	D65	E66	N67	M68	R69	S70	E71	Y72	L73	T74	N75	P76	K77	K78	Y79	I80	F81	G82	T83	K84	M85	A86	F87	G88	G89	L90	K91	K92	E93	K94	D95	R96	N97	D98	L99	I100	T101	Y102	L103	K104	K105	A106	S107	E108
-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	------	------	------	------	------	------	------	------	------

4 Data and refinement statistics

Property	Value	Source
Space group	P 1	Depositor
Cell constants a, b, c, α , β , γ	44.53Å 47.09Å 84.29Å 102.47° 98.74° 102.16°	Depositor
Resolution (Å)	19.61 – 1.70 19.61 – 1.70	Depositor EDS
% Data completeness (in resolution range)	100.0 (19.61-1.70) 93.1 (19.61-1.70)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	3.00 (at 1.70Å)	Xtrriage
Refinement program	REFMAC	Depositor
R, R_{free}	0.219 , 0.230 0.225 , 0.237	Depositor DCC
R_{free} test set	3335 reflections (5.09%)	wwPDB-VP
Wilson B-factor (Å ²)	19.1	Xtrriage
Anisotropy	0.068	Xtrriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.37 , 48.9	EDS
L-test for twinning ²	$\langle L \rangle = 0.48$, $\langle L^2 \rangle = 0.32$	Xtrriage
Estimated twinning fraction	No twinning to report.	Xtrriage
F_o, F_c correlation	0.92	EDS
Total number of atoms	6265	wwPDB-VP
Average B, all atoms (Å ²)	27.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 6.46% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality i

5.1 Standard geometry i

Bond lengths and bond angles in the following residue types are not validated in this section: HEC, ZNH

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	1.65	13/2438 (0.5%)	1.66	59/3302 (1.8%)
1	C	1.17	5/2438 (0.2%)	1.79	60/3302 (1.8%)
2	B	1.02	4/865 (0.5%)	1.82	50/1156 (4.3%)
All	All	1.38	22/5741 (0.4%)	1.74	169/7760 (2.2%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	A	0	8
1	C	1	6
All	All	1	14

The worst 5 of 22 bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	79	ASP	C-N	36.68	1.74	1.34
1	A	1	THR	C-N	-33.12	0.65	1.33
1	A	11	GLU	C-N	28.84	1.74	1.33
1	A	2	THR	C-N	27.33	1.69	1.33
1	C	79	ASP	C-N	25.20	1.66	1.33

The worst 5 of 169 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	C	279	ASP	O-C-N	-47.62	59.26	122.59
1	A	79	ASP	CA-C-N	-25.54	93.31	119.87
1	A	79	ASP	C-N-CA	-25.54	93.31	119.87

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	C	294	LEU	CB-CA-C	-19.89	72.31	110.10
1	A	1	THR	O-C-N	-17.41	95.15	123.00

All (1) chirality outliers are listed below:

Mol	Chain	Res	Type	Atom
1	C	279	ASP	CA

5 of 14 planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	A	1	THR	Mainchain,Peptide
1	A	136	ASP	Mainchain
1	A	3	PRO	Mainchain,Peptide
1	A	80	PRO	Mainchain
1	A	81	SER	Mainchain

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2371	0	2247	121	4
1	C	2371	0	2252	72	4
2	B	847	0	849	90	3
3	A	43	0	30	0	0
3	C	43	0	30	0	0
4	B	43	0	30	5	0
5	A	286	0	0	66	2
5	B	27	0	0	12	1
5	C	234	0	0	37	2
All	All	6265	0	5438	280	9

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 25.

The worst 5 of 280 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:2:THR:C	1:A:3:PRO:N	1.69	1.49
1:A:11:GLU:C	1:A:12:LYS:N	1.74	1.42
1:A:79:ASP:C	1:A:80:PRO:N	1.74	1.41
2:B:100:ILE:O	2:B:104:LYS:HG3	1.22	1.34
1:C:76:GLU:HB3	5:C:1082:HOH:O	1.17	1.34

The worst 5 of 9 symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:226:LYS:NZ	5:A:1233:HOH:O[1_445]	0.39	1.81
5:C:1321:HOH:O	5:C:1332:HOH:O[1_655]	0.68	1.52
1:A:226:LYS:CE	5:A:1233:HOH:O[1_445]	1.14	1.06
1:A:1:THR:OG1	5:C:1472:HOH:O[1_564]	1.86	0.34
1:A:17:GLU:OE1	2:B:49:GLU:OE1[1_665]	1.87	0.33

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	292/296 (99%)	281 (96%)	5 (2%)	6 (2%)	5	1
1	C	292/296 (99%)	284 (97%)	6 (2%)	2 (1%)	18	7
2	B	106/108 (98%)	97 (92%)	6 (6%)	3 (3%)	4	0
All	All	690/700 (99%)	662 (96%)	17 (2%)	11 (2%)	7	1

5 of 11 Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	4	LEU
1	C	279	ASP
1	C	280	ALA
1	A	2	THR

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
2	B	66	GLU

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	252/254 (99%)	232 (92%)	20 (8%)	11	2
1	C	252/254 (99%)	228 (90%)	24 (10%)	8	1
2	B	89/89 (100%)	69 (78%)	20 (22%)	1	0
All	All	593/597 (99%)	529 (89%)	64 (11%)	6	1

5 of 64 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	C	226	LYS
1	C	260	LYS
2	B	24	THR
2	B	21	GLN
1	C	269	LEU

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 34 such sidechains are listed below:

Mol	Chain	Res	Type
1	C	159	GLN
1	C	195	ASN
1	C	222	GLN
1	A	222	GLN
1	A	208	ASN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

3 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
4	HEC	B	109	2	46,50,50	1.29	3 (6%)	58,82,82	0.97	1 (1%)
3	ZNH	C	295	1	50,50,50	1.52	10 (20%)	65,82,82	1.44	9 (13%)
3	ZNH	A	295	1	50,50,50	1.09	2 (4%)	65,82,82	1.69	12 (18%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
4	HEC	B	109	2	-	5/14/54/54	-
3	ZNH	C	295	1	1/1/3/9	2/14/54/54	-
3	ZNH	A	295	1	1/1/3/9	4/14/54/54	-

The worst 5 of 15 bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
3	C	295	ZNH	C1A-C2A	-3.70	1.38	1.45
3	C	295	ZNH	CMC-C2C	3.48	1.57	1.50
4	B	109	HEC	CBB-CAB	-3.19	1.37	1.49

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
3	C	295	ZNH	C4A-C3A	-2.90	1.39	1.45
4	B	109	HEC	CBC-CAC	-2.90	1.38	1.49

The worst 5 of 22 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	A	295	ZNH	C4B-C3B-C2B	-4.86	102.82	107.28
3	A	295	ZNH	CAD-C3D-C4D	4.85	133.16	124.70
3	A	295	ZNH	CAD-C3D-C2D	-4.34	119.74	127.87
3	C	295	ZNH	C4B-C3B-C2B	-4.33	103.30	107.28
3	A	295	ZNH	CBD-CAD-C3D	4.19	124.11	112.53

All (2) chirality outliers are listed below:

Mol	Chain	Res	Type	Atom
3	A	295	ZNH	NA
3	C	295	ZNH	NA

5 of 11 torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
4	B	109	HEC	C4C-C3C-CAC-CBC
3	A	295	ZNH	C4D-C3D-CAD-CBD
3	A	295	ZNH	C2D-C3D-CAD-CBD
4	B	109	HEC	C2C-C3C-CAC-CBC
3	C	295	ZNH	CAA-CBA-CGA-O2A


There are no ring outliers.


1 monomer is involved in 5 short contacts:


Mol	Chain	Res	Type	Clashes	Symm-Clashes
4	B	109	HEC	5	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and

any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.







5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

The following chains have linkage breaks:

Mol	Chain	Number of breaks
1	A	7
1	C	2
2	B	1

The worst 5 of 10 chain breaks are listed below:

Model	Chain	Residue-1	Atom-1	Residue-2	Atom-2	Distance (Å)
1	A	11:GLU	C	12:LYS	N	1.74
1	A	79:ASP	C	80:PRO	N	1.74
1	A	2:THR	C	3:PRO	N	1.69
1	C	79:ASP	C	80:PRO	N	1.66
1	C	277:PRO	C	278:LYS	N	1.65

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	294/296 (99%)	1.05	42 (14%) 6 5	20, 24, 36, 50	0
1	C	294/296 (99%)	1.17	52 (17%) 4 3	18, 24, 39, 49	0
2	B	108/108 (100%)	2.98	87 (80%) 0 0	18, 34, 49, 50	0
All	All	696/700 (99%)	1.40	181 (26%) 1 1	18, 25, 43, 50	0

The worst 5 of 181 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	2	THR	7.6
2	B	55	ASP	6.8
2	B	3	PHE	6.7
1	C	138	THR	6.2
2	B	58	ILE	6.1

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)


There are no oligosaccharides in this entry.


6.4 Ligands [i](#)


In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled ‘Q < 0.9’ lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(\AA^2)	Q<0.9
4	HEC	B	109	43/43	0.83	0.13	17,20,21,23	0
3	ZNH	C	295	43/43	0.97	0.09	19,21,23,26	0
3	ZNH	A	295	43/43	0.98	0.08	19,22,26,28	0

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.







6.5 Other polymers [i](#)

There are no such residues in this entry.