



Full wwPDB X-ray Structure Validation Report ⓘ

Mar 5, 2026 – 03:12 PM UTC

PDB ID : 2BCO / pdb_00002bco
Title : X-ray structure of succinylglutamate desuccinalase from *Vibrio Parahaemolyticus* (RIMD 2210633) at the resolution 2.3 Å, Northeast Structural Genomics Target Vpr14
Authors : Kuzin, A.P.; Abashidze, M.; Forouhar, F.; Benach, J.; Zhou, W.; Acton, T.; Northeast Structural Genomics Consortium (NESG)
Deposited on : 2005-10-19
Resolution : 2.33 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4-5-2 with Phenix2.0
Mogul	:	2022.3.0, CSD as543be (2022)
Xtriage (Phenix)	:	2.0
EDS	:	3.0
Percentile statistics	:	20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4	:	9.0.010 (Gargrove)
Density-Fitness	:	1.0.12
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.33 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	3031 (2.36-2.32)
Clashscore	190562	3127 (2.36-2.32)
Ramachandran outliers	187476	3095 (2.36-2.32)
Sidechain outliers	187428	3095 (2.36-2.32)
RSRZ outliers	180081	3033 (2.36-2.32)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	350	<div> <div>9%</div> <div>72%</div> <div>22%</div> <div>.</div> </div>
1	B	350	<div> <div>23%</div> <div>61%</div> <div>21%</div> <div>.</div> <div>17%</div> </div>

2 Entry composition

There are 3 unique types of molecules in this entry. The entry contains 5393 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Succinylglutamate desuccinylase.

Mol	Chain	Residues	Atoms						ZeroOcc	AltConf	Trace
1	A	338	Total	C	N	O	S	Se	0	0	0
			2703	1710	470	513	5	5			
1	B	292	Total	C	N	O	S	Se	0	0	0
			2341	1485	404	443	5	4			

There are 28 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	1	MSE	MET	modified residue	UNP Q87Q40
A	70	MSE	MET	modified residue	UNP Q87Q40
A	218	MSE	MET	modified residue	UNP Q87Q40
A	276	MSE	MET	modified residue	UNP Q87Q40
A	302	MSE	MET	modified residue	UNP Q87Q40
A	326	MSE	MET	modified residue	UNP Q87Q40
A	343	LEU	-	expression tag	UNP Q87Q40
A	344	GLU	-	expression tag	UNP Q87Q40
A	345	HIS	-	expression tag	UNP Q87Q40
A	346	HIS	-	expression tag	UNP Q87Q40
A	347	HIS	-	expression tag	UNP Q87Q40
A	348	HIS	-	expression tag	UNP Q87Q40
A	349	HIS	-	expression tag	UNP Q87Q40
A	350	HIS	-	expression tag	UNP Q87Q40
B	1	MSE	MET	modified residue	UNP Q87Q40
B	70	MSE	MET	modified residue	UNP Q87Q40
B	218	MSE	MET	modified residue	UNP Q87Q40
B	276	MSE	MET	modified residue	UNP Q87Q40
B	302	MSE	MET	modified residue	UNP Q87Q40
B	326	MSE	MET	modified residue	UNP Q87Q40
B	343	LEU	-	expression tag	UNP Q87Q40
B	344	GLU	-	expression tag	UNP Q87Q40
B	345	HIS	-	expression tag	UNP Q87Q40
B	346	HIS	-	expression tag	UNP Q87Q40
B	347	HIS	-	expression tag	UNP Q87Q40

Continued on next page...

Continued from previous page...

Chain	Residue	Modelled	Actual	Comment	Reference
B	348	HIS	-	expression tag	UNP Q87Q40
B	349	HIS	-	expression tag	UNP Q87Q40
B	350	HIS	-	expression tag	UNP Q87Q40


- Molecule 2 is ZINC ION (CCD ID: ZN) (formula: Zn).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
2	A	1	Total 1	Zn 1	0	0
2	B	1	Total 1	Zn 1	0	0

- Molecule 3 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	A	244	Total 244	O 244	0	0
3	B	103	Total 103	O 103	0	0

- Molecule 1: Succinylglutamate desuccinylase



4 Data and refinement statistics

Property	Value	Source
Space group	P 1 21 1	Depositor
Cell constants a, b, c, α , β , γ	62.85Å 72.56Å 93.03Å 90.00° 105.66° 90.00°	Depositor
Resolution (Å)	30.26 – 2.33 30.26 – 2.33	Depositor EDS
% Data completeness (in resolution range)	87.0 (30.26-2.33) 92.0 (30.26-2.33)	Depositor EDS
R_{merge}	0.04	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	9.33 (at 2.34Å)	Xtriage
Refinement program	CNS 1.1	Depositor
R, R_{free}	0.213 , 0.267 0.225 , 0.280	Depositor DCC
R_{free} test set	3070 reflections (4.88%)	wwPDB-VP
Wilson B-factor (Å ²)	23.6	Xtriage
Anisotropy	0.241	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.30 , 47.2	EDS
L-test for twinning ²	$\langle L \rangle = 0.50$, $\langle L^2 \rangle = 0.33$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.92	EDS
Total number of atoms	5393	wwPDB-VP
Average B, all atoms (Å ²)	38.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 5.16% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: ZN

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.42	0/2755	0.91	15/3725 (0.4%)
1	B	0.34	0/2381	0.88	7/3216 (0.2%)
All	All	0.39	0/5136	0.90	22/6941 (0.3%)

There are no bond length outliers.

All (22) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	B	96	ALA	N-CA-C	11.36	123.75	111.36
1	A	67	THR	N-CA-C	8.43	122.86	112.23
1	B	67	THR	N-CA-C	8.41	121.58	111.82
1	A	217	THR	N-CA-C	-7.15	96.87	108.52
1	A	231	LEU	N-CA-C	6.73	119.47	111.33
1	B	95	ILE	N-CA-C	-6.68	97.36	106.85
1	A	64	GLY	N-CA-C	6.63	120.92	112.83
1	A	178	SER	N-CA-C	6.58	119.02	110.53
1	B	56	ILE	N-CA-C	6.49	117.21	107.80
1	A	149	ARG	N-CA-C	6.41	119.95	109.76
1	A	95	ILE	N-CA-C	-5.88	98.24	106.53
1	A	154	LEU	N-CA-C	5.52	117.74	108.96
1	A	191	GLU	N-CA-C	5.40	117.92	111.71
1	B	149	ARG	N-CA-C	5.39	117.50	109.25
1	A	215	ALA	N-CA-C	5.28	118.35	109.95
1	A	115	LEU	N-CA-C	5.20	118.65	112.72
1	A	56	ILE	N-CA-C	5.20	115.34	107.80
1	A	62	ILE	N-CA-C	-5.19	104.57	111.05
1	B	191	GLU	N-CA-C	5.09	117.49	111.33
1	A	162	LYS	N-CA-C	-5.09	106.91	113.01
1	A	97	HIS	N-CA-C	5.07	121.02	109.81
1	B	154	LEU	N-CA-C	5.03	116.95	109.25

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2703	0	2654	48	0
1	B	2341	0	2321	50	0
2	A	1	0	0	0	0
2	B	1	0	0	0	0
3	A	244	0	0	3	0
3	B	103	0	0	0	0
All	All	5393	0	4975	96	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 10.

All (96) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:66:GLU:HG3	1:B:156:CYS:HB2	1.35	1.03
1:B:152:LEU:HB3	1:B:218:MSE:HE1	1.50	0.93
1:B:262:VAL:HG12	1:B:328:CYS:HB3	1.60	0.81
1:B:33:GLN:HB2	1:B:45:ILE:HB	1.69	0.74
1:B:238:ASP:O	1:B:242:ARG:HG2	1.90	0.71
1:A:249:GLN:HE22	1:A:251:GLU:HB2	1.57	0.69
1:B:70:MSE:HE3	1:B:95:ILE:HG23	1.76	0.68
1:A:66:GLU:HG2	1:A:156:CYS:HB2	1.75	0.67
1:B:257:CYS:O	1:B:259:LYS:HD2	1.94	0.67
1:B:68:ALA:HB3	1:B:69:PRO:HD3	1.76	0.66
1:B:333:ARG:O	1:B:340:VAL:HG22	1.97	0.64
1:B:163:HIS:HE1	1:B:258:ILE:HD12	1.64	0.63
1:A:269:LEU:O	1:A:319:ILE:HD11	2.00	0.62
1:B:66:GLU:HG3	1:B:156:CYS:CB	2.22	0.61
1:A:166:PHE:CZ	1:A:219:GLU:HG3	2.36	0.60
1:A:185:LEU:HG	1:A:193:VAL:HG21	1.82	0.60
1:B:159:ARG:NH1	1:B:313:PRO:HG2	2.16	0.60

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:110:GLU:O	1:B:126:GLU:HG2	2.03	0.59
1:B:152:LEU:CB	1:B:218:MSE:HE1	2.28	0.59
1:A:330:VAL:HG13	1:A:332:THR:HG23	1.85	0.58
1:A:125:LYS:O	1:A:129:ILE:HG12	2.04	0.56
1:A:68:ALA:HB3	1:A:69:PRO:HD3	1.87	0.56
1:A:241:LEU:O	1:A:245:ILE:HG23	2.06	0.56
1:A:153:ASP:CG	1:A:155:HIS:HE2	2.14	0.56
1:A:248:ALA:HB2	3:A:640:HOH:O	2.04	0.55
1:B:163:HIS:CE1	1:B:258:ILE:HD12	2.42	0.54
1:A:317:VAL:CG1	1:A:321:GLN:HB2	2.39	0.53
1:A:341:TYR:O	1:A:342:ASP:HB2	2.09	0.53
1:B:55:ASN:OD1	1:B:90:ARG:HD3	2.10	0.52
1:B:261:ARG:CZ	1:B:341:TYR:HB3	2.40	0.52
1:A:330:VAL:CG2	1:A:341:TYR:HB2	2.40	0.51
1:A:317:VAL:HG13	1:A:321:GLN:HB2	1.94	0.50
1:A:188:ALA:O	1:A:189:HIS:HB2	2.12	0.50
1:B:335:GLU:HG2	1:B:336:GLU:HG2	1.94	0.49
1:B:233:ARG:C	1:B:234:LEU:HD12	2.38	0.49
1:A:229:ASN:HB2	1:A:231:LEU:HD13	1.93	0.49
1:A:28:LEU:HD12	1:A:32:VAL:HB	1.95	0.49
1:A:330:VAL:HG21	1:A:341:TYR:HB2	1.95	0.48
1:B:314:ASN:ND2	1:B:316:HIS:H	2.11	0.48
1:B:262:VAL:CG1	1:B:328:CYS:HB3	2.39	0.48
1:A:159:ARG:HD3	1:A:281:VAL:O	2.14	0.48
1:A:312:PHE:N	1:A:313:PRO:HD3	2.28	0.48
1:B:5:LEU:O	1:B:5:LEU:HD13	2.14	0.47
1:B:161:SER:HA	1:B:282:GLU:HB3	1.96	0.47
1:B:17:VAL:HG23	1:B:18:HIS:ND1	2.29	0.47
1:B:47:GLU:C	1:B:49:PRO:HD3	2.39	0.47
1:A:114:ARG:HA	1:A:201:SER:OG	2.13	0.47
1:A:322:ARG:HG3	1:A:322:ARG:HH11	1.80	0.47
1:B:30:ASN:HB2	1:B:32:VAL:HG23	1.97	0.47
1:A:330:VAL:HG13	1:A:332:THR:CG2	2.44	0.47
1:B:47:GLU:O	1:B:49:PRO:HD3	2.15	0.46
1:B:161:SER:HB3	1:B:283:ASN:HB2	1.97	0.46
1:B:222:ARG:HH11	1:B:222:ARG:HG3	1.81	0.46
1:B:14:THR:HG21	1:B:95:ILE:HG21	1.97	0.46
1:B:42:LEU:HD23	1:B:77:ILE:HD12	1.96	0.46
1:B:277:PHE:HB2	1:B:281:VAL:HG11	1.97	0.46
1:A:48:ASN:HB2	3:A:691:HOH:O	2.15	0.45
1:A:73:VAL:O	1:A:77:ILE:HG12	2.17	0.45

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:106:ARG:HG3	3:A:518:HOH:O	2.16	0.45
1:A:5:LEU:HD13	1:A:26:GLN:HB2	1.98	0.45
1:B:253:LEU:N	1:B:253:LEU:HD12	2.32	0.45
1:A:193:VAL:HG23	1:A:257:CYS:SG	2.57	0.44
1:A:314:ASN:HB3	1:A:317:VAL:HG23	1.99	0.44
1:A:110:GLU:O	1:A:126:GLU:HG2	2.18	0.44
1:B:92:LEU:HD13	1:B:94:ILE:HD11	1.99	0.44
1:B:193:VAL:O	1:B:193:VAL:HG13	2.18	0.44
1:A:70:MSE:HE1	1:A:101:THR:HG21	2.00	0.44
1:B:192:ALA:HB1	1:B:260:TYR:HD2	1.82	0.44
1:B:197:ASN:H	1:B:197:ASN:ND2	2.16	0.43
1:B:197:ASN:H	1:B:197:ASN:HD22	1.66	0.43
1:A:261:ARG:HD3	1:A:330:VAL:HG21	2.01	0.43
1:B:66:GLU:HG2	1:B:156:CYS:H	1.82	0.43
1:B:197:ASN:CG	1:B:263:SER:HB2	2.43	0.43
1:A:62:ILE:HD13	1:A:62:ILE:HA	1.92	0.43
1:B:11:LEU:CD1	1:B:70:MSE:HE2	2.49	0.43
1:B:220:LEU:HD21	1:B:237:PHE:CE2	2.54	0.42
1:A:224:ALA:HB3	1:A:229:ASN:OD1	2.19	0.42
1:A:130:ALA:O	1:A:134:LYS:HG3	2.19	0.42
1:A:311:VAL:HG12	1:A:312:PHE:CD1	2.54	0.42
1:B:69:PRO:HD3	1:B:156:CYS:SG	2.60	0.42
1:A:26:GLN:HE21	1:A:28:LEU:HD23	1.84	0.41
1:A:325:LEU:HD12	1:A:325:LEU:HA	1.92	0.41
1:A:261:ARG:HD3	1:A:330:VAL:CG2	2.51	0.41
1:A:67:THR:HB	1:A:226:ILE:HG12	2.02	0.41
1:A:264:ARG:NH2	1:A:302:MSE:O	2.54	0.41
1:B:13:ASP:HA	1:B:19:ILE:HD12	2.01	0.41
1:B:117:ASP:HB3	1:B:201:SER:HB3	2.03	0.41
1:A:135:LEU:HD21	1:B:125:LYS:HA	2.01	0.41
1:A:162:LYS:HE2	1:A:284:PHE:HB3	2.02	0.41
1:B:72:LEU:O	1:B:76:ILE:HG13	2.21	0.41
1:B:220:LEU:HD21	1:B:237:PHE:CZ	2.56	0.41
1:A:22:ALA:HB2	1:B:18:HIS:HA	2.03	0.41
1:A:310:ILE:HG12	1:A:313:PRO:HG3	2.03	0.40
1:A:70:MSE:CE	1:A:226:ILE:HD11	2.51	0.40
1:B:88:ASP:OD2	1:B:247:GLU:HB3	2.22	0.40
1:A:291:GLU:O	1:A:302:MSE:HA	2.21	0.40

There are no symmetry-related clashes.

5.3 Torsion angles

5.3.1 Protein backbone

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	334/350 (95%)	325 (97%)	9 (3%)	0	100	100
1	B	280/350 (80%)	247 (88%)	30 (11%)	3 (1%)	11	10
All	All	614/700 (88%)	572 (93%)	39 (6%)	3 (0%)	24	26

All (3) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	B	341	TYR
1	B	264	ARG
1	B	259	LYS

5.3.2 Protein sidechains

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	298/304 (98%)	286 (96%)	12 (4%)	28	36
1	B	261/304 (86%)	254 (97%)	7 (3%)	39	51
All	All	559/608 (92%)	540 (97%)	19 (3%)	32	42

All (19) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	15	LEU
1	A	42	LEU
1	A	97	HIS

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	A	112	LEU
1	A	127	LEU
1	A	131	ASP
1	A	135	LEU
1	A	194	LEU
1	A	220	LEU
1	A	231	LEU
1	A	245	ILE
1	A	325	LEU
1	B	5	LEU
1	B	28	LEU
1	B	53	THR
1	B	92	LEU
1	B	138	ARG
1	B	197	ASN
1	B	341	TYR

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (9) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	26	GLN
1	A	33	GLN
1	A	189	HIS
1	A	249	GLN
1	A	305	ASN
1	A	316	HIS
1	B	8	GLN
1	B	197	ASN
1	B	314	ASN

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

Of 2 ligands modelled in this entry, 2 are monoatomic - leaving 0 for Mogul analysis.

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	333/350 (95%)	0.21	32 (9%) 13 16	6, 20, 65, 114	0
1	B	288/350 (82%)	1.37	82 (28%) 1 1	13, 45, 98, 138	0
All	All	621/700 (88%)	0.75	114 (18%) 3 4	6, 30, 93, 138	0

All (114) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	B	263	SER	6.5
1	B	330	VAL	6.1
1	B	341	TYR	5.9
1	B	21	VAL	5.8
1	B	340	VAL	5.7
1	B	342	ASP	5.5
1	B	325	LEU	5.2
1	B	262	VAL	5.1
1	B	327	VAL	4.8
1	A	253	LEU	4.8
1	A	252	HIS	4.8
1	B	277	PHE	4.4
1	B	333	ARG	4.3
1	B	328	CYS	4.3
1	B	246	ALA	4.2
1	B	261	ARG	4.1
1	A	342	ASP	4.1
1	B	337	GLY	4.1
1	B	339	LEU	4.1
1	B	332	THR	4.0
1	B	317	VAL	4.0
1	B	50	THR	3.8
1	B	309	ALA	3.8
1	B	338	GLU	3.8

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	B	253	LEU	3.8
1	B	264	ARG	3.7
1	A	19	ILE	3.7
1	B	314	ASN	3.7
1	B	87	VAL	3.6
1	B	51	GLN	3.6
1	B	19	ILE	3.6
1	B	17	VAL	3.5
1	B	4	SER	3.4
1	B	260	TYR	3.4
1	A	305	ASN	3.3
1	B	279	ASP	3.3
1	B	280	ASN	3.2
1	A	251	GLU	3.2
1	B	30	ASN	3.2
1	A	319	ILE	3.2
1	B	18	HIS	3.1
1	B	265	THR	3.1
1	B	254	SER	3.1
1	A	21	VAL	3.1
1	B	310	ILE	3.0
1	B	189	HIS	2.9
1	A	18	HIS	2.9
1	B	255	LYS	2.9
1	B	281	VAL	2.9
1	A	317	VAL	2.9
1	B	27	VAL	2.9
1	A	304	LYS	2.8
1	B	8	GLN	2.8
1	B	334	PHE	2.8
1	B	329	GLU	2.8
1	A	48	ASN	2.8
1	B	283	ASN	2.8
1	B	9	SER	2.8
1	B	207	SER	2.8
1	B	120	GLU	2.8
1	B	83	GLY	2.7
1	B	178	SER	2.7
1	B	258	ILE	2.7
1	B	188	ALA	2.7
1	A	26	GLN	2.7
1	B	284	PHE	2.7

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	B	86	LYS	2.7
1	B	162	LYS	2.7
1	A	330	VAL	2.6
1	B	158	ILE	2.6
1	A	289	HIS	2.6
1	A	299	LYS	2.6
1	A	7	ARG	2.6
1	B	312	PHE	2.6
1	B	81	GLU	2.6
1	A	307	ASN	2.5
1	A	288	VAL	2.5
1	B	82	SER	2.5
1	A	306	ASP	2.5
1	B	33	GLN	2.5
1	B	221	GLY	2.5
1	A	272	ASP	2.4
1	B	84	PHE	2.4
1	B	331	LYS	2.4
1	B	256	PRO	2.4
1	A	273	PHE	2.3
1	B	232	ASP	2.3
1	B	236	ALA	2.3
1	B	316	HIS	2.3
1	B	52	GLU	2.3
1	B	159	ARG	2.3
1	A	249	GLN	2.3
1	B	194	LEU	2.3
1	B	29	SER	2.3
1	B	37	TYR	2.2
1	A	300	PRO	2.2
1	A	248	ALA	2.2
1	B	180	ALA	2.2
1	B	227	GLY	2.2
1	A	29	SER	2.2
1	A	250	PRO	2.2
1	B	335	GLU	2.1
1	A	270	HIS	2.1
1	B	198	SER	2.1
1	B	164	TYR	2.1
1	A	269	LEU	2.1
1	A	8	GLN	2.1
1	A	4	SER	2.1

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	B	161	SER	2.1
1	B	257	CYS	2.1
1	B	282	GLU	2.0
1	B	142	GLN	2.0
1	B	278	ASP	2.0
1	A	51	GLN	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(\AA^2)	Q<0.9
2	ZN	B	502	1/1	0.93	0.08	62,62,62,62	0
2	ZN	A	501	1/1	0.98	0.08	46,46,46,46	0

6.5 Other polymers [i](#)

There are no such residues in this entry.