



# wwPDB NMR Structure Validation Summary Report ⓘ

Mar 6, 2026 – 06:46 PM UTC

PDB ID : 7BLM / pdb\_00007blm  
BMRB ID : 34592  
Title : AG repetition attached to a compact i-motif clip at 3'-end  
Authors : Novotny, A.; Novotny, J.  
Deposited on : 2021-01-18

This is a wwPDB NMR Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<https://www.wwpdb.org/validation/2017/NMRValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<https://www.wwpdb.org/validation/2017/FAQs#types>.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4-5-2 with Phenix2.0  
Mogul : 2022.3.0, CSD as543be (2022)  
Percentile statistics : 20250101.v01 (using entries in the PDB archive January 1st 2025)  
wwPDB-RCI : v\_1n\_11\_5\_13\_A (Berjanski et al., 2005)  
PANAV : Wang et al. (2010)  
wwPDB-ShiftChecker : v1.2  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : 2.49

# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

*SOLUTION NMR*





The overall completeness of chemical shifts assignment is 7%.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	NMR archive (#Entries)
Clashscore	229148	14424

The table below summarises the geometric issues observed across the polymeric chains and their fit to the experimental data. The red, orange, yellow and green segments indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria. A cyan segment indicates the fraction of residues that are not part of the well-defined cores, and a grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$

Mol	Chain	Length	Quality of chain
1	A	7	 43% 43% 14%
1	B	7	 57% 29% 14%
2	C	7	 71% 29%
2	D	7	 57% 43%

## 2 Ensemble composition and analysis

This entry contains 4 models. This entry does not contain polypeptide chains, therefore identification of well-defined residues and clustering analysis are not possible. All residues are included in the validation scores.

### 3 Entry composition [i](#)

There are 2 unique types of molecules in this entry. The entry contains 882 atoms, of which 322 are hydrogens and 0 are deuteriums.

- Molecule 1 is a DNA chain called AGAG(CH<sup>+</sup>)C(CH<sup>+</sup>).

Mol	Chain	Residues	Atoms					Trace	
			Total	C	H	N	O		P
1	A	7	221	67	81	29	38	6	0
1	B	7	221	67	81	29	38	6	0

- Molecule 2 is a DNA chain called AGAGC(CH<sup>+</sup>)C.


Mol	Chain	Residues	Atoms					Trace	
			Total	C	H	N	O		P
2	C	7	220	67	80	29	38	6	0
2	D	7	220	67	80	29	38	6	0

## 4 Residue-property plots

### 4.1 Average score per residue in the NMR ensemble

These plots are provided for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic is the same as shown in the summary in section 1 of this report. The second graphic shows the sequence where residues are colour-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. Stretches of 2 or more consecutive residues without any outliers are shown as green connectors. Residues which are classified as ill-defined in the NMR ensemble, are shown in cyan with an underline colour-coded according to the previous scheme. Residues which were present in the experimental sample, but not modelled in the final structure are shown in grey.


- Molecule 1: AGAG(CH+)C(CH+)



- Molecule 1: AGAG(CH+)C(CH+)



- Molecule 2: AGAGC(CH+)C




- Molecule 2: AGAGC(CH+)C



### 4.2 Residue scores for the representative (author defined) model from the NMR ensemble

The representative model is number 1. Colouring as in section 4.1 above.

- Molecule 1: AGAG(CH+)C(CH+)

Chain A:  14% 71% 14%

 A1  
G2  
A3  
G4  
C5  
C6  
C7

• Molecule 1: AGAG(CH+)C(CH+)

Chain B:  43% 43% 14%

 A8  
G11  
C12  
C13  
C14


• Molecule 2: AGAGC(CH+)C

Chain C:  57% 43%

 A15  
G16  
A17  
G18  
C19  
C20  
C21

• Molecule 2: AGAGC(CH+)C

Chain D:  57% 43%

 A22  
G25  
C26  
C27  
C28

## 5 Refinement protocol and experimental data overview

The models were refined using the following method: *molecular dynamics*.

Of the 3000 calculated structures, 4 were deposited, based on the following criterion: *Snapshot every 100 nanoseconds taken from initial 300 ns of 1 microsecond unrestrained MD.*

The following table shows the software used for structure solution, optimisation and refinement.

Software name	Classification	Version
Amber	refinement	16
Amber	structure calculation	16

The following table shows chemical shift validation statistics as aggregates over all chemical shift files. Detailed validation can be found in section 7 of this report.

Chemical shift file(s)	working_cs.cif
Number of chemical shift lists	1
Total number of shifts	49
Number of shifts mapped to atoms	49
Number of unparsed shifts	0
Number of shifts with mapping errors	0
Number of shifts with mapping warnings	0
Assignment completeness (well-defined parts)	7%

## 6 Model quality i

### 6.1 Standard geometry i

Bond lengths and bond angles in the following residue types are not validated in this section: DNR

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the (average) root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	#Z>5	RMSZ	#Z>5
1	A	0.99±0.01	1±0/114 ( 0.7± 0.4%)	1.33±0.02	0±0/172 ( 0.3± 0.3%)
1	B	1.00±0.01	1±0/114 ( 0.9± 0.0%)	1.36±0.01	0±0/172 ( 0.0± 0.0%)
2	C	1.10±0.01	0±0/135 ( 0.2± 0.3%)	1.32±0.03	0±0/204 ( 0.1± 0.2%)
2	D	1.10±0.01	1±0/135 ( 0.6± 0.3%)	1.30±0.02	0±0/204 ( 0.1± 0.2%)
All	All	1.06	11/1992 ( 0.6%)	1.33	4/3008 ( 0.1%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	Chirality	Planarity
1	A	0.0±0.0	2.0±0.7
1	B	0.0±0.0	1.5±0.5
2	C	0.0±0.0	1.5±0.5
2	D	0.0±0.0	1.5±0.5
All	All	0	26

All unique bond outliers are listed below. They are sorted according to the Z-score of the worst occurrence in the ensemble.

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)	Models	
								Worst	Total
1	A	6	DC	C5-C6	5.20	1.44	1.34	3	3
1	B	13	DC	C5-C6	5.18	1.44	1.34	4	4
2	C	19	DC	C5-C6	5.02	1.44	1.34	2	1
2	D	26	DC	C5-C6	5.02	1.44	1.34	3	3

All unique angle outliers are listed below. They are sorted according to the Z-score of the worst occurrence in the ensemble.

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)	Models	
								Worst	Total
2	D	23	DG	O3'-P-O5'	-5.89	95.16	104.00	4	1
2	C	16	DG	O3'-P-O5'	-5.83	95.25	104.00	4	1
1	A	2	DG	O3'-P-O5'	-5.41	95.89	104.00	2	2

There are no chirality outliers.

5 of 13 unique planar outliers are listed below. They are sorted by the frequency of occurrence in the ensemble.

Mol	Chain	Res	Type	Group	Models (Total)
1	A	6	DC	Sidechain	4
2	D	25	DG	Sidechain	4
1	A	4	DG	Sidechain	3
1	B	13	DC	Sidechain	3
2	C	16	DG	Sidechain	3

## 6.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in each chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes averaged over the ensemble.

Mol	Chain	Non-H	H(model)	H(added)	Clashes
All	All	2240	1288	1288	-

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is -.

There are no clashes.

## 6.3 Torsion angles [i](#)

### 6.3.1 Protein backbone [i](#)

There are no protein molecules in this entry.

### 6.3.2 Protein sidechains [i](#)

There are no protein molecules in this entry.

### 6.3.3 RNA [i](#)

There are no RNA molecules in this entry.

## 6.4 Non-standard residues in protein, DNA, RNA chains [i](#)

6 non-standard protein/DNA/RNA residues are modelled in this entry.

In the following table, the Counts columns list the number of bonds for which Mogul statistics could be retrieved, the number of bonds that are observed in the model and the number of bonds that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length is the number of standard deviations the observed value is removed from the expected value. A bond length with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the average root-mean-square of all Z scores of the bond lengths.

Mol	Type	Chain	Res	Link	Bond lengths		
					Counts	RMSZ	#Z>2
1	DNR	B	14	2,1	17,20,21	1.44±0.01	2±0 (11±0%)
1	DNR	A	5	2,1	17,20,21	1.41±0.03	2±0 (11±0%)
2	DNR	D	27	2,1	17,20,21	1.38±0.01	2±0 (11±0%)
2	DNR	C	20	2,1	17,20,21	1.38±0.01	2±0 (11±0%)
1	DNR	B	12	2,1	17,20,21	1.41±0.01	2±0 (11±0%)
1	DNR	A	7	2,1	17,20,21	1.43±0.02	2±0 (11±0%)

In the following table, the Counts columns list the number of angles for which Mogul statistics could be retrieved, the number of angles that are observed in the model and the number of angles that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond angle is the number of standard deviations the observed value is removed from the expected value. A bond angle with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the average root-mean-square of all Z scores of the bond angles.

Mol	Type	Chain	Res	Link	Bond angles		
					Counts	RMSZ	#Z>2
1	DNR	B	14	2,1	23,28,31	1.09±0.02	2±1 (6±4%)
1	DNR	A	5	2,1	23,28,31	1.06±0.02	2±1 (7±3%)
2	DNR	D	27	2,1	23,28,31	1.09±0.01	0±0 (0±0%)
2	DNR	C	20	2,1	23,28,31	1.09±0.02	1±1 (3±3%)
1	DNR	B	12	2,1	23,28,31	1.19±0.04	1±0 (5±1%)
1	DNR	A	7	2,1	23,28,31	1.12±0.08	2±1 (8±6%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the chemical component dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	DNR	D	27	2,1	-	0±0,7,21,22	0±0,2,2,2
1	DNR	B	14	2,1	-	0±0,7,21,22	0±0,2,2,2
2	DNR	C	20	2,1	-	0±0,7,21,22	0±0,2,2,2
1	DNR	B	12	2,1	-	0±0,7,21,22	0±0,2,2,2
1	DNR	A	5	2,1	-	0±0,7,21,22	0±0,2,2,2
1	DNR	A	7	2,1	-	0±0,7,21,22	0±0,2,2,2

5 of 12 unique bond outliers are listed below. They are sorted according to the Z-score of the worst occurrence in the ensemble.

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)	Models	
								Worst	Total
1	B	14	DNR	C6-C5	4.22	1.44	1.35	3	4
1	A	5	DNR	C6-C5	4.15	1.44	1.35	4	4
1	B	12	DNR	C6-C5	4.12	1.44	1.35	2	4
1	A	7	DNR	C6-C5	4.10	1.44	1.35	2	4
2	C	20	DNR	C6-C5	4.06	1.44	1.35	2	4

5 of 14 unique angle outliers are listed below. They are sorted according to the Z-score of the worst occurrence in the ensemble.

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)	Models	
								Worst	Total
1	B	12	DNR	O4'-C1'-N1	3.47	114.01	107.86	2	4
1	A	7	DNR	O4'-C1'-C2'	2.47	101.62	106.25	1	1
1	A	7	DNR	O2-C2-N3	2.37	118.60	122.33	2	3
1	A	7	DNR	O2-C2-N1	2.29	123.39	118.90	2	3
1	A	5	DNR	O2-C2-N1	2.26	123.33	118.90	4	4

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

## 6.5 Carbohydrates (i)

There are no oligosaccharides in this entry.

## 6.6 Ligand geometry [i](#)

There are no ligands in this entry.

## 6.7 Other polymers [i](#)

There are no such molecules in this entry.

## 6.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

## 7 Chemical shift validation [i](#)

The completeness of assignment taking into account all chemical shift lists is 7% for the well-defined parts and 7% for the entire structure.

### 7.1 Chemical shift list 1

File name: working\_cs.cif

Chemical shift list name: *starch\_output*

#### 7.1.1 Bookkeeping [i](#)

The following table shows the results of parsing the chemical shift list and reports the number of nuclei with statistically unusual chemical shifts.

Total number of shifts	49
Number of shifts mapped to atoms	49
Number of unparsed shifts	0
Number of shifts with mapping errors	0
Number of shifts with mapping warnings	0
Number of shift outliers (ShiftChecker)	1

#### 7.1.2 Chemical shift referencing [i](#)

No chemical shift referencing corrections were calculated (not enough data).

#### 7.1.3 Completeness of resonance assignments [i](#)

The following table shows the completeness of the chemical shift assignments for the well-defined regions of the structure. The overall completeness is 7%, i.e. 31 atoms were assigned a chemical shift out of a possible 448. 0 out of 0 assigned methyl groups (LEU and VAL) were assigned stereospecifically.

	Total	<sup>1</sup> H	<sup>13</sup> C	<sup>15</sup> N
Sugar	15/264 (6%)	15/154 (10%)	0/110 (0%)	0/0 (—%)
Base	16/184 (9%)	16/118 (14%)	0/36 (0%)	0/30 (0%)
Overall	31/448 (7%)	31/272 (11%)	0/146 (0%)	0/30 (0%)

#### 7.1.4 Statistically unusual chemical shifts [i](#)

The following table lists the statistically unusual chemical shifts. These are statistical measures, and large deviations from the mean do not necessarily imply incorrect assignments. Molecules con-

taining paramagnetic centres or hemes are expected to give rise to anomalous chemical shifts.

List Id	Chain	Res	Type	Atom	Shift, <i>ppm</i>	Expected range, <i>ppm</i>	Z-score
1	A	6	DC	H3'	15.57	3.90 – 5.62	62.9

### 7.1.5 Random Coil Index (RCI) plots [i](#)

No *random coil index*(RCI) plot could be generated from the current chemical shift list. RCI is only applicable to proteins