



wwPDB X-ray Structure Validation Summary Report ⓘ

Mar 8, 2026 – 03:39 PM UTC

PDB ID : 5BRM / pdb_00005brm
Title : Structural basis for Mob1-dependent activation of the core Mst-Lats kinase cascade in Hippo signaling
Authors : Luo, X.; Ni, L.
Deposited on : 2015-05-31
Resolution : 2.65 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4-5-2 with Phenix2.0
Mogul	:	2022.3.0, CSD as543be (2022)
Xtriage (Phenix)	:	2.0
EDS	:	3.0
Percentile statistics	:	20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4	:	9.0.010 (Gargrove)
Density-Fitness	:	1.0.12
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.65 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	1110 (2.66-2.66)
Clashscore	190562	1141 (2.66-2.66)
Ramachandran outliers	187476	1126 (2.66-2.66)
Sidechain outliers	187428	1126 (2.66-2.66)
RSRZ outliers	180081	1110 (2.66-2.66)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	177	
1	B	177	
1	C	177	
1	D	177	
1	E	177	

Continued on next page...

Continued from previous page...

Mol	Chain	Length	Quality of chain
1	F	177	
2	G	31	
2	H	31	
2	I	31	
2	J	31	
2	K	31	
2	L	31	
2	M	31	
2	N	31	
2	O	31	

2 Entry composition

There are 4 unique types of molecules in this entry. The entry contains 17464 atoms, of which 8552 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called MOB kinase activator 1A.

Mol	Chain	Residues	Atoms						ZeroOcc	AltConf	Trace
1	A	154	Total	C	H	N	O	S	0	0	0
			2538	842	1247	209	232	8			
1	B	158	Total	C	H	N	O	S	0	0	0
			2586	857	1271	213	237	8			
1	C	160	Total	C	H	N	O	S	0	0	0
			2629	869	1297	216	239	8			
1	D	144	Total	C	H	N	O	S	0	0	0
			2384	796	1166	196	218	8			
1	E	160	Total	C	H	N	O	S	0	0	0
			2628	869	1296	216	239	8			
1	F	160	Total	C	H	N	O	S	0	0	0
			2628	869	1296	216	239	8			

There are 6 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	40	GLY	-	expression tag	UNP Q9H8S9
B	40	GLY	-	expression tag	UNP Q9H8S9
C	40	GLY	-	expression tag	UNP Q9H8S9
D	40	GLY	-	expression tag	UNP Q9H8S9
E	40	GLY	-	expression tag	UNP Q9H8S9
F	40	GLY	-	expression tag	UNP Q9H8S9

- Molecule 2 is a protein called Serine/threonine-protein kinase 3.

Mol	Chain	Residues	Atoms						ZeroOcc	AltConf	Trace
2	G	22	Total	C	H	N	O	P	S	0	0
			350	111	167	32	37	1	2		
2	H	17	Total	C	H	N	O	P	S	0	0
			288	92	137	27	29	1	2		
2	I	12	Total	C	H	N	O	P	S	0	0
			207	69	97	16	22	1	2		

Continued on next page...

Continued from previous page...

Mol	Chain	Residues	Atoms							ZeroOcc	AltConf	Trace
2	J	13	Total	C	H	N	O	P	S	0	0	0
			220	73	102	18	24	1	2			
2	K	11	Total	C	H	N	O	P	S	0	0	0
			183	63	84	12	21	1	2			
2	L	12	Total	C	H	N	O	P	S	0	0	0
			208	69	98	17	21	1	2			
2	M	17	Total	C	H	N	O	S		0	0	0
			277	87	136	27	26	1				
2	N	12	Total	C	H	N	O	S		0	0	0
			193	64	92	17	19	1				
2	O	8	Total	C	H	N	O			0	0	0
			133	43	66	13	11					

- Molecule 3 is ZINC ION (CCD ID: ZN) (formula: Zn).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	A	1	Total	Zn	0	0
			1	1		
3	B	1	Total	Zn	0	0
			1	1		
3	C	1	Total	Zn	0	0
			1	1		
3	D	1	Total	Zn	0	0
			1	1		
3	E	1	Total	Zn	0	0
			1	1		
3	F	1	Total	Zn	0	0
			1	1		

- Molecule 4 is water.


Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	A	1	Total	O	0	0
			1	1		
4	B	1	Total	O	0	0
			1	1		
4	C	1	Total	O	0	0
			1	1		
4	F	3	Total	O	0	0
			3	3		

3 Residue-property plots

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: MOB kinase activator 1A

Chain A: 




- Molecule 1: MOB kinase activator 1A

Chain B: 




- Molecule 1: MOB kinase activator 1A

Chain C: 




- Molecule 1: MOB kinase activator 1A

Chain D: 



- Molecule 1: MOB kinase activator 1A

Chain E: 



- Molecule 1: MOB kinase activator 1A

Chain F:  84% 6% 10%





- Molecule 2: Serine/threonine-protein kinase 3

Chain G:  65% . . 29%




- Molecule 2: Serine/threonine-protein kinase 3

Chain H:  42% 13% 45%




- Molecule 2: Serine/threonine-protein kinase 3

Chain I:  35% . 61%




- Molecule 2: Serine/threonine-protein kinase 3

Chain J:  3% 35% . . 58%




- Molecule 2: Serine/threonine-protein kinase 3

Chain K:  32% . 65%



- Molecule 2: Serine/threonine-protein kinase 3

Chain L:  26% 13% 61%



- Molecule 2: Serine/threonine-protein kinase 3

Chain M: 39% 16% 45%

ASP	GLU	GLU	GLU	GLU	ASP	GLY	TPO	MET	K380	R390	P391	S392	F393	M394	D395	Y396	PHE	ASP	LYS	GLN	ASP
-----	-----	-----	-----	-----	-----	-----	-----	-----	------	------	------	------	------	------	------	------	-----	-----	-----	-----	-----

● Molecule 2: Serine/threonine-protein kinase 3

Chain N: 32% 6% 61%

ASP	GLU	GLU	GLU	GLU	ASP	GLY	TPO	MET	LYS	ARG	ASN	ALA	THR	S385	P386	Y396	PHE	ASP	LYS	GLN	ASP
-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	------	------	------	-----	-----	-----	-----	-----

● Molecule 2: Serine/threonine-protein kinase 3

Chain O: 26% 74%

ASP	GLU	GLU	GLU	GLY	ASP	GLY	TPO	MET	LYS	ARG	ASN	ALA	THR	SER	P386	F393	MET	ASP	TYR	PHE	ASP	LYS	GLN	ASP
-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	------	------	-----	-----	-----	-----	-----	-----	-----	-----

4 Data and refinement statistics

Property	Value	Source
Space group	P 31 2 1	Depositor
Cell constants a, b, c, α , β , γ	142.19Å 142.19Å 135.98Å 90.00° 90.00° 120.00°	Depositor
Resolution (Å)	39.30 – 2.65 39.30 – 2.65	Depositor EDS
% Data completeness (in resolution range)	89.3 (39.30-2.65) 89.3 (39.30-2.65)	Depositor EDS
R_{merge}	0.09	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	3.96 (at 2.65Å)	Xtriage
Refinement program	PHENIX	Depositor
R, R_{free}	0.244 , 0.268 0.247 , 0.268	Depositor DCC
R_{free} test set	2004 reflections (4.83%)	wwPDB-VP
Wilson B-factor (Å ²)	38.4	Xtriage
Anisotropy	0.181	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.39 , 44.4	EDS
L-test for twinning ²	$\langle L \rangle = 0.42$, $\langle L^2 \rangle = 0.24$	Xtriage
Estimated twinning fraction	0.067 for -h,-k,l	Xtriage
F_o, F_c correlation	0.91	EDS
Total number of atoms	17464	wwPDB-VP
Average B, all atoms (Å ²)	56.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 6.41% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: TPO, ZN

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.33	0/1328	0.73	0/1800
1	B	0.35	0/1353	0.78	2/1835 (0.1%)
1	C	0.35	0/1371	0.73	0/1860
1	D	0.36	0/1253	0.75	1/1697 (0.1%)
1	E	0.34	0/1371	0.78	1/1860 (0.1%)
1	F	0.36	0/1371	0.81	1/1860 (0.1%)
2	G	0.34	0/175	0.70	0/232
2	H	0.33	0/141	0.54	0/181
2	I	0.33	0/101	0.56	0/130
2	J	0.35	0/109	0.75	0/141
2	K	0.35	0/90	0.76	0/116
2	L	0.30	0/101	0.68	0/130
2	M	0.28	0/144	0.93	0/193
2	N	0.28	0/104	0.73	0/140
2	O	0.24	0/69	0.61	0/92
All	All	0.34	0/9081	0.76	5/12267 (0.0%)

There are no bond length outliers.

All (5) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	E	100	GLY	N-CA-C	-8.20	101.50	112.82
1	B	99	ASP	N-CA-C	6.28	120.57	111.02
1	D	208	ILE	N-CA-C	5.83	116.57	110.62
1	B	100	GLY	N-CA-C	-5.32	97.87	113.30
1	F	98	ALA	N-CA-C	5.06	115.81	107.20

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts ⓘ

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1291	1247	1247	6	0
1	B	1315	1271	1271	10	0
1	C	1332	1297	1296	11	0
1	D	1218	1166	1166	14	0
1	E	1332	1296	1296	14	0
1	F	1332	1296	1296	9	0
2	G	183	167	167	2	0
2	H	151	137	138	2	0
2	I	110	97	98	0	0
2	J	118	102	104	2	0
2	K	99	84	85	2	0
2	L	110	98	100	2	0
2	M	141	136	136	4	0
2	N	101	92	92	1	0
2	O	67	66	66	0	0
3	A	1	0	0	0	0
3	B	1	0	0	0	0
3	C	1	0	0	0	0
3	D	1	0	0	0	0
3	E	1	0	0	0	0
3	F	1	0	0	0	0
4	A	1	0	0	0	0
4	B	1	0	0	0	0
4	C	1	0	0	0	0
4	F	3	0	0	0	0
All	All	8912	8552	8558	62	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 4.

The worst 5 of 62 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:52:ASP:N	1:B:55:GLU:OE2	2.18	0.76
1:E:99:ASP:OD2	2:K:378:TPO:O2P	2.05	0.73

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:199:ARG:NH2	1:A:200:GLU:OE2	2.25	0.69
1:E:69:ASN:ND2	1:E:119:MET:SD	2.65	0.69
1:E:175:GLU:OE1	1:E:178:HIS:ND1	2.25	0.68

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	150/177 (85%)	146 (97%)	4 (3%)	0	100	100
1	B	154/177 (87%)	148 (96%)	5 (3%)	1 (1%)	21	34
1	C	158/177 (89%)	156 (99%)	2 (1%)	0	100	100
1	D	138/177 (78%)	137 (99%)	1 (1%)	0	100	100
1	E	158/177 (89%)	153 (97%)	5 (3%)	0	100	100
1	F	158/177 (89%)	154 (98%)	4 (2%)	0	100	100
2	G	19/31 (61%)	18 (95%)	1 (5%)	0	100	100
2	H	12/31 (39%)	12 (100%)	0	0	100	100
2	I	8/31 (26%)	7 (88%)	1 (12%)	0	100	100
2	J	9/31 (29%)	8 (89%)	1 (11%)	0	100	100
2	K	7/31 (23%)	5 (71%)	2 (29%)	0	100	100
2	L	8/31 (26%)	7 (88%)	1 (12%)	0	100	100
2	M	15/31 (48%)	14 (93%)	1 (7%)	0	100	100
2	N	10/31 (32%)	10 (100%)	0	0	100	100
2	O	6/31 (19%)	6 (100%)	0	0	100	100
All	All	1010/1341 (75%)	981 (97%)	28 (3%)	1 (0%)	48	66

All (1) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	B	106	PRO

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	142/160 (89%)	141 (99%)	1 (1%)	76	86
1	B	145/160 (91%)	145 (100%)	0	100	100
1	C	147/160 (92%)	145 (99%)	2 (1%)	59	76
1	D	134/160 (84%)	134 (100%)	0	100	100
1	E	147/160 (92%)	145 (99%)	2 (1%)	59	76
1	F	147/160 (92%)	146 (99%)	1 (1%)	76	86
2	G	19/28 (68%)	19 (100%)	0	100	100
2	H	15/28 (54%)	15 (100%)	0	100	100
2	I	11/28 (39%)	11 (100%)	0	100	100
2	J	12/28 (43%)	12 (100%)	0	100	100
2	K	10/28 (36%)	10 (100%)	0	100	100
2	L	11/28 (39%)	11 (100%)	0	100	100
2	M	16/28 (57%)	16 (100%)	0	100	100
2	N	12/28 (43%)	12 (100%)	0	100	100
2	O	8/28 (29%)	8 (100%)	0	100	100
All	All	976/1212 (80%)	970 (99%)	6 (1%)	78	88

5 of 6 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	E	99	ASP
1	E	187	ILE
1	F	187	ILE
1	C	99	ASP
1	A	104	LYS

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 11 such sidechains are listed below:

Mol	Chain	Res	Type
1	D	205	GLN
1	F	96	HIS
1	F	205	GLN
1	F	178	HIS
1	C	165	GLN

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

6 non-standard protein/DNA/RNA residues are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
2	TPO	I	378	2	8,10,11	1.19	0	10,14,16	1.56	1 (10%)
2	TPO	L	378	2	8,10,11	1.14	0	10,14,16	1.50	1 (10%)
2	TPO	G	378	2	8,10,11	1.10	0	10,14,16	1.74	2 (20%)
2	TPO	H	378	2	8,10,11	1.17	0	10,14,16	1.84	2 (20%)
2	TPO	K	378	2	8,10,11	1.13	0	10,14,16	1.58	2 (20%)
2	TPO	J	378	2	8,10,11	1.17	0	10,14,16	1.63	2 (20%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	TPO	I	378	2	-	4/9/11/13	-
2	TPO	L	378	2	-	2/9/11/13	-

Continued on next page...

Continued from previous page...

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	TPO	G	378	2	-	1/9/11/13	-
2	TPO	H	378	2	-	1/9/11/13	-
2	TPO	K	378	2	-	0/9/11/13	-
2	TPO	J	378	2	-	4/9/11/13	-

There are no bond length outliers.

The worst 5 of 10 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed($^{\circ}$)	Ideal($^{\circ}$)
2	H	378	TPO	P-OG1-CB	-4.38	111.41	123.33
2	G	378	TPO	P-OG1-CB	-4.28	111.71	123.33
2	I	378	TPO	P-OG1-CB	-4.05	112.33	123.33
2	K	378	TPO	P-OG1-CB	-3.99	112.48	123.33
2	L	378	TPO	P-OG1-CB	-3.99	112.48	123.33

There are no chirality outliers.

5 of 12 torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
2	I	378	TPO	N-CA-CB-OG1
2	I	378	TPO	C-CA-CB-CG2
2	J	378	TPO	N-CA-CB-OG1
2	J	378	TPO	C-CA-CB-CG2
2	L	378	TPO	N-CA-CB-OG1

There are no ring outliers.

3 monomers are involved in 4 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	G	378	TPO	1	0
2	K	378	TPO	2	0
2	J	378	TPO	1	0

5.5 Carbohydrates

There are no oligosaccharides in this entry.

5.6 Ligand geometry

Of 6 ligands modelled in this entry, 6 are monoatomic - leaving 0 for Mogul analysis.

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

5.7 Other polymers

There are no such residues in this entry.

5.8 Polymer linkage issues

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	154/177 (87%)	-1.26	0 100 100	14, 29, 64, 106	0
1	B	158/177 (89%)	-1.05	0 100 100	16, 33, 85, 118	0
1	C	160/177 (90%)	-0.63	0 100 100	21, 54, 97, 121	0
1	D	144/177 (81%)	-0.24	0 100 100	44, 67, 114, 132	0
1	E	160/177 (90%)	-0.41	1 (0%) 85 83	28, 55, 111, 139	0
1	F	160/177 (90%)	-0.36	0 100 100	26, 66, 117, 143	0
2	G	21/31 (67%)	-1.06	0 100 100	26, 42, 59, 80	0
2	H	16/31 (51%)	-0.94	0 100 100	36, 56, 88, 111	0
2	I	11/31 (35%)	-0.43	0 100 100	50, 68, 97, 102	0
2	J	12/31 (38%)	0.36	1 (8%) 17 12	76, 97, 132, 143	0
2	K	10/31 (32%)	-0.28	0 100 100	59, 79, 103, 104	0
2	L	11/31 (35%)	0.17	0 100 100	80, 94, 108, 108	0
2	M	17/31 (54%)	-0.47	0 100 100	34, 64, 110, 144	0
2	N	12/31 (38%)	-0.82	0 100 100	39, 46, 72, 83	0
2	O	8/31 (25%)	-0.19	0 100 100	57, 71, 84, 101	0
All	All	1054/1341 (78%)	-0.64	2 (0%) 91 90	14, 53, 106, 144	0

All (2) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	E	108	LYS	2.6
2	J	391	PRO	2.5

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
2	TPO	I	378	11/12	0.98	0.07	25,55,83,84	0
2	TPO	L	378	11/12	0.98	0.06	21,72,102,102	0
2	TPO	J	378	11/12	0.99	0.04	18,59,77,84	0
2	TPO	K	378	11/12	0.99	0.04	10,32,43,44	0
2	TPO	G	378	11/12	0.99	0.03	23,28,31,33	0
2	TPO	H	378	11/12	1.00	0.02	25,34,45,47	0

6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
3	ZN	D	301	1/1	0.99	0.02	39,39,39,39	0
3	ZN	B	301	1/1	1.00	0.02	14,14,14,14	0
3	ZN	C	301	1/1	1.00	0.01	49,49,49,49	0
3	ZN	A	301	1/1	1.00	0.02	14,14,14,14	0
3	ZN	E	301	1/1	1.00	0.02	41,41,41,41	0
3	ZN	F	301	1/1	1.00	0.02	77,77,77,77	0

6.5 Other polymers [i](#)

There are no such residues in this entry.