



Full wwPDB X-ray Structure Validation Report ⓘ

Aug 25, 2026 – 09:14 am BST

PDB ID : 2C1V / pdb_00002c1v
Title : CRYSTAL STRUCTURE OF THE DI-HAEM CYTOCHROME C PEROXIDASE FROM PARACOCCLUS PANTOTROPHUS - Mixed VALENCE FORM
Authors : Echalier, A.; Fulop, V.
Deposited on : 2005-09-21
Resolution : 1.20 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4-5-2 with Phenix2.0
Mogul	:	1.8.4, CSD as541be (2020)
Xtriage (Phenix)	:	NOT EXECUTED
EDS	:	NOT EXECUTED
Buster-report	:	wwPDB partial adaption of 1.1.7 (2018)
Percentile statistics	:	20250101.v01 (using entries in the PDB archive January 1st 2025)
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.50


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 1.20 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
Clashscore	190562	1265 (1.20-1.20)
Ramachandran outliers	187476	1226 (1.20-1.20)
Sidechain outliers	187428	1226 (1.20-1.20)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$.

Note EDS was not executed.

Mol	Chain	Length	Quality of chain
1	A	338	96% . .
1	B	338	94% 5% .

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
2	HEC	A	401	X	-	-	-
2	HEC	A	402	X	-	-	-
2	HEC	B	401	X	-	-	-
2	HEC	B	402	X	-	-	-

Continued on next page...

Continued from previous page...

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
4	EDO	B	1340	-	-	X	-

2 Entry composition [i](#)


There are 5 unique types of molecules in this entry. The entry contains 6305 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called DI-HAEM CYTOCHROME C PEROXIDASE.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	335	Total	C	N	O	S	0	0	0
			2528	1594	422	499	13			
1	B	335	Total	C	N	O	S	0	0	0
			2528	1594	422	499	13			

- Molecule 2 is HEME C (CCD ID: HEC) (formula: C₃₄H₃₆FeN₄O₄).




Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
2	A	1	Total	C	Fe	N	O	0	0
			43	34	1	4	4		
2	A	1	Total	C	Fe	N	O	0	0
			43	34	1	4	4		
2	B	1	Total	C	Fe	N	O	0	0
			43	34	1	4	4		
2	B	1	Total	C	Fe	N	O	0	0
			43	34	1	4	4		

- Molecule 3 is CALCIUM ION (CCD ID: CA) (formula: Ca).

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
3	A	1	Total Ca 1 1	0	0
3	B	1	Total Ca 1 1	0	0

- Molecule 4 is 1,2-ETHANEDIOL (CCD ID: EDO) (formula: C₂H₆O₂).



Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
4	A	1	Total C O 6 3 3	0	1
4	A	1	Total C O 4 2 2	0	0
4	B	1	Total C O 6 3 3	0	1
4	B	1	Total C O 4 2 2	0	0

- Molecule 5 is water.


Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
5	A	509	Total O 509 509	0	0
5	B	546	Total O 546 546	0	0

3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


Note EDS was not executed.

- Molecule 1: DI-HAEM CYTOCHROME C PEROXIDASE

Chain A:  96%



- Molecule 1: DI-HAEM CYTOCHROME C PEROXIDASE

Chain B:  94% 5%



4 Data and refinement statistics

Xtriage (Phenix) and EDS were not executed - this section is therefore incomplete.

Property	Value	Source
Space group	P 21 21 2	Depositor
Cell constants a, b, c, α , β , γ	101.37Å 141.20Å 51.58Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	48.22 – 1.20	Depositor
% Data completeness (in resolution range)	92.5 (48.22-1.20)	Depositor
R_{merge}	0.07	Depositor
R_{sym}	(Not available)	Depositor
Refinement program	REFMAC 5.1.9999	Depositor
R, R_{free}	0.173 , 0.188	Depositor
Estimated twinning fraction	No twinning to report.	Xtriage
Total number of atoms	6305	wwPDB-VP
Average B, all atoms (Å ²)	18.0	wwPDB-VP

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: HEC, CA, EDO

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.74	0/2587	0.88	0/3523
1	B	0.78	3/2587 (0.1%)	0.86	1/3523 (0.0%)
All	All	0.76	3/5174 (0.1%)	0.87	1/7046 (0.0%)

All (3) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	B	183	ALA	C-O	6.87	1.32	1.24
1	B	143	MET	SD-CE	-5.50	1.65	1.79
1	B	28	ILE	C-O	5.09	1.29	1.23

All (1) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	B	328	ARG	NE-CZ-NH2	-5.57	114.19	119.20

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2528	0	2432	10	0
1	B	2528	0	2432	18	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
2	A	86	0	60	0	0
2	B	86	0	60	1	0
3	A	1	0	0	0	0
3	B	1	0	0	0	0
4	A	10	0	18	2	0
4	B	10	0	18	11	0
5	A	509	0	0	3	0
5	B	546	0	0	3	1
All	All	6305	0	5020	28	1

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 3.

All (28) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:251:ARG:HH11	1:A:251:ARG:HG2	1.29	0.97
1:A:294:ILE:HG22	1:A:296:THR:HG23	1.58	0.86
1:B:327:VAL:O	4:B:1340:EDO:H12	1.85	0.77
1:B:143:MET:HA	4:B:1340:EDO:H11	1.68	0.75
1:A:251:ARG:HG2	1:A:251:ARG:NH1	1.97	0.74
4:B:1340:EDO:H22	5:B:2308:HOH:O	1.92	0.70
1:B:326:PRO:HB3	4:B:1340:EDO:H21	1.78	0.65
1:B:143:MET:CA	4:B:1340:EDO:H11	2.26	0.64
1:B:205:ALA:HA	1:B:208:GLU:HG2	1.79	0.63
1:A:218:VAL:H	4:A:1340:EDO:C2	2.23	0.52
1:A:251:ARG:HH11	1:A:251:ARG:CG	2.13	0.51
1:B:144:PRO:CD	4:B:1340:EDO:H11	2.41	0.51
1:B:327:VAL:O	4:B:1340:EDO:C1	2.58	0.50
1:B:144:PRO:HD2	4:B:1340:EDO:H11	1.93	0.49
1:B:140:ILE:HG23	1:B:146:TYR:HB3	1.97	0.46
1:B:205:ALA:CB	5:B:2480:HOH:O	2.64	0.46
1:B:251:ARG:O	1:B:251:ARG:HD2	2.15	0.46
1:A:327:VAL:HG23	5:A:2281:HOH:O	2.15	0.46
1:B:95:PRO:HG2	2:B:401:HEC:HBA1	1.99	0.45
1:B:145:GLU:HG3	4:B:1340:EDO:O2	2.17	0.45
1:A:44:LYS:HE2	5:A:2290:HOH:O	2.17	0.44
1:A:218:VAL:H	4:A:1340:EDO:H21	1.83	0.44
1:B:205:ALA:HB1	5:B:2480:HOH:O	2.19	0.42
1:B:326:PRO:HB3	4:B:1340:EDO:C2	2.47	0.42
1:B:143:MET:HB3	4:B:1340:EDO:C1	2.50	0.41

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:12:GLU:HG2	5:A:2016:HOH:O	2.21	0.41
1:A:87:TRP:HB2	1:B:87:TRP:HB2	2.01	0.41
1:B:5:ILE:HG21	1:B:200:LYS:HD3	2.04	0.40

All (1) symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
5:B:2193:HOH:O	5:B:2193:HOH:O[2_565]	2.06	0.14

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	333/338 (98%)	326 (98%)	7 (2%)	0	100	100
1	B	333/338 (98%)	326 (98%)	7 (2%)	0	100	100
All	All	666/676 (98%)	652 (98%)	14 (2%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	264/267 (99%)	261 (99%)	3 (1%)	65	31

Continued on next page...

Continued from previous page...

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	B	264/267 (99%)	264 (100%)	0	100	100
All	All	528/534 (99%)	525 (99%)	3 (1%)	78	52

All (3) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	33	ASP
1	A	156	GLU
1	A	251	ARG

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (5) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	124	GLN
1	A	198	GLN
1	A	317	GLN
1	B	30	GLN
1	B	124	GLN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

Of 12 ligands modelled in this entry, 2 are monoatomic - leaving 10 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The

Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
4	EDO	B	1339[B]	-	3,3,3	0.40	0	2,2,2	0.25	0
4	EDO	A	1339[A]	-	3,3,3	0.46	0	2,2,2	0.35	0
4	EDO	A	1339[B]	-	3,3,3	0.51	0	2,2,2	0.05	0
4	EDO	B	1340	-	3,3,3	0.23	0	2,2,2	0.37	0
2	HEC	B	402	1	50,50,50	1.35	7 (14%)	68,82,82	1.21	3 (4%)
4	EDO	A	1340	-	3,3,3	0.36	0	2,2,2	0.31	0
2	HEC	B	401	5,1	50,50,50	1.21	4 (8%)	68,82,82	1.24	11 (16%)
2	HEC	A	401	5,1	50,50,50	1.31	6 (12%)	68,82,82	1.23	8 (11%)
4	EDO	B	1339[A]	-	3,3,3	0.47	0	2,2,2	0.30	0
2	HEC	A	402	1	50,50,50	1.30	6 (12%)	68,82,82	1.25	7 (10%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
4	EDO	B	1339[B]	-	-	0/1/1/1	-
4	EDO	A	1339[A]	-	-	0/1/1/1	-
4	EDO	A	1339[B]	-	-	1/1/1/1	-
4	EDO	B	1340	-	-	0/1/1/1	-
2	HEC	B	402	1	1/1/3/7	6/14/54/54	-
4	EDO	A	1340	-	-	1/1/1/1	-
2	HEC	B	401	5,1	1/1/3/7	4/14/54/54	-
2	HEC	A	401	5,1	1/1/3/7	4/14/54/54	-
4	EDO	B	1339[A]	-	-	0/1/1/1	-
2	HEC	A	402	1	1/1/3/7	8/14/54/54	-

All (23) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	B	402	HEC	C3D-C2D	5.03	1.47	1.36
2	A	402	HEC	C3D-C2D	4.37	1.46	1.36
2	A	401	HEC	C3D-C2D	4.31	1.45	1.36
2	B	401	HEC	C3D-C2D	4.26	1.45	1.36
2	A	402	HEC	FE-NC	2.96	2.05	1.95

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	B	402	HEC	FE-NC	2.90	2.05	1.95
2	B	402	HEC	FE-NA	2.67	2.04	1.95
2	A	402	HEC	FE-ND	2.58	2.02	1.94
2	B	401	HEC	FE-NC	2.44	2.03	1.95
2	A	401	HEC	FE-NB	2.33	2.02	1.94
2	B	401	HEC	FE-NA	2.29	2.02	1.95
2	A	402	HEC	FE-NB	2.28	2.01	1.94
2	A	402	HEC	CMC-C2C	2.26	1.55	1.50
2	A	401	HEC	CMB-C2B	2.24	1.55	1.50
2	B	402	HEC	CMA-C3A	2.18	1.55	1.50
2	A	401	HEC	CAB-C3B	2.10	1.56	1.51
2	B	401	HEC	FE-ND	2.08	2.01	1.94
2	A	402	HEC	C1D-ND	-2.05	1.36	1.40
2	A	401	HEC	O1A-CGA	2.04	1.28	1.22
2	A	401	HEC	O2D-CGD	-2.04	1.23	1.30
2	B	402	HEC	CMC-C2C	2.03	1.55	1.50
2	B	402	HEC	CAB-C3B	2.02	1.56	1.51
2	B	402	HEC	CMB-C2B	2.01	1.55	1.50

All (29) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	A	401	HEC	CBC-CAC-C3C	3.94	123.30	112.43
2	A	401	HEC	C1D-ND-C4D	3.94	109.14	105.07
2	B	402	HEC	CBD-CAD-C3D	-3.49	102.94	112.63
2	A	402	HEC	C1D-ND-C4D	3.32	108.50	105.07
2	B	401	HEC	C1D-ND-C4D	2.93	108.09	105.07
2	A	402	HEC	CAC-C3C-C2C	2.90	131.51	126.86
2	A	402	HEC	CAC-C3C-C4C	-2.87	120.94	124.92
2	A	402	HEC	CBD-CAD-C3D	-2.84	104.74	112.63
2	A	401	HEC	CMB-C2B-C1B	-2.54	121.17	125.04
2	B	401	HEC	CBC-CAC-C3C	2.54	119.43	112.43
2	B	401	HEC	C4A-NA-C1A	2.52	107.81	105.35
2	B	402	HEC	C1D-C2D-C3D	-2.48	104.35	106.96
2	B	401	HEC	CAD-C3D-C4D	2.37	128.80	124.66
2	A	402	HEC	C4C-NC-C1C	2.33	107.63	105.35
2	B	401	HEC	C1D-C2D-C3D	-2.33	104.50	106.96
2	B	401	HEC	C3C-C4C-NC	-2.33	107.54	110.15
2	B	401	HEC	O1D-CGD-CBD	-2.30	115.70	123.08
2	B	401	HEC	CAC-C3C-C4C	-2.28	121.76	124.92
2	B	401	HEC	C2A-C1A-NA	-2.25	108.13	110.32
2	A	401	HEC	CAD-C3D-C4D	2.25	128.60	124.66

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	A	402	HEC	O2A-CGA-CBA	2.21	121.13	114.03
2	A	401	HEC	CHB-C1B-C2B	2.19	128.41	124.98
2	A	402	HEC	CAD-CBD-CGD	-2.19	108.89	113.60
2	A	401	HEC	O2D-CGD-CBD	2.17	121.00	114.03
2	B	401	HEC	CMD-C2D-C1D	2.16	128.34	125.04
2	A	401	HEC	C4B-NB-C1B	2.16	107.30	105.07
2	B	402	HEC	C4C-NC-C1C	2.07	107.38	105.35
2	A	401	HEC	C2B-C1B-NB	-2.02	107.46	109.88
2	B	401	HEC	O2D-CGD-CBD	2.02	120.52	114.03

All (4) chirality outliers are listed below:

Mol	Chain	Res	Type	Atom
2	A	401	HEC	NA
2	A	402	HEC	NA
2	B	401	HEC	NA
2	B	402	HEC	NA

All (24) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
2	B	401	HEC	C4C-C3C-CAC-CBC
2	A	401	HEC	C4C-C3C-CAC-CBC
2	A	402	HEC	C4C-C3C-CAC-CBC
2	B	402	HEC	C2B-C3B-CAB-CBB
2	A	402	HEC	C2C-C3C-CAC-CBC
2	B	402	HEC	C4C-C3C-CAC-CBC
2	A	402	HEC	C2B-C3B-CAB-CBB
2	A	401	HEC	C2C-C3C-CAC-CBC
2	A	401	HEC	C2B-C3B-CAB-CBB
2	B	402	HEC	C4B-C3B-CAB-CBB
2	B	401	HEC	C2B-C3B-CAB-CBB
2	B	402	HEC	C2C-C3C-CAC-CBC
2	A	402	HEC	C4B-C3B-CAB-CBB
2	B	401	HEC	C2C-C3C-CAC-CBC
4	A	1340	EDO	O1-C1-C2-O2
2	B	401	HEC	C4B-C3B-CAB-CBB
2	A	402	HEC	CAD-CBD-CGD-O1D
2	A	402	HEC	CAD-CBD-CGD-O2D
2	B	402	HEC	CAD-CBD-CGD-O1D
2	B	402	HEC	CAD-CBD-CGD-O2D
2	A	401	HEC	C4B-C3B-CAB-CBB

Continued on next page...

Continued from previous page...


Mol	Chain	Res	Type	Atoms
4	A	1339[B]	EDO	O1-C1-C2-O2
2	A	402	HEC	CAA-CBA-CGA-O2A
2	A	402	HEC	CAA-CBA-CGA-O1A


There are no ring outliers.


3 monomers are involved in 14 short contacts:


Mol	Chain	Res	Type	Clashes	Symm-Clashes
4	B	1340	EDO	11	0
4	A	1340	EDO	2	0
2	B	401	HEC	1	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.









5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

EDS was not executed - this section is therefore empty.

6.2 Non-standard residues in protein, DNA, RNA chains

EDS was not executed - this section is therefore empty.

6.3 Carbohydrates

EDS was not executed - this section is therefore empty.

6.4 Ligands

EDS was not executed - this section is therefore empty.

6.5 Other polymers

EDS was not executed - this section is therefore empty.