



Full wwPDB X-ray Structure Validation Report ⓘ

Aug 5, 2026 – 08:16 PM EDT

PDB ID : 1C2R / pdb_00001c2r
Title : MOLECULAR STRUCTURE OF CYTOCHROME C2 ISOLATED FROM
RHODOBACTER CAPSULATUS DETERMINED AT 2.5 ANGSTROMS
RESOLUTION
Authors : Benning, M.M.; Wesenberg, G.; Caffrey, M.S.; Bartsch, R.G.; Meyer, T.E.;
Cusanovich, M.A.; Rayment, I.; Holden, H.M.
Deposited on : 1991-03-19
Resolution : 2.50 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references](#) ①) were used in the production of this report:

MolProbity	:	4-5-2 with Phenix2.0
Mogul	:	2022.3.0, CSD as543be (2022)
Xtriage (Phenix)	:	2.0
EDS	:	3.0
Buster-report	:	wwPDB partial adaption of 1.1.7 (2018)
Percentile statistics	:	20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4	:	9.0.015 (Gargrove)
Density-Fitness	:	1.0.12
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.50


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.50 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
Clashscore	190562	6492 (2.50-2.50)
Ramachandran outliers	187476	6378 (2.50-2.50)
Sidechain outliers	187428	6380 (2.50-2.50)
RSRZ outliers	180081	5833 (2.50-2.50)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	116	
1	B	116	

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:


Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
2	HEC	A	120	X	-	-	-
2	HEC	B	120	X	-	-	-

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called CYTOCHROME C2.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	116	Total 864	C 552	N 141	O 168	S 3	0	0	0
1	B	116	Total 864	C 552	N 141	O 168	S 3	0	0	0

- Molecule 2 is HEME C (CCD ID: HEC) (formula: $\text{C}_{34}\text{H}_{36}\text{FeN}_4\text{O}_4$).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
2	A	1	Total 43	C 34	Fe 1	N 4	O 4	0	0
2	B	1	Total 43	C 34	Fe 1	N 4	O 4	0	0


- Molecule 3 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	A	48	Total 48	O 48	0	0
3	B	44	Total 44	O 44	0	0


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

• Molecule 1: CYTOCHROME C2



• Molecule 1: CYTOCHROME C2



4 Data and refinement statistics

Property	Value	Source
Space group	H 3 2	Depositor
Cell constants a, b, c, α , β , γ	100.03Å 100.03Å 162.10Å 90.00° 90.00° 120.00°	Depositor
Resolution (Å)	(Not available) – 2.50 76.40 – 2.50	Depositor EDS
% Data completeness (in resolution range)	(Not available) ((Not available)-2.50) 86.3 (76.40-2.50)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	5.99 (at 2.50Å)	Xtriage
Refinement program	TNT	Depositor
R, R_{free}	0.168 , (Not available) 0.153 , (Not available)	Depositor DCC
R_{free} test set	No test flags present.	wwPDB-VP
Wilson B-factor (Å ²)	31.2	Xtriage
Anisotropy	0.090	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.37 , 129.8	EDS
L-test for twinning ²	$\langle L \rangle = 0.50$, $\langle L^2 \rangle = 0.33$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.96	EDS
Total number of atoms	1906	wwPDB-VP
Average B, all atoms (Å ²)	33.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 5.51% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: HEC

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	1.43	5/880 (0.6%)	1.80	21/1181 (1.8%)
1	B	1.32	1/880 (0.1%)	1.78	21/1181 (1.8%)
All	All	1.38	6/1760 (0.3%)	1.79	42/2362 (1.8%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	A	1	0

All (6) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	115	VAL	C-N	6.23	1.42	1.33
1	A	54	LYS	C-N	-5.42	1.27	1.33
1	A	82	PHE	N-CA	-5.31	1.40	1.46
1	B	115	VAL	CA-CB	-5.16	1.48	1.54
1	A	91	LYS	N-CA	-5.07	1.39	1.46
1	A	115	VAL	CA-C	5.04	1.59	1.52

All (42) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	55	ASP	N-CA-C	10.49	122.29	111.07
1	A	51	PHE	CA-CB-CG	-9.20	104.60	113.80
1	B	51	PHE	CA-CB-CG	-9.04	104.76	113.80
1	A	5	LYS	CB-CA-C	7.54	125.77	110.38
1	B	2	ASP	N-CA-C	7.29	120.15	108.41

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	B	55	ASP	N-CA-C	7.17	120.00	111.33
1	B	8	LYS	N-CA-CB	7.05	121.16	110.30
1	B	10	PHE	CA-CB-CG	-6.84	106.96	113.80
1	B	76	VAL	N-CA-C	6.63	119.33	111.05
1	B	70	GLU	CB-CG-CD	6.62	123.86	112.60
1	A	5	LYS	N-CA-CB	6.50	120.32	110.30
1	B	114	VAL	N-CA-C	6.50	118.03	110.62
1	A	113	SER	N-CA-C	6.48	118.35	111.28
1	B	17	HIS	CA-CB-CG	6.41	120.21	113.80
1	B	56	SER	N-CA-C	6.35	118.00	111.14
1	A	54	LYS	N-CA-C	-6.30	102.05	110.55
1	B	105	GLU	N-CA-C	6.25	119.06	111.82
1	A	114	VAL	N-CA-C	5.96	119.39	111.17
1	B	88	ASP	N-CA-C	-5.95	103.94	111.92
1	A	102	LYS	N-CA-CB	-5.83	100.58	111.13
1	A	15	THR	N-CA-CB	-5.78	101.31	110.22
1	A	11	ASN	CA-CB-CG	5.72	118.32	112.60
1	A	65	PHE	N-CA-C	5.70	118.51	109.96
1	B	84	LYS	CA-CB-CG	-5.70	102.70	114.10
1	A	8	LYS	CA-C-N	5.63	131.00	121.14
1	A	8	LYS	C-N-CA	5.63	131.00	121.14
1	A	101	ALA	N-CA-C	5.61	118.24	111.40
1	B	22	PRO	N-CA-CB	5.55	109.39	103.39
1	B	3	ALA	CA-C-N	5.49	127.57	120.44
1	B	3	ALA	C-N-CA	5.49	127.57	120.44
1	B	12	LYS	CB-CA-C	5.40	120.53	109.67
1	B	14	LYS	CB-CA-C	5.35	122.16	110.18
1	B	54	LYS	N-CA-C	-5.28	103.42	110.55
1	A	29	LYS	N-CA-CB	-5.28	101.66	110.37
1	A	83	LEU	N-CA-C	5.27	116.71	111.07
1	B	88	ASP	CA-CB-CG	-5.19	107.41	112.60
1	A	105	GLU	N-CA-CB	-5.15	102.29	110.28
1	B	12	LYS	N-CA-CB	5.14	118.04	110.33
1	A	76	VAL	CB-CA-C	5.12	118.75	112.04
1	A	48	TYR	CA-C-N	5.09	125.02	119.78
1	A	48	TYR	C-N-CA	5.09	125.02	119.78
1	A	29	LYS	CA-CB-CG	5.01	124.11	114.10

All (1) chirality outliers are listed below:

Mol	Chain	Res	Type	Atom
1	A	5	LYS	CA

There are no planarity outliers.

5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	864	0	872	25	1
1	B	864	0	872	40	1
2	A	43	0	31	7	0
2	B	43	0	31	8	0
3	A	48	0	0	1	1
3	B	44	0	0	3	1
All	All	1906	0	1806	65	2

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 18.

All (65) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:16:CYS:SG	2:B:120:HEC:HAC	1.38	1.60
1:A:16:CYS:SG	2:A:120:HEC:CAC	2.08	1.41
1:A:16:CYS:SG	2:A:120:HEC:HAC	1.63	1.38
1:B:16:CYS:SG	2:B:120:HEC:CAC	2.35	1.15
1:B:2:ASP:OD2	1:B:4:ALA:HB3	1.60	0.99
1:A:11:ASN:HD21	1:A:12:LYS:NZ	1.76	0.84
1:A:11:ASN:HD21	1:A:12:LYS:HZ1	1.31	0.75
1:A:60:LEU:O	1:A:63:SER:OG	2.06	0.74
1:B:69:GLU:OE1	3:B:145:HOH:O	2.07	0.72
1:B:48:TYR:O	3:B:150:HOH:O	2.09	0.70
1:A:20:ILE:HD13	1:A:26:GLU:HG3	1.76	0.67
1:B:16:CYS:CB	2:B:120:HEC:HAC	2.25	0.66
1:A:76:VAL:HG21	2:A:120:HEC:HMB1	1.77	0.66
1:A:50:GLU:HA	1:B:62:ALA:O	1.96	0.66
1:A:16:CYS:SG	2:A:120:HEC:C3C	2.85	0.63
1:B:87:LEU:O	1:B:88:ASP:CB	2.46	0.63
1:B:16:CYS:HA	1:B:31:ALA:HB3	1.84	0.60
1:B:16:CYS:SG	1:B:33:THR:HG21	2.42	0.60

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:38:TYR:CZ	1:A:115:VAL:HG12	2.38	0.59
1:B:2:ASP:HB3	1:B:5:LYS:HB2	1.86	0.56
1:B:96:MET:HG3	2:B:120:HEC:CHD	2.37	0.54
1:B:96:MET:HG3	2:B:120:HEC:C4C	2.37	0.54
1:B:111:LEU:O	1:B:114:VAL:HG22	2.07	0.54
1:A:11:ASN:ND2	1:A:12:LYS:NZ	2.54	0.53
1:B:36:ASN:ND2	1:B:38:TYR:H	2.07	0.53
1:B:68:THR:HB	1:B:70:GLU:OE2	2.08	0.53
1:A:78:ASP:CG	1:A:99:LYS:HE2	2.34	0.53
1:B:16:CYS:HB2	2:B:120:HEC:C3C	2.39	0.52
1:A:5:LYS:O	1:A:9:GLU:HG2	2.10	0.51
1:A:16:CYS:SG	2:A:120:HEC:CBC	2.93	0.50
1:A:100:LEU:HD23	3:A:133:HOH:O	2.11	0.49
1:B:17:HIS:HE1	1:B:35:PRO:HD2	1.76	0.49
1:B:44:THR:O	3:B:147:HOH:O	2.19	0.49
1:A:76:VAL:HG21	2:A:120:HEC:CMB	2.42	0.48
1:B:17:HIS:CE1	1:B:35:PRO:HD2	2.49	0.48
1:B:16:CYS:SG	1:B:33:THR:CG2	3.01	0.48
1:B:73:ALA:CB	1:B:105:GLU:HG3	2.43	0.47
1:B:110:TYR:O	1:B:114:VAL:HG13	2.14	0.47
1:A:16:CYS:HA	1:A:31:ALA:HB3	1.96	0.47
1:B:73:ALA:HB1	1:B:105:GLU:HG3	1.97	0.47
1:B:86:LYS:HD3	1:B:86:LYS:HA	1.73	0.47
1:A:9:GLU:C	1:A:11:ASN:H	2.22	0.47
1:B:104:GLY:O	1:B:107:VAL:HG12	2.15	0.46
1:B:9:GLU:OE1	1:B:12:LYS:HE3	2.16	0.46
1:A:46:GLY:HA2	1:A:53:TYR:CE2	2.51	0.45
1:A:84:LYS:HG2	1:A:89:ASP:O	2.16	0.45
1:B:22:PRO:HD3	1:B:38:TYR:CE1	2.51	0.45
1:B:84:LYS:CG	1:B:92:ALA:HB3	2.47	0.45
1:A:23:ASP:OD1	1:A:23:ASP:C	2.59	0.44
1:B:3:ALA:O	1:B:7:GLU:N	2.42	0.44
1:B:103:GLY:O	1:B:104:GLY:C	2.58	0.44
1:A:9:GLU:C	1:A:11:ASN:N	2.76	0.44
1:B:20:ILE:H	1:B:36:ASN:HD21	1.66	0.44
1:B:23:ASP:C	1:B:23:ASP:OD2	2.58	0.44
1:B:84:LYS:HG2	1:B:92:ALA:HB3	2.00	0.43
1:B:36:ASN:HD22	1:B:37:LEU:N	2.16	0.43
1:B:79:PRO:HG3	1:B:96:MET:HE2	2.00	0.43
1:B:10:PHE:CZ	1:B:19:ILE:HD11	2.54	0.43
1:B:13:CYS:HA	2:B:120:HEC:HHC	2.01	0.43

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
2:A:120:HEC:HMB3	2:A:120:HEC:HBB3	2.01	0.42
1:A:71:ASP:O	1:A:72:ILE:C	2.62	0.42
1:A:48:TYR:CG	1:A:49:PRO:HD2	2.54	0.42
1:B:96:MET:HG3	2:B:120:HEC:C1D	2.51	0.41
1:A:48:TYR:HA	1:A:49:PRO:HD3	1.93	0.40
1:B:54:LYS:HD2	1:B:54:LYS:HA	1.84	0.40

All (2) symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:5:LYS:NZ	1:B:114:VAL:O[5_565]	1.93	0.27
3:A:161:HOH:O	3:B:154:HOH:O[4_555]	2.00	0.20

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	114/116 (98%)	113 (99%)	1 (1%)	0	100	100
1	B	114/116 (98%)	106 (93%)	8 (7%)	0	100	100
All	All	228/232 (98%)	219 (96%)	9 (4%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was

analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	86/86 (100%)	71 (83%)	15 (17%)	2	3
1	B	86/86 (100%)	80 (93%)	6 (7%)	14	29
All	All	172/172 (100%)	151 (88%)	21 (12%)	5	10

All (21) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	5	LYS
1	A	11	ASN
1	A	14	LYS
1	A	20	ILE
1	A	26	GLU
1	A	32	LYS
1	A	55	ASP
1	A	76	VAL
1	A	86	LYS
1	A	93	LYS
1	A	102	LYS
1	A	105	GLU
1	A	113	SER
1	A	114	VAL
1	A	116	LYS
1	B	2	ASP
1	B	12	LYS
1	B	14	LYS
1	B	54	LYS
1	B	63	SER
1	B	111	LEU

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (2) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	11	ASN
1	B	36	ASN

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

2 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
2	HEC	B	120	1	50,50,50	1.35	7 (14%)	68,82,82	1.89	18 (26%)
2	HEC	A	120	1	50,50,50	1.74	13 (26%)	68,82,82	1.69	17 (25%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	HEC	B	120	1	1/1/3/7	2/14/54/54	-
2	HEC	A	120	1	1/1/3/7	5/14/54/54	-

All (20) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	A	120	HEC	C4B-NB	-5.10	1.31	1.40
2	B	120	HEC	C4B-NB	-4.39	1.32	1.40
2	A	120	HEC	O2A-CGA	-3.29	1.19	1.30
2	A	120	HEC	C4B-C3B	-3.28	1.39	1.45
2	A	120	HEC	CHA-C1A	-3.14	1.32	1.38
2	B	120	HEC	O2A-CGA	-2.96	1.21	1.30

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	A	120	HEC	C4D-ND	-2.92	1.33	1.38
2	A	120	HEC	CHC-C1C	-2.91	1.32	1.39
2	B	120	HEC	C3C-C2C	-2.83	1.31	1.38
2	A	120	HEC	C3B-C2B	-2.71	1.30	1.36
2	A	120	HEC	C3C-C2C	-2.54	1.32	1.38
2	A	120	HEC	CAB-C3B	-2.42	1.44	1.51
2	B	120	HEC	CMB-C2B	2.25	1.55	1.50
2	A	120	HEC	C3D-C2D	-2.24	1.31	1.36
2	A	120	HEC	CBA-CGA	2.20	1.55	1.50
2	B	120	HEC	CBA-CGA	2.09	1.55	1.50
2	B	120	HEC	CHA-C1A	-2.06	1.34	1.38
2	B	120	HEC	C3B-C2B	-2.06	1.32	1.36
2	A	120	HEC	CMB-C2B	2.03	1.54	1.50
2	A	120	HEC	C2A-C3A	-2.01	1.32	1.36

All (35) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	B	120	HEC	CBB-CAB-C3B	6.19	129.21	112.42
2	B	120	HEC	CBC-CAC-C3C	5.80	128.15	112.42
2	A	120	HEC	CBC-CAC-C3C	5.34	126.89	112.42
2	A	120	HEC	CBB-CAB-C3B	4.55	124.74	112.42
2	A	120	HEC	CHB-C1B-NB	4.38	129.13	124.42
2	B	120	HEC	CAD-CBD-CGD	3.75	123.61	113.67
2	B	120	HEC	CBD-CAD-C3D	-3.43	103.06	112.53
2	B	120	HEC	O2A-CGA-CBA	3.39	124.70	114.00
2	B	120	HEC	O1A-CGA-CBA	-3.38	112.38	123.09
2	B	120	HEC	CHD-C4C-NC	-3.25	117.97	123.86
2	B	120	HEC	C3D-C4D-ND	3.02	113.27	110.35
2	A	120	HEC	CHC-C4B-NB	2.97	128.05	124.37
2	A	120	HEC	O2A-CGA-CBA	2.90	123.17	114.00
2	A	120	HEC	O1A-CGA-CBA	-2.87	113.97	123.09
2	B	120	HEC	CHA-C1A-NA	-2.85	121.36	124.45
2	A	120	HEC	CBD-CAD-C3D	-2.61	105.31	112.53
2	A	120	HEC	CHB-C4A-NA	-2.58	121.64	124.45
2	B	120	HEC	C1B-C2B-C3B	-2.47	104.38	106.98
2	B	120	HEC	CHA-C4D-ND	-2.47	121.77	124.42
2	A	120	HEC	C2C-C1C-NC	2.40	113.99	110.14
2	B	120	HEC	CHB-C4A-NA	-2.37	121.87	124.45
2	B	120	HEC	C4A-NA-C1A	-2.28	102.11	105.82
2	A	120	HEC	C4C-NC-C1C	-2.26	102.13	105.82
2	A	120	HEC	CHC-C1C-NC	-2.23	119.81	123.86

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	B	120	HEC	CMB-C2B-C3B	2.20	132.10	126.15
2	A	120	HEC	CAD-CBD-CGD	2.19	119.48	113.67
2	A	120	HEC	C4B-C3B-C2B	2.19	110.08	106.89
2	A	120	HEC	C3D-C4D-ND	2.15	112.43	110.35
2	A	120	HEC	C4B-NB-C1B	2.12	107.72	105.21
2	B	120	HEC	C4C-NC-C1C	-2.11	102.38	105.82
2	B	120	HEC	CHD-C4C-C3C	2.10	129.90	125.30
2	A	120	HEC	CAA-CBA-CGA	2.09	119.21	113.67
2	A	120	HEC	CMC-C2C-C3C	2.08	130.03	125.62
2	B	120	HEC	C2B-C1B-NB	2.02	112.24	109.90
2	B	120	HEC	CAA-CBA-CGA	2.01	118.99	113.67

All (2) chirality outliers are listed below:

Mol	Chain	Res	Type	Atom
2	A	120	HEC	NA
2	B	120	HEC	NA

All (7) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
2	A	120	HEC	C2B-C3B-CAB-CBB
2	A	120	HEC	C2C-C3C-CAC-CBC
2	A	120	HEC	C3D-CAD-CBD-CGD
2	B	120	HEC	C2B-C3B-CAB-CBB
2	A	120	HEC	C4B-C3B-CAB-CBB
2	A	120	HEC	C4C-C3C-CAC-CBC
2	B	120	HEC	C4B-C3B-CAB-CBB


There are no ring outliers.


2 monomers are involved in 15 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	B	120	HEC	8	0
2	A	120	HEC	7	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be

highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.





5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	116/116 (100%)	-0.71	0 100 100	9, 24, 63, 100	0
1	B	116/116 (100%)	-0.54	0 100 100	9, 29, 65, 80	0
All	All	232/232 (100%)	-0.62	0 100 100	9, 26, 65, 100	0

There are no RSRZ outliers to report.

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.4 Ligands [i](#)




In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled ‘Q< 0.9’ lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
2	HEC	B	120	43/43	0.97	0.07	1,20,38,41	0
2	HEC	A	120	43/43	0.98	0.06	3,14,27,35	0

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.



Electron density around HEC B 120:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)



Electron density around HEC A 120:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)



6.5 Other polymers ⓘ

There are no such residues in this entry.