Full wwPDB X-ray Structure Validation Report (i)

Aug 25, 2026 – 11:17 am BST

PDB ID	:	8C7P / pdb_00008c7p
Title	:	Tagless BtuM in complex with cyanocobalamin
Authors	:	Martinez-Felices, J.M.; Slotboom, D.J.
Deposited on	:	2023-01-16
Resolution	:	3.00 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at *validation@mail.wwpdb.org*
A user guide is available at
https://www.wwpdb.org/validation/2017/XrayValidationReportHelp
with specific help available everywhere you see the (i) symbol.

The types of validation reports are described at
http://www.wwpdb.org/validation/2017/FAQs#types.

The following versions of software and data (see references (i)) were used in the production of this report:


MolProbity	:	4-5-2 with Phenix2.0
Mogul	:	1.8.4, CSD as541be (2020)
Xtriage (Phenix)	:	2.0
EDS	:	3.0
Buster-report	:	wwPDB partial adaption of 1.1.7 (2018)
Percentile statistics	:	20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4	:	9.0.015 (Gargrove)
Density-Fitness	:	1.0.12
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.50

1 Overall quality at a glance (i)

The following experimental techniques were used to determine the structure:
X-RAY DIFFRACTION

The reported resolution of this entry is 3.00 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	2672 (3.00-3.00)
Clashscore	190562	2977 (3.00-3.00)
Ramachandran outliers	187476	2877 (3.00-3.00)
Sidechain outliers	187428	2880 (3.00-3.00)
RSRZ outliers	180081	2671 (3.00-3.00)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for >=3, 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions <=5% The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	199	10%, 63%, 24%, 10%

2 Entry composition

There are 2 unique types of molecules in this entry. The entry contains 1474 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.


- Molecule 1 is a protein called Cobalamin ABC transporter.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	179	1383	932	226	220	5	0	1	0

There are 2 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	1	GLY	-	expression tag	UNP Q3SFD8
A	2	PRO	-	expression tag	UNP Q3SFD8

- Molecule 2 is COBALAMIN (CCD ID: B12) (formula: $C_{62}H_{89}CoN_{13}O_{14}P$) (labeled as "Ligand of Interest" by depositor).




Mol	Chain	Residues	Atoms						ZeroOcc	AltConf
			Total	C	Co	N	O	P		
2	A	1	91	62	1	13	14	1	0	0

3 Residue-property plots (i)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density (RSRZ > 2). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: Cobalamin ABC transporter



4 Data and refinement statistics (i)

Property	Value	Source
Space group	P 32 2 1	Depositor
Cell constants a, b, c, α, β, γ	91.32Å 91.32Å 77.39Å 90.00° 90.00° 120.00°	Depositor
Resolution (Å)	79.08 – 3.00 79.08 – 3.00	Depositor EDS
% Data completeness (in resolution range)	80.7 (79.08-3.00) 80.7 (79.08-3.00)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	(Not available)	Depositor
$< I/\sigma(I) >$ [1]	3.07 (at 3.01Å)	Xtriage
Refinement program	REFMAC 5.8.0405	Depositor
R, R_{free}	0.225 , 0.263 0.223 , 0.261	Depositor DCC
R_{free} test set	295 reflections (3.80%)	wwPDB-VP
Wilson B-factor (Å^2)	56.8	Xtriage
Anisotropy	0.336	Xtriage
Bulk solvent k_{sol}(e/Å^3), B_{sol}(Å^2)	0.36 , 80.8	EDS
L-test for twinning[2]	$< \|L\| > = 0.49$, $< L^2 > = 0.32$	Xtriage
Estimated twinning fraction	0.029 for -h,-k,l	Xtriage
F_o,F_c correlation	0.90	EDS
Total number of atoms	1474	wwPDB-VP
Average B, all atoms (Å^2)	81.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 6.34% of the height of the origin peak. No significant pseudotranslation is detected.*

[1]Intensities estimated from amplitudes.

[2]Theoretical values of $< |L| >$, $< L^2 >$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: B12

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	#$\|Z\|>5$	RMSZ	#$\|Z\|>5$
1	A	0.82	0/1428	1.37	10/1955 (0.5%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand.A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	A	0	3

There are no bond length outliers.

All (10) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(^{o})	Ideal(^{o})
1	A	165	PHE	CA-C-N	6.05	129.19	120.79
1	A	165	PHE	C-N-CA	6.05	129.19	120.79
1	A	24	THR	CA-CB-OG1	-5.53	101.30	109.60
1	A	65	VAL	N-CA-CB	-5.50	104.41	110.51
1	A	68	TYR	CB-CA-C	-5.48	102.04	110.81
1	A	143	ALA	CA-C-N	-5.44	116.98	122.74
1	A	143	ALA	C-N-CA	-5.44	116.98	122.74
1	A	60	ILE	N-CA-C	-5.27	105.26	111.00
1	A	57	PRO	CB-CA-C	5.24	119.33	111.68
1	A	153	ARG	CB-CG-CD	5.08	122.99	111.30

There are no chirality outliers.

All (3) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	A	153	ARG	Sidechain
1	A	25	ARG	Sidechain
1	A	8	ARG	Sidechain

5.2 Too-close contacts (i)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1383	0	1428	26	0
2	A	91	0	88	11	0
All	All	1474	0	1516	34	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 11.

All (34) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:98:GLY:HA3	1:A:123:VAL:HG11	1.52	0.91
2:A:201:B12:H552	2:A:201:B12:H531	1.61	0.83
1:A:147:LEU:O	1:A:151:VAL:HG23	1.91	0.70
1:A:137:PHE:HE2	1:A:150:LEU:HD22	1.56	0.69
1:A:158:PHE:HB3	1:A:159:PRO:HD3	1.87	0.57
2:A:201:B12:H252	2:A:201:B12:C61	2.35	0.57
1:A:128:GLN:NE2	1:A:132:THR:OG1	2.36	0.55
1:A:122:ALA:O	1:A:126:VAL:HG23	2.07	0.53
1:A:98:GLY:HA3	1:A:123:VAL:CG1	2.33	0.53
2:A:201:B12:H351	2:A:201:B12:H371	1.93	0.51
1:A:161:MET:HE1	2:A:201:B12:H601	1.94	0.49
1:A:31:SER:HB3	1:A:34:SER:HB2	1.95	0.48
1:A:88:LEU:HD11	2:A:201:B12:C35	2.44	0.48
1:A:178:ALA:O	1:A:182:VAL:HG23	2.14	0.47
2:A:201:B12:H531	2:A:201:B12:C55	2.32	0.47
1:A:76:VAL:HG11	2:A:201:B12:H451	1.81	0.46
1:A:56:VAL:N	1:A:57:PRO:HD2	2.31	0.45
1:A:31:SER:HB3	1:A:34:SER:CB	2.46	0.45
1:A:9:GLN:O	1:A:13:ILE:HG12	2.18	0.44

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:106:HIS:CG	1:A:107:SER:H	2.36	0.44
1:A:24:THR:C	1:A:26:SER:H	2.26	0.43
2:A:201:B12:C15	2:A:201:B12:H492	2.47	0.42
1:A:134:GLY:O	1:A:138:TYR:HB2	2.19	0.42
2:A:201:B12:H262	2:A:201:B12:H91	1.82	0.42
1:A:175:VAL:O	1:A:179:LEU:HG	2.20	0.42
1:A:123:VAL:HG23	1:A:124:VAL:N	2.33	0.42
1:A:157:TYR:HB3	1:A:161:MET:HE3	2.01	0.41
1:A:84:ALA:O	1:A:85:TYR:C	2.63	0.41
1:A:46:LEU:HD11	1:A:59:LEU:HD12	2.03	0.41
1:A:79:PHE:HA	1:A:141:ARG:HH11	1.85	0.41
2:A:201:B12:H562	2:A:201:B12:H18	1.79	0.41
2:A:201:B12:H421	2:A:201:B12:H361	2.02	0.40
1:A:44:PHE:HA	1:A:95:LEU:O	2.22	0.40
1:A:175:VAL:O	1:A:175:VAL:CG1	2.69	0.40

There are no symmetry-related clashes.

5.3 Torsion angles (i)

5.3.1 Protein backbone (i)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
1	A	178/199 (89%)	162 (91%)	16 (9%)	0	100 100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains (i)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was

analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles
1	A	138/154 (90%)	131 (95%)	7 (5%)	21 55

All (7) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	35	LEU
1	A	64	VAL
1	A	65	VAL
1	A	115	ARG
1	A	146	THR
1	A	147	LEU
1	A	153	ARG

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (2) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	27	HIS
1	A	128	GLN

5.3.3 RNA

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates

There are no oligosaccharides in this entry.

5.6 Ligand geometry

1 ligand is modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The

Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\#\|Z\| > 2$	Counts	RMSZ	$\#\|Z\| > 2$
2	B12	A	201	1	93,101,101	1.06	4 (4%)	146,166,166	1.43	24 (16%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	B12	A	201	1	-	14/56/223/223	0/3/11/11

All (4) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	A	201	B12	C14-N23	6.78	1.43	1.35
2	A	201	B12	C9-N22	4.02	1.41	1.30
2	A	201	B12	C26-C27	2.37	1.59	1.51
2	A	201	B12	C1P-C2P	2.36	1.57	1.51

All (24) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(^{o})	Ideal(^{o})
2	A	201	B12	C2P-C1P-N59	-6.25	103.72	112.93
2	A	201	B12	C55-C56-C57	-3.93	102.64	111.23
2	A	201	B12	C36-C7-C37	-3.55	104.95	110.80
2	A	201	B12	C8B-N1B-C1R	-3.25	119.85	125.39
2	A	201	B12	C17-C18-C19	3.23	107.30	102.36
2	A	201	B12	C56-C55-C17	-3.21	109.32	115.52
2	A	201	B12	C5-C6-N22	-3.12	119.12	123.88
2	A	201	B12	C25-C2-C1	2.90	118.15	113.78
2	A	201	B12	O3-P-O4	2.86	120.22	109.47
2	A	201	B12	C20-C1-C19	-2.80	106.66	109.36
2	A	201	B12	C12-C13-C14	-2.66	97.81	102.26
2	A	201	B12	C7-C6-N22	2.63	112.75	107.94
2	A	201	B12	C12-C11-C10	2.57	126.71	123.37
2	A	201	B12	C25-C2-C26	-2.56	104.54	109.71
2	A	201	B12	C48-C13-C12	-2.50	109.63	116.63
2	A	201	B12	O2-P-O4	2.41	118.50	109.47

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(^{o})	Ideal(^{o})
2	A	201	B12	C30-C3-C2	-2.38	114.04	119.09
2	A	201	B12	C3P-C2P-C1P	2.36	115.98	111.39
2	A	201	B12	C3-C4-C5	-2.29	119.95	123.81
2	A	201	B12	C3-C4-N21	2.13	114.64	111.97
2	A	201	B12	C20-C1-C2	2.10	116.80	113.28
2	A	201	B12	C37-C7-C6	2.10	113.73	107.12
2	A	201	B12	C13-C12-C11	2.08	103.32	100.97
2	A	201	B12	C1-C19-N24	2.05	108.55	106.24

There are no chirality outliers.

All (14) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
2	A	201	B12	C56-C57-N59-C1P
2	A	201	B12	O58-C57-N59-C1P
2	A	201	B12	O6R-C4R-C5R-O8R
2	A	201	B12	C3R-C4R-C5R-O8R
2	A	201	B12	C13-C48-C49-C50
2	A	201	B12	C41-C42-C43-O44
2	A	201	B12	C41-C42-C43-N45
2	A	201	B12	C19-C18-C60-C61
2	A	201	B12	C18-C60-C61-O63
2	A	201	B12	C55-C56-C57-O58
2	A	201	B12	C55-C56-C57-N59
2	A	201	B12	C17-C18-C60-C61
2	A	201	B12	C2-C3-C30-C31
2	A	201	B12	C18-C60-C61-N62

There are no ring outliers.


1 monomer is involved in 11 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	A	201	B12	11	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less then 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the

average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.

Ligand B12 A 201



5.7 Other polymers (i)

There are no such residues in this entry.

5.8 Polymer linkage issues (i)

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

In the following table, the column labelled '#RSRZ> 2' contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95^{th} percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled 'Q< 0.9' lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å^2)	Q<0.9
1	A	179/199 (89%)	0.59	20 (11%) 10 6	39, 73, 151, 169	1 (0%)

All (20) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	108	LEU	6.4
1	A	111	LEU	5.8
1	A	113	LEU	4.3
1	A	110	LEU	4.2
1	A	106	HIS	4.1
1	A	32	ILE	4.0
1	A	182	VAL	4.0
1	A	114	PHE	3.4
1	A	107	SER	3.4
1	A	177	VAL	3.2
1	A	178	ALA	2.9
1	A	105	GLY	2.6
1	A	180	ALA	2.3
1	A	138	TYR	2.3
1	A	104	ARG	2.3
1	A	14	GLY	2.2
1	A	179	LEU	2.2
1	A	109	SER	2.1
1	A	175	VAL	2.1
1	A	13	ILE	2.0

6.2 Non-standard residues in protein, DNA, RNA chains

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates (i)

There are no oligosaccharides in this entry.

6.4 Ligands (i)


In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95^{th} percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.


Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å^2)	Q<0.9
2	B12	A	201	91/91	0.96	0.10	60,72,133,140	0


The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.

Electron density around B12 A 201:

$2mF_o$-DF_c (at 0.7 rmsd) in gray
mF_o-DF_c (at 3 rmsd) in purple (negative) and green (positive)







6.5 Other polymers (i)

There are no such residues in this entry.