



Full wwPDB X-ray Structure Validation Report ⓘ

Mar 5, 2026 – 03:13 PM UTC

PDB ID : 3CA0 / pdb_00003ca0
Title : Sambucus nigra agglutinin II (SNA-II), hexagonal crystal form
Authors : Maveyraud, L.; Niwa, H.; Guillet, V.; Palmer, R.A.; Reynolds, C.D.; Mourey, L.
Deposited on : 2008-02-19
Resolution : 1.95 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4-5-2 with Phenix2.0
Mogul	:	2022.3.0, CSD as543be (2022)
Xtriage (Phenix)	:	2.0
EDS	:	3.0
Percentile statistics	:	20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4	:	9.0.010 (Gargrove)
Density-Fitness	:	1.0.12
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 1.95 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	3494 (1.96-1.96)
Clashscore	190562	3612 (1.96-1.96)
Ramachandran outliers	187476	3587 (1.96-1.96)
Sidechain outliers	187428	3587 (1.96-1.96)
RSRZ outliers	180081	3495 (1.96-1.96)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	258	37% 92% 7%
2	B	7	100%
3	C	2	100%
4	D	2	100%
5	E	3	100%

2 Entry composition [i](#)

There are 8 unique types of molecules in this entry. The entry contains 2570 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.


- Molecule 1 is a protein called Agglutinin II.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	257	2026	1254	363	395	14	0	6	0

There is a discrepancy between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	224	LEU	HIS	SEE REMARK 999	UNP P33183

- Molecule 2 is an oligosaccharide called beta-D-xylopyranose-(1-2)-[alpha-D-mannopyranose-(1-3)][alpha-D-mannopyranose-(1-6)]beta-D-mannopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-[alpha-L-fucopyranose-(1-3)]2-acetamido-2-deoxy-beta-D-glucopyranose.



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	Trace
			Total	C	N	O			
2	B	7	80	45	2	33	0	0	0

- Molecule 3 is an oligosaccharide called alpha-L-fucopyranose-(1-3)-2-acetamido-2-deoxy-beta-D-glucopyranose.




Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	Trace
			Total	C	N	O			
3	C	2	24	14	1	9	0	0	0

- Molecule 4 is an oligosaccharide called 2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose.




Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	Trace
4	D	2	Total	C	N	O	0	0	0
			28	16	2	10			

- Molecule 5 is an oligosaccharide called alpha-L-fucopyranose-(1-3)-[2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)]2-acetamido-2-deoxy-beta-D-glucopyranose.



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	Trace
5	E	3	Total	C	N	O	0	0	0
			38	22	2	14			

- Molecule 6 is SULFATE ION (CCD ID: SO4) (formula: O₄S).




Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
6	A	1	Total	O	S	0	0
			5	4	1		

Continued on next page...

Continued from previous page...

Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
6	A	1	Total	O	S	0	0
			5	4	1		
6	A	1	Total	O	S	0	0
			5	4	1		
6	A	1	Total	O	S	0	0
			5	4	1		
6	A	1	Total	O	S	0	0
			5	4	1		
6	A	1	Total	O	S	0	0
			5	4	1		

- Molecule 7 is ACETATE ION (CCD ID: ACT) (formula: $C_2H_3O_2$).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
7	A	1	Total	C	O	0	0
			4	2	2		


- Molecule 8 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
8	A	335	Total	O	0	0
			335	335		


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: Agglutinin II




- Molecule 2: beta-D-xylopyranose-(1-2)-[alpha-D-mannopyranose-(1-3)][alpha-D-mannopyranose-(1-6)]beta-D-mannopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-[alpha-L-fucopyranose-(1-3)]2-acetamido-2-deoxy-beta-D-glucopyranose



- Molecule 3: alpha-L-fucopyranose-(1-3)-2-acetamido-2-deoxy-beta-D-glucopyranose



- Molecule 4: 2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose



- Molecule 5: α -L-fucopyranose-(1-3)-[2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)]2-acetamido-2-deoxy-beta-D-glucopyranose

Chain E:

100%

MAG1
FUC2
MAG3

4 Data and refinement statistics

Property	Value	Source
Space group	P 64 2 2	Depositor
Cell constants a, b, c, α , β , γ	120.20Å 120.20Å 177.34Å 90.00° 90.00° 120.00°	Depositor
Resolution (Å)	20.00 – 1.95 20.00 – 1.95	Depositor EDS
% Data completeness (in resolution range)	99.0 (20.00-1.95) 98.9 (20.00-1.95)	Depositor EDS
R_{merge}	0.08	Depositor
R_{sym}	0.08	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.96 (at 1.94Å)	Xtriage
Refinement program	REFMAC 5.2.0019	Depositor
R, R_{free}	0.171 , 0.194 0.181 , 0.201	Depositor DCC
R_{free} test set	2762 reflections (4.97%)	wwPDB-VP
Wilson B-factor (Å ²)	28.6	Xtriage
Anisotropy	0.051	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.36 , 52.8	EDS
L-test for twinning ²	$\langle L \rangle = 0.50$, $\langle L^2 \rangle = 0.33$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.92	EDS
Total number of atoms	2570	wwPDB-VP
Average B, all atoms (Å ²)	55.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 3.09% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: NAG, ACT, FUC, BMA, MAN, SO4, XYP

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	1.09	2/2063 (0.1%)	0.92	0/2810

All (2) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	128	ASN	CB-CG	5.17	1.65	1.52
1	A	184	CYS	N-CA	5.15	1.52	1.46

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2026	0	1962	11	0
2	B	80	0	60	0	0
3	C	24	0	22	0	0
4	D	28	0	25	0	0
5	E	38	0	34	0	0
6	A	35	0	0	0	0
7	A	4	0	3	0	0
8	A	335	0	0	4	0
All	All	2570	0	2106	11	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 3.

All (11) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:77:GLU:HG2	8:A:1082:HOH:O	1.95	0.66
1:A:250:GLN:NE2	8:A:1084:HOH:O	2.29	0.65
1:A:84:VAL:CG1	8:A:931:HOH:O	2.57	0.52
1:A:84:VAL:HG13	8:A:931:HOH:O	2.11	0.51
1:A:241:PHE:CG	1:A:242:PRO:HD2	2.47	0.50
1:A:196:ILE:HD12	1:A:238:ILE:HG22	1.95	0.48
1:A:85:PRO:HD2	1:A:89:SER:O	2.17	0.45
1:A:66:ASN:ND2	1:A:109:ILE:HD12	2.33	0.43
1:A:218:VAL:HG22	1:A:225:VAL:HG12	2.03	0.41
1:A:31:TRP:CD2	1:A:32:PRO:HD2	2.55	0.41
1:A:66:ASN:HD21	1:A:109:ILE:HD12	1.86	0.41

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	261/258 (101%)	257 (98%)	4 (2%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was

analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles
1	A	226/223 (101%)	223 (99%)	3 (1%)	61 58

All (3) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	83[A]	GLU
1	A	83[B]	GLU
1	A	195	LEU

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (2) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	66	ASN
1	A	254	GLN

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates ⓘ

14 monosaccharides are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
2	NAG	B	1	2,1	14,14,15	1.67	4 (28%)	17,19,21	1.75	6 (35%)
2	NAG	B	2	2	14,14,15	0.81	1 (7%)	17,19,21	1.07	1 (5%)

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
2	BMA	B	3	2	11,11,12	1.08	1 (9%)	15,15,17	1.67	4 (26%)
2	XYP	B	4	2	9,9,10	1.41	2 (22%)	10,12,14	1.09	1 (10%)
2	MAN	B	5	2	11,11,12	0.74	0	15,15,17	1.18	2 (13%)
2	MAN	B	6	2	11,11,12	0.65	0	15,15,17	1.05	1 (6%)
2	FUC	B	7	2	10,10,11	1.03	0	14,14,16	1.80	2 (14%)
3	NAG	C	1	3,1	14,14,15	0.77	1 (7%)	17,19,21	1.50	4 (23%)
3	FUC	C	2	3	10,10,11	0.71	0	14,14,16	1.25	2 (14%)
4	NAG	D	1	1,4	14,14,15	0.66	0	17,19,21	1.89	4 (23%)
4	NAG	D	2	4	14,14,15	0.58	0	17,19,21	1.53	3 (17%)
5	NAG	E	1	1,5	14,14,15	0.58	0	17,19,21	1.95	4 (23%)
5	FUC	E	2	5	10,10,11	0.61	0	14,14,16	1.56	3 (21%)
5	NAG	E	3	5	14,14,15	0.63	0	17,19,21	1.30	1 (5%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	NAG	B	1	2,1	-	0/6/23/26	0/1/1/1
2	NAG	B	2	2	-	0/6/23/26	0/1/1/1
2	BMA	B	3	2	-	0/2/19/22	0/1/1/1
2	XYP	B	4	2	-	-	0/1/1/1
2	MAN	B	5	2	-	2/2/19/22	0/1/1/1
2	MAN	B	6	2	-	0/2/19/22	0/1/1/1
2	FUC	B	7	2	-	-	0/1/1/1
3	NAG	C	1	3,1	-	2/6/23/26	0/1/1/1
3	FUC	C	2	3	-	-	0/1/1/1
4	NAG	D	1	1,4	-	1/6/23/26	0/1/1/1
4	NAG	D	2	4	-	0/6/23/26	0/1/1/1
5	NAG	E	1	1,5	-	1/6/23/26	0/1/1/1
5	FUC	E	2	5	-	-	0/1/1/1
5	NAG	E	3	5	-	4/6/23/26	0/1/1/1

All (9) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	B	1	NAG	C2-N2	3.45	1.52	1.46
2	B	4	XYP	O5-C1	-3.00	1.37	1.43

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	B	1	NAG	C8-C7	2.90	1.56	1.50
2	B	3	BMA	O5-C1	-2.54	1.39	1.43
2	B	2	NAG	O5-C1	-2.52	1.39	1.43
2	B	1	NAG	C4-C3	2.48	1.58	1.52
3	C	1	NAG	O5-C1	-2.22	1.40	1.43
2	B	4	XYP	C2-C3	2.20	1.55	1.52
2	B	1	NAG	C6-C5	2.01	1.58	1.51

All (38) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	B	7	FUC	C1-C2-C3	-5.36	101.85	109.64
4	D	1	NAG	O5-C1-C2	-4.76	103.93	111.29
5	E	1	NAG	C1-C2-N2	4.44	117.43	110.43
4	D	2	NAG	C1-O5-C5	4.11	117.70	112.19
5	E	1	NAG	C4-C3-C2	-3.96	105.21	111.02
2	B	1	NAG	O4-C4-C3	-3.90	101.17	110.38
5	E	1	NAG	C1-O5-C5	3.38	116.72	112.19
5	E	3	NAG	C2-N2-C7	3.25	127.26	122.90
5	E	1	NAG	O5-C1-C2	-3.23	106.29	111.29
2	B	3	BMA	C3-C4-C5	-3.17	104.48	110.23
4	D	1	NAG	C1-C2-N2	3.15	115.39	110.43
2	B	3	BMA	O3-C3-C4	-3.07	103.14	110.38
5	E	2	FUC	C1-O5-C5	2.97	119.97	112.97
5	E	2	FUC	O5-C5-C4	2.91	114.79	109.55
4	D	1	NAG	O3-C3-C2	-2.82	103.54	109.40
2	B	3	BMA	C1-C2-C3	-2.74	105.65	109.64
2	B	1	NAG	C3-C4-C5	-2.74	105.27	110.23
3	C	1	NAG	O3-C3-C2	2.68	114.98	109.40
3	C	2	FUC	O5-C5-C4	2.68	114.38	109.55
2	B	7	FUC	O2-C2-C3	-2.61	104.75	110.15
2	B	1	NAG	O3-C3-C4	-2.49	104.50	110.38
2	B	2	NAG	O3-C3-C2	-2.46	104.29	109.40
2	B	5	MAN	O3-C3-C4	2.43	116.11	110.38
4	D	2	NAG	O5-C1-C2	2.42	115.03	111.29
2	B	6	MAN	C1-O5-C5	2.39	115.39	112.19
4	D	2	NAG	O7-C7-C8	-2.34	117.89	122.05
2	B	5	MAN	C2-C3-C4	-2.33	106.76	110.86
2	B	1	NAG	O5-C1-C2	-2.32	107.69	111.29
5	E	2	FUC	C1-C2-C3	-2.29	106.31	109.64
2	B	1	NAG	O5-C5-C6	2.22	111.98	107.66
2	B	3	BMA	C6-C5-C4	-2.19	107.63	113.02

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	C	1	NAG	O5-C1-C2	-2.09	108.06	111.29
3	C	1	NAG	C1-C2-N2	-2.09	107.14	110.43
3	C	1	NAG	C4-C3-C2	-2.07	107.98	111.02
3	C	2	FUC	C1-O5-C5	2.07	117.84	112.97
2	B	4	XYP	O2-C2-C1	2.06	113.93	109.22
4	D	1	NAG	O3-C3-C4	2.05	115.20	110.38
2	B	1	NAG	O3-C3-C2	-2.04	105.16	109.40

There are no chirality outliers.


All (10) torsion outliers are listed below:


Mol	Chain	Res	Type	Atoms
5	E	1	NAG	C1-C2-N2-C7
5	E	3	NAG	O5-C5-C6-O6
5	E	3	NAG	C4-C5-C6-O6
2	B	5	MAN	C4-C5-C6-O6
3	C	1	NAG	C4-C5-C6-O6
3	C	1	NAG	O5-C5-C6-O6
2	B	5	MAN	O5-C5-C6-O6
4	D	1	NAG	C3-C2-N2-C7
5	E	3	NAG	C1-C2-N2-C7
5	E	3	NAG	C3-C2-N2-C7


There are no ring outliers.


No monomer is involved in short contacts.

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for oligosaccharide.









5.6 Ligand geometry [i](#)

8 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
6	SO4	A	905	-	4,4,4	0.31	0	6,6,6	0.34	0
6	SO4	A	903	-	4,4,4	0.55	0	6,6,6	0.55	0
6	SO4	A	904	-	4,4,4	0.40	0	6,6,6	0.36	0
6	SO4	A	906	-	4,4,4	0.35	0	6,6,6	0.37	0
6	SO4	A	907	-	4,4,4	0.27	0	6,6,6	0.79	0
6	SO4	A	901	-	4,4,4	0.20	0	6,6,6	0.78	0
6	SO4	A	902	-	4,4,4	0.25	0	6,6,6	0.50	0

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
7	ACT	A	910	-	3,3,3	0.92	0	3,3,3	1.50	0

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data ⓘ

6.1 Protein, DNA and RNA chains ⓘ

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	257/258 (99%)	1.81	96 (37%) 1 0	26, 53, 61, 68	6 (2%)

All (96) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	1	THR	6.1
1	A	257	PRO	5.6
1	A	184	CYS	4.8
1	A	236	ARG	4.8
1	A	116	ILE	3.8
1	A	154	TRP	3.8
1	A	201	CYS	3.7
1	A	73	SER	3.6
1	A	130	LYS	3.5
1	A	103[A]	ARG	3.5
1	A	119	ALA	3.5
1	A	213	SER	3.4
1	A	72	CYS	3.3
1	A	117	TYR	3.3
1	A	214	ASP	3.2
1	A	234	SER	3.2
1	A	132	ILE	3.1
1	A	203	GLY	3.1
1	A	133	VAL	3.1
1	A	215	GLY	3.1
1	A	235	LEU	3.0
1	A	254	GLN	3.0
1	A	218	VAL	2.9
1	A	171	GLY	2.9
1	A	216	ALA	2.9
1	A	204	LEU	2.8
1	A	209	TRP	2.8

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	A	129	VAL	2.8
1	A	255	VAL	2.7
1	A	211	PHE	2.7
1	A	161	THR	2.7
1	A	91	ILE	2.7
1	A	170	TYR	2.7
1	A	175	ILE	2.7
1	A	118	ALA	2.6
1	A	205	PRO	2.6
1	A	177	VAL	2.6
1	A	37	ARG	2.6
1	A	188	ASN	2.6
1	A	174	THR	2.6
1	A	98	VAL	2.6
1	A	104	ALA	2.6
1	A	106	SER	2.6
1	A	149	GLU	2.5
1	A	210	PHE	2.5
1	A	97	LEU	2.5
1	A	199	LEU	2.5
1	A	128	ASN	2.5
1	A	190	TYR	2.5
1	A	112	LEU	2.5
1	A	256	LEU	2.5
1	A	62[A]	ASN	2.5
1	A	252	VAL	2.5
1	A	189	GLY	2.4
1	A	195	LEU	2.4
1	A	86	ILE	2.4
1	A	45	SER	2.3
1	A	173	ARG	2.3
1	A	101	ALA	2.3
1	A	160	ALA	2.3
1	A	182	GLY	2.3
1	A	233	VAL	2.3
1	A	105	ALA	2.3
1	A	134	ALA	2.3
1	A	33	CYS	2.3
1	A	187	THR	2.3
1	A	123	TRP	2.2
1	A	36	GLN	2.2
1	A	111	LEU	2.2

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	A	168	ALA	2.2
1	A	172	ASP	2.2
1	A	109	ILE	2.2
1	A	100	THR	2.2
1	A	167	TRP	2.2
1	A	158	CYS	2.2
1	A	169	LEU	2.2
1	A	102	PRO	2.2
1	A	150	ASN	2.1
1	A	69	ILE	2.1
1	A	245	GLY	2.1
1	A	208	ARG	2.1
1	A	159	GLU	2.1
1	A	212	ASN	2.1
1	A	164	GLN	2.1
1	A	250	GLN	2.1
1	A	217	ILE	2.1
1	A	178	ASN	2.1
1	A	11	ASP	2.1
1	A	146	SER	2.1
1	A	77	GLU	2.1
1	A	200	LYS	2.1
1	A	84	VAL	2.1
1	A	14	CYS	2.0
1	A	31	TRP	2.0
1	A	83[A]	GLU	2.0
1	A	221	LYS	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q<0.9' lists the number of atoms with occupancy less than 0.9.


Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors($\text{\AA}^2$)	Q<0.9
5	FUC	E	2	10/11	0.18	0.32	106,108,109,110	0

Continued on next page...

Continued from previous page...




Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors($\text{\AA}^2$)	Q<0.9
5	NAG	E	3	14/15	0.37	0.28	105,107,108,109	0
3	FUC	C	2	10/11	0.51	0.28	86,92,94,94	0
2	MAN	B	5	11/12	0.58	0.30	79,83,85,86	0
4	NAG	D	2	14/15	0.65	0.23	79,81,86,86	0
5	NAG	E	1	14/15	0.68	0.23	85,92,100,101	0
2	MAN	B	6	11/12	0.78	0.19	64,68,75,78	0
3	NAG	C	1	14/15	0.83	0.17	58,65,76,79	0
4	NAG	D	1	14/15	0.87	0.17	46,56,64,73	0
2	BMA	B	3	11/12	0.90	0.15	47,56,63,68	0
2	FUC	B	7	10/11	0.94	0.14	43,46,50,53	0
2	NAG	B	2	14/15	0.94	0.15	38,42,47,47	0
2	XYP	B	4	9/10	0.95	0.16	38,43,45,46	0
2	NAG	B	1	14/15	0.96	0.15	34,38,43,46	0

The following is a graphical depiction of the model fit to experimental electron density for oligosaccharide. Each fit is shown from different orientation to approximate a three-dimensional view.






Electron density around Chain C:


$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)



Electron density around Chain D:

$2mF_o - DF_c$ (at 0.7 rmsd) in gray
 $mF_o - DF_c$ (at 3 rmsd) in purple (negative)
and green (positive)





6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors($\text{\AA}^2$)	Q<0.9
6	SO4	A	905	5/5	0.57	0.23	111,113,115,115	0
6	SO4	A	904	5/5	0.75	0.24	90,92,92,94	0
6	SO4	A	903	5/5	0.80	0.20	53,71,75,76	0
6	SO4	A	906	5/5	0.85	0.20	74,77,78,80	0
6	SO4	A	907	5/5	0.90	0.17	44,49,55,57	5
6	SO4	A	902	5/5	0.94	0.14	58,59,65,68	0
7	ACT	A	910	4/4	0.95	0.23	44,45,45,46	0
6	SO4	A	901	5/5	0.98	0.18	37,37,44,48	0

6.5 Other polymers ⓘ

There are no such residues in this entry.