



wwPDB X-ray Structure Validation Summary Report ⓘ

Mar 8, 2026 – 05:20 PM UTC

PDB ID : 2CF2 / pdb_00002cf2
Title : Architecture of mammalian fatty acid synthase
Authors : Maier, T.; Jenni, S.; Ban, N.
Deposited on : 2006-02-14
Resolution : 4.30 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4-5-2 with Phenix2.0
Xtriage (Phenix) : **NOT EXECUTED**
EDS : **NOT EXECUTED**
Percentile statistics : 20250101.v01 (using entries in the PDB archive January 1st 2025)
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 4.30 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
Clashscore	190562	1097 (4.70-3.90)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$.

Note EDS was not executed.

Mol	Chain	Length	Quality of chain
1	A	406	98% .
1	J	406	98% .
2	B	304	99% .
2	K	304	99% .
3	C	342	92% 8%
3	L	342	91% 9%
4	D	295	98% .
4	M	295	99% .
5	E	226	100%
5	N	226	100%

2 Entry composition

There are 5 unique types of molecules in this entry. The entry contains 3146 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called FATTY ACID SYNTHASE, KS DOMAIN.

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf	Trace
1	A	406	Total C 406 406	0	0	406
1	J	406	Total C 406 406	0	0	406

- Molecule 2 is a protein called FATTY ACID SYNTHASE, MAT DOMAIN.

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf	Trace
2	B	304	Total C 304 304	0	0	304
2	K	304	Total C 304 304	0	0	304

- Molecule 3 is a protein called FATTY ACID SYNTHASE, DH DOMAIN.

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf	Trace
3	C	342	Total C 342 342	0	0	342
3	L	342	Total C 342 342	0	0	342

- Molecule 4 is a protein called FATTY ACID SYNTHASE, ER DOMAIN.

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf	Trace
4	D	295	Total C 295 295	0	0	295
4	M	295	Total C 295 295	0	0	295

- Molecule 5 is a protein called FATTY ACID SYNTHASE, KR DOMAIN.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf	Trace
5	E	226	Total 226	C 226	0	0	226
5	N	226	Total 226	C 226	0	0	226

3 Residue-property plots

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

Note EDS was not executed.

- Molecule 1: FATTY ACID SYNTHASE, KS DOMAIN

Chain A:  98%



- Molecule 1: FATTY ACID SYNTHASE, KS DOMAIN

Chain J:  98%



- Molecule 2: FATTY ACID SYNTHASE, MAT DOMAIN

Chain B:  99%



- Molecule 2: FATTY ACID SYNTHASE, MAT DOMAIN

Chain K:  99%




- Molecule 3: FATTY ACID SYNTHASE, DH DOMAIN

Chain C:  92% 8%



- Molecule 3: FATTY ACID SYNTHASE, DH DOMAIN

Chain L:  91% 9%




- Molecule 4: FATTY ACID SYNTHASE, ER DOMAIN

Chain D:  98%




- Molecule 4: FATTY ACID SYNTHASE, ER DOMAIN

Chain M:  99%




- Molecule 5: FATTY ACID SYNTHASE, KR DOMAIN

Chain E:  100%



- Molecule 5: FATTY ACID SYNTHASE, KR DOMAIN

Chain N:  100%



4 Data and refinement statistics

Xtriage (Phenix) and EDS were not executed - this section is therefore incomplete.

Property	Value	Source
Space group	P 1 21 1	Depositor
Cell constants a, b, c, α , β , γ	96.10Å 245.00Å 134.10Å 90.00° 101.20° 90.00°	Depositor
Resolution (Å)	50.00 – 4.30	Depositor
% Data completeness (in resolution range)	(Not available) (50.00-4.30)	Depositor
R_{merge}	0.12	Depositor
R_{sym}	(Not available)	Depositor
Refinement program		Depositor
R, R_{free}	(Not available) , (Not available)	Depositor
Estimated twinning fraction	No twinning to report.	Xtriage
Total number of atoms	3146	wwPDB-VP
Average B, all atoms (Å ²)	60.0	wwPDB-VP

5 Model quality [i](#)

5.1 Standard geometry [i](#)

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

There are no protein, RNA or DNA chains available to summarize Z scores of covalent bonds and angles.

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	406	0	0	10	0
1	J	406	0	0	8	0
2	B	304	0	0	1	0
2	K	304	0	0	1	0
3	C	342	0	0	26	0
3	L	342	0	0	30	0
4	D	295	0	0	4	0
4	M	295	0	0	2	0
5	E	226	0	0	1	0
5	N	226	0	0	1	0
All	All	3146	0	0	49	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 16.

The worst 5 of 49 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:121:ARG:CA	3:C:1005:GLU:CA	1.75	1.60
1:A:43:GLY:CA	3:L:1020:LEU:CA	1.83	1.55
1:J:126:LEU:CA	3:L:1014:ALA:CA	1.77	1.53
3:L:2137:ARG:CA	4:M:52:ARG:CA	1.87	1.50
3:C:1047:GLY:CA	3:L:1127:ARG:CA	1.92	1.47

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

There are no protein backbone outliers to report in this entry.

5.3.2 Protein sidechains [i](#)

There are no protein residues with a non-rotameric sidechain to report in this entry.

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

There are no ligands in this entry.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues ⓘ

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

EDS was not executed - this section is therefore empty.

6.2 Non-standard residues in protein, DNA, RNA chains

EDS was not executed - this section is therefore empty.

6.3 Carbohydrates

EDS was not executed - this section is therefore empty.

6.4 Ligands

EDS was not executed - this section is therefore empty.

6.5 Other polymers

EDS was not executed - this section is therefore empty.