



wwPDB X-ray Structure Validation Summary Report ⓘ

Mar 9, 2026 – 07:31 PM UTC

PDB ID : 6COX / pdb_00006cox
Title : CYCLOOXYGENASE-2 (PROSTAGLANDIN SYNTHASE-2) COM-
PLEXED WITH A SELECTIVE INHIBITOR, SC-558 IN I222 SPACE
GROUP
Authors : Kurumbail, R.; Stallings, W.
Deposited on : 1996-12-18
Resolution : 2.80 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4-5-2 with Phenix2.0
Mogul : 2022.3.0, CSD as543be (2022)
Xtriage (Phenix) : **NOT EXECUTED**
EDS : **NOT EXECUTED**
Buster-report : wwPDB partial adaption of 1.1.7 (2018)
Percentile statistics : 20250101.v01 (using entries in the PDB archive January 1st 2025)
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.80 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
Clashscore	190562	4276 (2.80-2.80)
Ramachandran outliers	187476	4196 (2.80-2.80)
Sidechain outliers	187428	4198 (2.80-2.80)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$

Note EDS was not executed.

Mol	Chain	Length	Quality of chain
1	A	587	35% 43% 14% • 6%
1	B	587	36% 42% 14% • 6%

2 Entry composition [i](#)

There are 4 unique types of molecules in this entry. The entry contains 9168 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.


- Molecule 1 is a protein called CYCLOOXYGENASE-2.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	552	Total	C	N	O	S	0	0	0
			4473	2886	748	814	25			
1	B	552	Total	C	N	O	S	0	0	0
			4473	2886	748	814	25			

There are 4 discrepancies between the modelled and reference sequences:


Chain	Residue	Modelled	Actual	Comment	Reference
A	310	GLN	ASN	conflict	UNP Q05769
A	333	LYS	ARG	conflict	UNP Q05769
B	310	GLN	ASN	conflict	UNP Q05769
B	333	LYS	ARG	conflict	UNP Q05769

- Molecule 2 is 2-acetamido-2-deoxy-beta-D-glucopyranose (CCD ID: NAG) (formula: $C_8H_{15}NO_6$).




Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
2	A	1	Total	C	N	O	0	0
			14	8	1	5		
2	A	1	Total	C	N	O	0	0
			14	8	1	5		
2	A	1	Total	C	N	O	0	0
			14	8	1	5		
2	B	1	Total	C	N	O	0	0
			14	8	1	5		
2	B	1	Total	C	N	O	0	0
			14	8	1	5		
2	B	1	Total	C	N	O	0	0
			14	8	1	5		

- Molecule 3 is PROTOPORPHYRIN IX CONTAINING FE (CCD ID: HEM) (formula: $C_{34}H_{32}FeN_4O_4$).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
3	A	1	Total 43	C 34	Fe 1	N 4	O 4	0	0
3	B	1	Total 43	C 34	Fe 1	N 4	O 4	0	0

- Molecule 4 is 1-PHENYLSULFONAMIDE-3-TRIFLUOROMETHYL-5-PARABROMOPH ENYLPYRAZOLE (CCD ID: S58) (formula: $C_{16}H_{11}BrF_3N_3O_2S$).




Mol	Chain	Residues	Atoms							ZeroOcc	AltConf
			Total	Br	C	F	N	O	S		
4	A	1	26	1	16	3	3	2	1	0	0
4	B	1	26	1	16	3	3	2	1	0	0

3 Residue-property plots


These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

Note EDS was not executed.

• Molecule 1: CYCLOOXYGENASE-2



• Molecule 1: CYCLOOXYGENASE-2





4 Data and refinement statistics

Xtriage (Phenix) and EDS were not executed - this section is therefore incomplete.

Property	Value	Source
Space group	I 2 2 2	Depositor
Cell constants a, b, c, α , β , γ	181.17Å 132.81Å 122.74Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	8.00 – 2.80	Depositor
% Data completeness (in resolution range)	88.5 (8.00-2.80)	Depositor
R_{merge}	0.09	Depositor
R_{sym}	0.09	Depositor
Refinement program	X-PLOR 3.1	Depositor
R, R_{free}	0.220 , 0.309	Depositor
Estimated twinning fraction	No twinning to report.	Xtriage
Total number of atoms	9168	wwPDB-VP
Average B, all atoms (Å ²)	20.0	wwPDB-VP

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: HEM, NAG, S58

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.80	1/4600 (0.0%)	1.26	55/6237 (0.9%)
1	B	0.79	2/4600 (0.0%)	1.27	46/6237 (0.7%)
All	All	0.80	3/9200 (0.0%)	1.27	101/12474 (0.8%)

All (3) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	B	279	ILE	CA-CB	6.07	1.58	1.54
1	A	197	MET	SD-CE	-5.78	1.65	1.79
1	B	388	HIS	CG-CD2	5.49	1.41	1.35

The worst 5 of 101 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	B	269	THR	N-CA-C	-9.50	97.72	110.55
1	A	319	GLU	N-CA-C	-9.46	102.55	114.56
1	B	137	LYS	N-CA-C	9.29	122.36	111.02
1	A	137	LYS	N-CA-C	9.25	122.30	111.02
1	B	427	THR	N-CA-C	-9.04	101.43	111.28

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	4473	0	4375	330	0
1	B	4473	0	4375	322	0
2	A	42	0	39	1	0
2	B	42	0	39	0	0
3	A	43	0	30	2	0
3	B	43	0	30	1	0
4	A	26	0	11	3	0
4	B	26	0	11	2	0
All	All	9168	0	8910	634	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 35.

The worst 5 of 634 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:283:LEU:HD22	1:A:411:ASN:HB2	1.32	1.11
1:B:280:PRO:HD2	1:B:283:LEU:HD23	1.35	1.09
1:A:279:ILE:HG23	1:A:283:LEU:HG	1.32	1.08
1:B:279:ILE:HG23	1:B:283:LEU:HG	1.37	1.07
1:A:458:MET:HE3	1:A:460:TYR:HE1	1.17	1.05

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
1	A	550/587 (94%)	435 (79%)	84 (15%)	31 (6%)	1 4
1	B	550/587 (94%)	431 (78%)	86 (16%)	33 (6%)	1 3
All	All	1100/1174 (94%)	866 (79%)	170 (16%)	64 (6%)	1 4

5 of 64 Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	138	SER
1	A	348	TYR
1	A	398	GLU
1	A	510	GLU
1	A	514	PRO

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	493/525 (94%)	428 (87%)	65 (13%)	4	14
1	B	493/525 (94%)	430 (87%)	63 (13%)	4	15
All	All	986/1050 (94%)	858 (87%)	128 (13%)	4	14

5 of 128 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	B	444	VAL
1	B	484	GLU
1	A	405	LYS
1	A	396	ASN
1	B	509	VAL

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 22 such sidechains are listed below:

Mol	Chain	Res	Type
1	B	320	HIS
1	B	369	GLN
1	B	351	HIS
1	B	417	HIS
1	A	351	HIS

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

10 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# $ Z > 2$	Counts	RMSZ	# $ Z > 2$
4	S58	B	701	-	28,28,28	2.33	7 (25%)	43,43,43	2.29	11 (25%)
2	NAG	B	671	1	14,14,15	0.58	0	17,19,21	0.94	1 (5%)
2	NAG	B	661	1	14,14,15	0.57	0	17,19,21	0.59	0
2	NAG	B	681	1	14,14,15	0.71	0	17,19,21	0.86	0
3	HEM	B	682	1	50,50,50	1.21	4 (8%)	67,82,82	0.94	2 (2%)
4	S58	A	701	-	28,28,28	2.41	7 (25%)	43,43,43	2.45	13 (30%)
2	NAG	A	671	1	14,14,15	0.42	0	17,19,21	0.85	1 (5%)
2	NAG	A	681	1	14,14,15	0.33	0	17,19,21	0.98	1 (5%)
3	HEM	A	682	1	50,50,50	1.21	4 (8%)	67,82,82	0.86	0
2	NAG	A	661	1	14,14,15	0.67	0	17,19,21	0.54	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
4	S58	B	701	-	-	2/20/20/20	0/3/3/3
2	NAG	B	671	1	-	2/6/23/26	0/1/1/1
2	NAG	B	661	1	-	2/6/23/26	0/1/1/1

Continued on next page...

Continued from previous page...

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	NAG	B	681	1	-	0/6/23/26	0/1/1/1
3	HEM	B	682	1	-	7/14/54/54	-
4	S58	A	701	-	-	2/20/20/20	0/3/3/3
2	NAG	A	671	1	-	2/6/23/26	0/1/1/1
2	NAG	A	681	1	-	0/6/23/26	0/1/1/1
3	HEM	A	682	1	-	7/14/54/54	-
2	NAG	A	661	1	-	2/6/23/26	0/1/1/1

The worst 5 of 22 bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
4	A	701	S58	S1-N3	8.19	1.76	1.60
4	B	701	S58	S1-N3	7.40	1.74	1.60
4	B	701	S58	C1-C3	5.07	1.47	1.39
4	A	701	S58	C1-C3	4.83	1.47	1.39
4	B	701	S58	C8-S1	4.25	1.83	1.77

The worst 5 of 29 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
4	B	701	S58	C2-N1-N2	-7.43	107.47	112.19
4	A	701	S58	C2-N1-N2	-7.36	107.52	112.19
4	A	701	S58	O2-S1-O1	-5.65	110.11	118.80
4	B	701	S58	C4-C3-N2	5.29	126.29	119.20
4	A	701	S58	C4-C3-N2	5.11	126.05	119.20

There are no chirality outliers.

5 of 26 torsion outliers are listed below:


Mol	Chain	Res	Type	Atoms
3	A	682	HEM	C2B-C3B-CAB-CBB
3	B	682	HEM	C2A-CAA-CBA-CGA
3	B	682	HEM	C2B-C3B-CAB-CBB
3	A	682	HEM	C2A-CAA-CBA-CGA
2	A	671	NAG	O5-C5-C6-O6


There are no ring outliers.


5 monomers are involved in 9 short contacts:


Mol	Chain	Res	Type	Clashes	Symm-Clashes
4	B	701	S58	2	0
3	B	682	HEM	1	0
4	A	701	S58	3	0
3	A	682	HEM	2	0
2	A	661	NAG	1	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.









5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

EDS was not executed - this section is therefore empty.

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

EDS was not executed - this section is therefore empty.

6.3 Carbohydrates [i](#)

EDS was not executed - this section is therefore empty.

6.4 Ligands [i](#)

EDS was not executed - this section is therefore empty.

6.5 Other polymers [i](#)

EDS was not executed - this section is therefore empty.