



wwPDB X-ray Structure Validation Summary Report ⓘ

Mar 8, 2026 – 03:46 AM UTC

PDB ID : 5D93 / pdb_00005d93
Title : Oxidoreductase Fragment of Mouse QSOX1 in Complex with a FAb Fragment from a Mouse QSOX1-Specific Antibody
Authors : Grossman, I.; Fass, D.
Deposited on : 2015-08-18
Resolution : 2.20 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4-5-2 with Phenix2.0
Xtriage (Phenix)	:	2.0
EDS	:	3.0
Percentile statistics	:	20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4	:	9.0.010 (Gargrove)
Density-Fitness	:	1.0.12
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION



The reported resolution of this entry is 2.20 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.




Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	6164 (2.20-2.20)
Clashscore	190562	6851 (2.20-2.20)
Ramachandran outliers	187476	6768 (2.20-2.20)
Sidechain outliers	187428	6769 (2.20-2.20)
RSRZ outliers	180081	6166 (2.20-2.20)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	244	 85% 12% ..
1	D	244	 83% 12% ..
2	C	218	 86% 8% 5%
2	F	218	 94% 5% .
3	B	213	 95% .

Continued on next page...

Continued from previous page...

Mol	Chain	Length	Quality of chain
3	E	213	 97%

2 Entry composition

There are 4 unique types of molecules in this entry. The entry contains 10907 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Sulfhydryl oxidase 1.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	240	Total	C	N	O	S	0	0	0
			1868	1190	319	351	8			
1	D	235	Total	C	N	O	S	0	0	0
			1828	1168	311	342	7			

There are 8 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	32	GLY	-	expression tag	UNP Q8BND5
A	33	SER	-	expression tag	UNP Q8BND5
A	34	HIS	-	expression tag	UNP Q8BND5
A	35	MET	-	expression tag	UNP Q8BND5
D	32	GLY	-	expression tag	UNP Q8BND5
D	33	SER	-	expression tag	UNP Q8BND5
D	34	HIS	-	expression tag	UNP Q8BND5
D	35	MET	-	expression tag	UNP Q8BND5

- Molecule 2 is a protein called Heavy Chain of Fab Fragment from a Mouse QSOX1-Specific Antibody.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
2	C	207	Total	C	N	O	S	0	0	0
			1578	1009	255	309	5			
2	F	216	Total	C	N	O	S	0	0	0
			1638	1043	266	324	5			

- Molecule 3 is a protein called Light Chain of Fab Fragment from a Mouse QSOX1-Specific Antibody.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
3	B	212	Total	C	N	O	S	0	0	0
			1622	1018	268	329	7			


Continued on next page...

Continued from previous page...

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
3	E	212	Total	C	N	O	S	0	0	0
			1622	1018	268	329	7			


- Molecule 4 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	A	153	Total	O	0	0
			153	153		
4	C	120	Total	O	0	0
			120	120		
4	B	126	Total	O	0	0
			126	126		
4	D	85	Total	O	0	0
			85	85		
4	F	138	Total	O	0	0
			138	138		
4	E	129	Total	O	0	0
			129	129		

Chain B:  95% .



- Molecule 3: Light Chain of Fab Fragment from a Mouse QSOX1-Specific Antibody

Chain E:  97% .



4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, α , β , γ	65.54Å 112.72Å 193.42Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	23.20 – 2.20 23.20 – 2.20	Depositor EDS
% Data completeness (in resolution range)	98.9 (23.20-2.20) 98.8 (23.20-2.20)	Depositor EDS
R_{merge}	0.09	Depositor
R_{sym}	0.09	Depositor
$\langle I/\sigma(I) \rangle$ ¹	2.27 (at 2.19Å)	Xtriage
Refinement program	PHENIX 1.9_1692	Depositor
R, R_{free}	0.174 , 0.236 0.182 , 0.184	Depositor DCC
R_{free} test set	3627 reflections (4.93%)	wwPDB-VP
Wilson B-factor (Å ²)	30.0	Xtriage
Anisotropy	0.352	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.34 , 31.4	EDS
L-test for twinning ²	$\langle L \rangle = 0.48$, $\langle L^2 \rangle = 0.31$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.95	EDS
Total number of atoms	10907	wwPDB-VP
Average B, all atoms (Å ²)	32.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 3.80% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.52	0/1915	0.83	2/2613 (0.1%)
1	D	0.49	0/1874	0.76	0/2557
2	C	0.50	0/1622	0.79	0/2215
2	F	0.50	0/1684	0.79	1/2298 (0.0%)
3	B	0.47	0/1662	0.80	0/2259
3	E	0.47	0/1662	0.77	0/2259
All	All	0.49	0/10419	0.79	3/14201 (0.0%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
2	C	0	1
2	F	0	1
All	All	0	2

There are no bond length outliers.

All (3) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed($^{\circ}$)	Ideal($^{\circ}$)
2	F	134	SER	N-CA-C	6.54	118.07	111.07
1	A	54	ARG	CA-C-N	-5.68	114.25	119.82
1	A	54	ARG	C-N-CA	-5.68	114.25	119.82

There are no chirality outliers.

All (2) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
2	C	99	SER	Peptide
2	F	99	SER	Peptide

5.2 Too-close contacts ⓘ

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1868	0	1818	22	0
1	D	1828	0	1781	22	0
2	C	1578	0	1524	10	0
2	F	1638	0	1592	7	0
3	B	1622	0	1565	6	0
3	E	1622	0	1565	2	0
4	A	153	0	0	5	5
4	B	126	0	0	0	2
4	C	120	0	0	0	2
4	D	85	0	0	2	1
4	E	129	0	0	0	0
4	F	138	0	0	2	0
All	All	10907	0	9845	66	5

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 3.

The worst 5 of 66 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:136:GLY:O	4:A:301:HOH:O	2.02	0.77
2:C:212:ARG:NH2	2:C:214:GLU:OE2	2.21	0.72
2:C:197:ILE:HG12	2:C:212:ARG:HG2	1.75	0.68
1:D:65:ALA:HB3	1:D:99:LEU:HD22	1.75	0.67
1:A:172:GLU:OE1	4:A:302:HOH:O	2.15	0.64

All (5) symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
4:A:301:HOH:O	4:B:306:HOH:O[4_475]	1.71	0.49
4:A:444:HOH:O	4:C:301:HOH:O[4_475]	1.77	0.43
4:A:304:HOH:O	4:C:352:HOH:O[4_475]	1.86	0.34
4:A:425:HOH:O	4:D:334:HOH:O[2_474]	2.04	0.16

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
4:A:444:HOH:O	4:B:355:HOH:O[4_475]	2.05	0.15

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	238/244 (98%)	233 (98%)	5 (2%)	0	100	100
1	D	231/244 (95%)	223 (96%)	8 (4%)	0	100	100
2	C	203/218 (93%)	194 (96%)	9 (4%)	0	100	100
2	F	214/218 (98%)	206 (96%)	8 (4%)	0	100	100
3	B	210/213 (99%)	203 (97%)	7 (3%)	0	100	100
3	E	210/213 (99%)	203 (97%)	7 (3%)	0	100	100
All	All	1306/1350 (97%)	1262 (97%)	44 (3%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	203/207 (98%)	198 (98%)	5 (2%)	42	56
1	D	198/207 (96%)	193 (98%)	5 (2%)	42	56
2	C	175/186 (94%)	173 (99%)	2 (1%)	65	79

Continued on next page...

Continued from previous page...

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
2	F	184/186 (99%)	181 (98%)	3 (2%)	55	71
3	B	184/185 (100%)	183 (100%)	1 (0%)	81	90
3	E	184/185 (100%)	183 (100%)	1 (0%)	81	90
All	All	1128/1156 (98%)	1111 (98%)	17 (2%)	57	73

5 of 17 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
2	F	152	VAL
3	E	1	GLN
3	B	1	GLN
1	D	110	SER
1	D	157	LEU

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (3) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	117	ASN
3	B	159	GLN
1	D	177	ASN

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates ⓘ

There are no oligosaccharides in this entry.

5.6 Ligand geometry ⓘ

There are no ligands in this entry.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	240/244 (98%)	-0.41	1 (0%) 88 87	21, 29, 43, 52	0
1	D	235/244 (96%)	-0.19	2 (0%) 81 79	24, 35, 55, 63	0
2	C	207/218 (94%)	-0.33	5 (2%) 59 56	19, 30, 56, 77	0
2	F	216/218 (99%)	-0.54	2 (0%) 81 79	21, 28, 45, 59	0
3	B	212/213 (99%)	-0.35	0 100 100	19, 31, 53, 60	0
3	E	212/213 (99%)	-0.41	0 100 100	24, 31, 45, 57	0
All	All	1322/1350 (97%)	-0.37	10 (0%) 82 80	19, 30, 51, 77	0

The worst 5 of 10 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
2	C	193	THR	3.0
1	D	164	TRP	2.9
2	C	138	ALA	2.7
2	C	129	SER	2.6
1	A	36	SER	2.4

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.4 Ligands [i](#)

There are no ligands in this entry.

6.5 Other polymers [i](#)

There are no such residues in this entry.