



# wwPDB X-ray Structure Validation Summary Report ⓘ

Mar 7, 2026 – 04:27 AM UTC

PDB ID : 3DC2 / pdb\_00003dc2  
Title : Crystal structure of serine bound D-3-phosphoglycerate dehydrogenase from Mycobacterium tuberculosis  
Authors : Dey, S.; Sacchettini, J.C.  
Deposited on : 2008-06-03  
Resolution : 2.70 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4-5-2 with Phenix2.0  
Mogul : 2022.3.0, CSD as543be (2022)  
Xtriage (Phenix) : 2.0  
EDS : 3.0  
Percentile statistics : 20250101.v01 (using entries in the PDB archive January 1st 2025)  
CCP4 : 9.0.010 (Gargrove)  
Density-Fitness : 1.0.12  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : 2.49


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

*X-RAY DIFFRACTION*

The reported resolution of this entry is 2.70 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	180053	3538 (2.70-2.70)
Clashscore	190562	3843 (2.70-2.70)
Ramachandran outliers	187476	3778 (2.70-2.70)
Sidechain outliers	187428	3778 (2.70-2.70)
RSRZ outliers	180081	3538 (2.70-2.70)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	529	 5% 59% 33% 6% ..
1	B	529	 8% 56% 35% 8% ..

## 2 Entry composition [i](#)

There are 4 unique types of molecules in this entry. The entry contains 7761 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.


- Molecule 1 is a protein called D-3-phosphoglycerate dehydrogenase.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	526	Total 3830	C 2413	N 672	O 743	S 2	0	0	0
1	B	523	Total 3810	C 2401	N 669	O 738	S 2	0	0	0

There are 4 discrepancies between the modelled and reference sequences:


Chain	Residue	Modelled	Actual	Comment	Reference
A	1	MET	-	expression tag	UNP P0A544
A	2	VAL	-	expression tag	UNP P0A544
B	1	MET	-	expression tag	UNP P0A544
B	2	VAL	-	expression tag	UNP P0A544

- Molecule 2 is SERINE (CCD ID: SER) (formula: C<sub>3</sub>H<sub>7</sub>NO<sub>3</sub>).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
2	A	1	Total	C	N	O	0	0
			7	3	1	3		
2	B	1	Total	C	N	O	0	0
			7	3	1	3		

- Molecule 3 is L(+)-TARTARIC ACID (CCD ID: TLA) (formula: C<sub>4</sub>H<sub>6</sub>O<sub>6</sub>).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
3	A	1	Total	C	O	0	0
			10	4	6		
3	B	1	Total	C	O	0	0
			10	4	6		
3	B	1	Total	C	O	0	0
			10	4	6		


- Molecule 4 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	A	40	Total	O	0	0
			40	40		
4	B	37	Total	O	0	0
			37	37		


### 3 Residue-property plots [i](#)


These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: D-3-phosphoglycerate dehydrogenase



- Molecule 1: D-3-phosphoglycerate dehydrogenase





## 4 Data and refinement statistics

Property	Value	Source
Space group	P 65 2 2	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	165.19Å 165.19Å 218.89Å 90.00° 90.00° 120.00°	Depositor
Resolution (Å)	46.57 – 2.70 46.57 – 2.70	Depositor EDS
% Data completeness (in resolution range)	99.2 (46.57-2.70) 99.1 (46.57-2.70)	Depositor EDS
$R_{merge}$	0.04	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	2.65 (at 2.69Å)	Xtrriage
Refinement program	REFMAC 5.2.0019	Depositor
R, $R_{free}$	0.222 , 0.263 0.210 , 0.251	Depositor DCC
$R_{free}$ test set	2452 reflections (5.01%)	wwPDB-VP
Wilson B-factor (Å <sup>2</sup> )	67.7	Xtrriage
Anisotropy	0.632	Xtrriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.30 , 66.0	EDS
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.46$ , $\langle L^2 \rangle = 0.29$	Xtrriage
Estimated twinning fraction	No twinning to report.	Xtrriage
$F_o, F_c$ correlation	0.92	EDS
Total number of atoms	7761	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	67.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 6.10% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

## 5 Model quality [i](#)

### 5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: TLA

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z  >5	RMSZ	# Z  >5
1	A	1.23	10/3884 (0.3%)	1.23	16/5308 (0.3%)
1	B	1.16	21/3863 (0.5%)	1.23	26/5277 (0.5%)
All	All	1.20	31/7747 (0.4%)	1.23	42/10585 (0.4%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	B	0	1

The worst 5 of 31 bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	128	HIS	CE1-NE2	36.00	1.68	1.32
1	A	128	HIS	CG-ND1	20.74	1.61	1.38
1	B	250	HIS	CE1-NE2	19.03	1.51	1.32
1	B	250	HIS	CG-CD2	13.34	1.50	1.35
1	A	446	ARG	CZ-NH1	11.25	1.48	1.32

The worst 5 of 42 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	128	HIS	ND1-CG-CD2	10.30	116.41	106.10
1	B	433	GLY	CA-C-N	8.94	128.53	119.24
1	B	433	GLY	C-N-CA	8.94	128.53	119.24
1	B	210	THR	N-CA-C	-8.78	101.64	111.82
1	B	223	LYS	CA-C-N	-8.01	111.75	119.76

There are no chirality outliers.

All (1) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	B	209	GLU	Peptide

## 5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	3830	0	3916	169	0
1	B	3810	0	3897	186	0
2	A	7	0	4	0	0
2	B	7	0	4	0	0
3	A	10	0	4	0	0
3	B	20	0	8	0	0
4	A	40	0	0	6	0
4	B	37	0	0	5	0
All	All	7761	0	7833	347	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 22.

The worst 5 of 347 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:66:LYS:CE	1:B:66:LYS:NZ	1.73	1.49
1:A:128:HIS:CE1	1:A:128:HIS:NE2	1.68	1.45
1:A:285:THR:HG22	1:A:288:ALA:H	1.06	1.20
1:A:332:ARG:HG2	1:A:332:ARG:HH21	1.18	1.08
1:B:285:THR:HG22	1:B:288:ALA:H	1.16	1.07

There are no symmetry-related clashes.

## 5.3 Torsion angles [i](#)

### 5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	524/529 (99%)	471 (90%)	47 (9%)	6 (1%)	11	29
1	B	519/529 (98%)	466 (90%)	50 (10%)	3 (1%)	21	44
All	All	1043/1058 (99%)	937 (90%)	97 (9%)	9 (1%)	14	35

5 of 9 Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	489	GLU
1	A	491	ALA
1	A	12	LYS
1	A	22	GLY
1	A	128	HIS

### 5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	396/399 (99%)	339 (86%)	57 (14%)	3	8
1	B	394/399 (99%)	341 (86%)	53 (14%)	4	9
All	All	790/798 (99%)	680 (86%)	110 (14%)	3	9

5 of 110 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	B	4	LEU

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type
1	B	134	SER
1	B	528	LEU
1	B	442	GLN
1	B	18	VAL

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 14 such sidechains are listed below:

Mol	Chain	Res	Type
1	B	99	ASN
1	B	128	HIS
1	B	456	ASN
1	B	321	ASN
1	B	352	GLN

### 5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

### 5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

### 5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

### 5.6 Ligand geometry [i](#)

5 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z  > 2	Counts	RMSZ	# Z  > 2
3	TLA	B	700	-	9,9,9	0.83	0	12,12,12	1.35	3 (25%)
2	SER	A	600	-	4,6,6	1.03	0	2,7,7	2.23	1 (50%)
2	SER	B	600	-	4,6,6	1.05	0	2,7,7	2.47	1 (50%)
3	TLA	A	700	-	9,9,9	0.83	0	12,12,12	1.56	3 (25%)
3	TLA	B	701	-	9,9,9	0.74	0	12,12,12	1.17	1 (8%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	TLA	B	700	-	-	0/12/12/12	-
2	SER	A	600	-	-	1/6/6/6	-
2	SER	B	600	-	-	0/6/6/6	-
3	TLA	A	700	-	-	4/12/12/12	-
3	TLA	B	701	-	-	7/12/12/12	-

There are no bond length outliers.

The worst 5 of 9 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	B	600	SER	OXT-C-O	-3.33	116.52	124.08
2	A	600	SER	OXT-C-O	-3.12	117.00	124.08
3	A	700	TLA	O41-C4-C3	3.10	121.92	113.31
3	A	700	TLA	O41-C4-O4	-2.40	118.62	124.08
3	A	700	TLA	O11-C1-C2	2.34	119.80	113.31

There are no chirality outliers.

5 of 12 torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
3	A	700	TLA	C2-C3-C4-O41
3	B	701	TLA	C2-C3-C4-O4
3	B	701	TLA	C2-C3-C4-O41
3	B	701	TLA	O3-C3-C4-O4
3	B	701	TLA	O3-C3-C4-O41

There are no ring outliers.

No monomer is involved in short contacts.

## 5.7 Other polymers [i](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

## 6 Fit of model and data [i](#)

### 6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	526/529 (99%)	0.39	24 (4%) 37 34	46, 65, 83, 101	8 (1%)
1	B	523/529 (98%)	0.63	44 (8%) 17 14	16, 67, 81, 99	18 (3%)
All	All	1049/1058 (99%)	0.51	68 (6%) 25 22	16, 66, 82, 101	26 (2%)

The worst 5 of 68 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	B	395	GLY	10.8
1	B	396	VAL	8.2
1	B	394	ARG	7.9
1	B	385	VAL	5.6
1	B	384	PHE	5.0

### 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

### 6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

### 6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled ‘Q < 0.9’ lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors( $\text{\AA}^2$ )	Q<0.9
3	TLA	B	701	10/10	0.78	0.18	63,79,81,81	10
3	TLA	A	700	10/10	0.79	0.11	82,94,101,102	0
3	TLA	B	700	10/10	0.79	0.14	81,93,100,100	0
2	SER	A	600	7/7	0.79	0.25	80,80,84,85	0
2	SER	B	600	7/7	0.90	0.19	72,82,90,90	0

## 6.5 Other polymers [i](#)

There are no such residues in this entry.