



wwPDB X-ray Structure Validation Summary Report ⓘ

Mar 8, 2026 – 01:44 PM UTC

PDB ID : 8DLA / pdb_00008dla
Title : ClpP2 from Chlamydia trachomatis bound by MAS1-12
Authors : Azadmanesh, J.; Struble, L.R.; Seleem, M.A.; Ouellette, S.; Conda-Sheridan, M.; Borgstahl, G.E.O.
Deposited on : 2022-07-07
Resolution : 2.66 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4-5-2 with Phenix2.0
Mogul	:	2022.3.0, CSD as543be (2022)
Xtriage (Phenix)	:	2.0
EDS	:	3.0
Buster-report	:	wwPDB partial adaption of 1.1.7 (2018)
Percentile statistics	:	20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4	:	9.0.010 (Gargrove)
Density-Fitness	:	1.0.12
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION


The reported resolution of this entry is 2.66 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	1110 (2.66-2.66)
Clashscore	190562	1141 (2.66-2.66)
Ramachandran outliers	187476	1126 (2.66-2.66)
Sidechain outliers	187428	1126 (2.66-2.66)
RSRZ outliers	180081	1110 (2.66-2.66)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	203	
1	B	203	
1	C	203	
1	D	203	
1	E	203	

Continued on next page...

Continued from previous page...

Mol	Chain	Length	Quality of chain
1	F	203	 69%28%..
1	G	203	 63%33%..
1	H	203	 67%29%..
1	I	203	 67%29%.
1	J	203	 64%33%.
1	K	203	 64%35%.
1	L	203	 68%28%.
1	M	203	 67%28%.
1	N	203	 68%28%..

2 Entry composition

There are 5 unique types of molecules in this entry. The entry contains 22282 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called ATP-dependent Clp protease proteolytic subunit 2.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	203	Total	C	N	O	S	0	0	0
			1542	974	256	301	11			
1	B	203	Total	C	N	O	S	0	0	0
			1542	974	256	301	11			
1	C	203	Total	C	N	O	S	0	0	0
			1542	974	256	301	11			
1	D	203	Total	C	N	O	S	0	0	0
			1542	974	256	301	11			
1	E	203	Total	C	N	O	S	0	0	0
			1542	974	256	301	11			
1	F	203	Total	C	N	O	S	0	0	0
			1542	974	256	301	11			
1	G	203	Total	C	N	O	S	0	0	0
			1542	974	256	301	11			
1	H	203	Total	C	N	O	S	0	0	0
			1542	974	256	301	11			
1	I	203	Total	C	N	O	S	0	0	0
			1542	974	256	301	11			
1	J	203	Total	C	N	O	S	0	0	0
			1542	974	256	301	11			
1	K	203	Total	C	N	O	S	0	0	0
			1542	974	256	301	11			
1	L	203	Total	C	N	O	S	0	0	0
			1542	974	256	301	11			
1	M	203	Total	C	N	O	S	0	0	0
			1542	974	256	301	11			
1	N	203	Total	C	N	O	S	0	0	0
			1542	974	256	301	11			

- Molecule 2 is 1-{4-[(4-chlorophenyl)methyl]piperazin-1-yl}-2-methyl-2-[5-(trifluoromethyl)pyridine-2-sulfonyl]propan-1-one (CCD ID: UER) (formula: C₂₁H₂₃ClF₃N₃O₃S) (labeled as "Ligand of Interest" by depositor).



Mol	Chain	Residues	Atoms							ZeroOcc	AltConf
2	A	1	Total 32	C 21	Cl 1	F 3	N 3	O 3	S 1	0	0
2	A	1	Total 32	C 21	Cl 1	F 3	N 3	O 3	S 1	0	0
2	B	1	Total 32	C 21	Cl 1	F 3	N 3	O 3	S 1	0	0
2	D	1	Total 32	C 21	Cl 1	F 3	N 3	O 3	S 1	0	0
2	G	1	Total 32	C 21	Cl 1	F 3	N 3	O 3	S 1	0	0
2	M	1	Total 32	C 21	Cl 1	F 3	N 3	O 3	S 1	0	0

- Molecule 3 is SODIUM ION (CCD ID: NA) (formula: Na).

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
3	B	1	Total Na 1 1	0	0
3	D	1	Total Na 1 1	0	0
3	E	1	Total Na 1 1	0	0
3	F	1	Total Na 1 1	0	0
3	G	1	Total Na 1 1	0	0
3	H	1	Total Na 1 1	0	0

Continued on next page...

Continued from previous page...

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	L	1	Total	Na	0	0
			1	1		
3	N	1	Total	Na	0	0
			1	1		

- Molecule 4 is GLYCEROL (CCD ID: GOL) (formula: $C_3H_8O_3$).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
4	B	1	Total	C	O	0	0
			6	3	3		
4	C	1	Total	C	O	0	0
			6	3	3		
4	G	1	Total	C	O	0	0
			6	3	3		
4	N	1	Total	C	O	0	0
			6	3	3		

- Molecule 5 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
5	A	31	Total	O	0	0
			31	31		
5	B	40	Total	O	0	0
			40	40		
5	C	22	Total	O	0	0
			22	22		

Continued on next page...


Continued from previous page...

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
5	D	34	Total 34	O 34	0	0
5	E	37	Total 37	O 37	0	0
5	F	49	Total 49	O 49	0	0
5	G	31	Total 31	O 31	0	0
5	H	30	Total 30	O 30	0	0
5	I	31	Total 31	O 31	0	0
5	J	30	Total 30	O 30	0	0
5	K	43	Total 43	O 43	0	0
5	L	29	Total 29	O 29	0	0
5	M	36	Total 36	O 36	0	0
5	N	27	Total 27	O 27	0	0


3 Residue-property plots

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: ATP-dependent Clp protease proteolytic subunit 2



- Molecule 1: ATP-dependent Clp protease proteolytic subunit 2




- Molecule 1: ATP-dependent Clp protease proteolytic subunit 2




- Molecule 1: ATP-dependent Clp protease proteolytic subunit 2





- Molecule 1: ATP-dependent Clp protease proteolytic subunit 2

Chain E: 72% 25% •




- Molecule 1: ATP-dependent Clp protease proteolytic subunit 2

Chain F: 69% 28% ••




- Molecule 1: ATP-dependent Clp protease proteolytic subunit 2

Chain G: 63% 33% ••




- Molecule 1: ATP-dependent Clp protease proteolytic subunit 2


Chain H: 67% 29% ••




- Molecule 1: ATP-dependent Clp protease proteolytic subunit 2




- Molecule 1: ATP-dependent Clp protease proteolytic subunit 2




- Molecule 1: ATP-dependent Clp protease proteolytic subunit 2




- Molecule 1: ATP-dependent Clp protease proteolytic subunit 2



- Molecule 1: ATP-dependent Clp protease proteolytic subunit 2



- Chain N: 68% 28% ..



4 Data and refinement statistics

Property	Value	Source
Space group	P 1	Depositor
Cell constants a, b, c, α , β , γ	97.00Å 98.01Å 97.97Å 97.16° 114.15° 114.16°	Depositor
Resolution (Å)	33.09 – 2.66 33.09 – 2.66	Depositor EDS
% Data completeness (in resolution range)	78.2 (33.09-2.66) 78.4 (33.09-2.66)	Depositor EDS
R_{merge}	0.08	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	2.22 (at 2.65Å)	Xtriage
Refinement program	PHENIX 1.20.1_4487	Depositor
R, R_{free}	0.173 , 0.210 (Not available) , 0.216	Depositor DCC
R_{free} test set	2033 reflections (2.97%)	wwPDB-VP
Wilson B-factor (Å ²)	36.4	Xtriage
Anisotropy	0.018	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.36 , 21.3	EDS
L-test for twinning ²	$\langle L \rangle = 0.49$, $\langle L^2 \rangle = 0.32$	Xtriage
Estimated twinning fraction	0.449 for -h,-l,-k	Xtriage
F_o, F_c correlation	0.97	EDS
Total number of atoms	22282	wwPDB-VP
Average B, all atoms (Å ²)	44.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 5.36% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality ⓘ

5.1 Standard geometry ⓘ

Bond lengths and bond angles in the following residue types are not validated in this section: NA, GOL, UER

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.14	0/1562	0.45	0/2106
1	B	0.20	0/1562	0.47	1/2106 (0.0%)
1	C	0.37	3/1562 (0.2%)	0.84	7/2106 (0.3%)
1	D	0.18	0/1562	0.45	0/2106
1	E	0.23	0/1562	0.56	2/2106 (0.1%)
1	F	0.18	0/1562	0.49	3/2106 (0.1%)
1	G	0.38	1/1562 (0.1%)	0.64	7/2106 (0.3%)
1	H	0.47	1/1562 (0.1%)	0.60	5/2106 (0.2%)
1	I	0.27	0/1562	1.00	13/2106 (0.6%)
1	J	0.31	0/1562	0.60	3/2106 (0.1%)
1	K	0.22	0/1562	0.47	0/2106
1	L	0.22	0/1562	0.46	0/2106
1	M	0.25	0/1562	0.52	4/2106 (0.2%)
1	N	0.29	0/1562	0.65	7/2106 (0.3%)
All	All	0.28	5/21868 (0.0%)	0.61	52/29484 (0.2%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	D	0	2
1	E	0	1
1	F	0	3
1	G	0	1
1	H	0	4
1	I	0	1
1	J	0	1
1	M	0	1
1	N	0	4

Continued on next page...

Continued from previous page...

Mol	Chain	#Chirality outliers	#Planarity outliers
All	All	0	18

All (5) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	H	13	ARG	C-N	13.76	1.42	1.33
1	G	9	GLU	CD-OE1	-6.59	1.12	1.25
1	C	16	ARG	CB-CG	-5.90	1.34	1.52
1	C	16	ARG	CG-CD	-5.61	1.35	1.52
1	C	54	GLU	C-N	-5.60	1.29	1.33

The worst 5 of 52 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	C	16	ARG	CG-CD-NE	-24.58	57.92	112.00
1	I	9	GLU	OE1-CD-OE2	-18.95	77.42	122.90
1	I	9	GLU	CG-CD-OE1	18.82	161.69	118.40
1	C	16	ARG	CB-CG-CD	14.59	144.85	111.30
1	I	9	GLU	CA-CB-CG	12.01	138.11	114.10

There are no chirality outliers.

5 of 18 planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	D	125	PRO	Peptide
1	D	127	GLY	Peptide
1	E	16	ARG	Sidechain
1	F	8	VAL	Peptide
1	F	9	GLU	Sidechain

5.2 Too-close contacts ⓘ

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1542	0	1578	82	0
1	B	1542	0	1578	51	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	C	1542	0	1577	68	0
1	D	1542	0	1578	65	0
1	E	1542	0	1578	58	0
1	F	1542	0	1577	73	0
1	G	1542	0	1577	78	0
1	H	1542	0	1578	70	0
1	I	1542	0	1578	65	0
1	J	1542	0	1578	70	0
1	K	1542	0	1578	81	0
1	L	1542	0	1578	53	0
1	M	1542	0	1578	71	0
1	N	1542	0	1578	79	0
2	A	64	0	0	3	0
2	B	32	0	0	0	0
2	D	32	0	0	1	0
2	G	32	0	0	1	0
2	M	32	0	0	1	0
3	B	1	0	0	0	0
3	D	1	0	0	0	0
3	E	1	0	0	0	0
3	F	1	0	0	0	0
3	G	1	0	0	0	0
3	H	1	0	0	0	0
3	L	1	0	0	0	0
3	N	1	0	0	0	0
4	B	6	0	8	0	0
4	C	6	0	8	0	0
4	G	6	0	8	3	0
4	N	6	0	8	2	0
5	A	31	0	0	4	0
5	B	40	0	0	1	0
5	C	22	0	0	1	0
5	D	34	0	0	3	0
5	E	37	0	0	2	0
5	F	49	0	0	1	0
5	G	31	0	0	1	0
5	H	30	0	0	8	0
5	I	31	0	0	2	0
5	J	30	0	0	0	0
5	K	43	0	0	5	0
5	L	29	0	0	1	0
5	M	36	0	0	2	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
5	N	27	0	0	2	0
All	All	22282	0	22121	802	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 18.

The worst 5 of 802 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:C:9:GLU:OE2	1:C:23:ARG:NE	1.83	1.12
1:J:34:GLN:HG3	1:J:40:LEU:CD1	1.84	1.07
1:C:129:ILE:HD12	1:H:138:LEU:HD21	1.40	1.01
1:J:34:GLN:HG3	1:J:40:LEU:HD12	1.46	0.96
1:L:124:GLN:HB3	1:M:139:GLN:NE2	1.81	0.94

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	201/203 (99%)	185 (92%)	16 (8%)	0	100	100
1	B	201/203 (99%)	182 (90%)	19 (10%)	0	100	100
1	C	201/203 (99%)	180 (90%)	20 (10%)	1 (0%)	24	39
1	D	201/203 (99%)	184 (92%)	17 (8%)	0	100	100
1	E	201/203 (99%)	180 (90%)	18 (9%)	3 (2%)	8	13
1	F	201/203 (99%)	184 (92%)	14 (7%)	3 (2%)	8	13
1	G	201/203 (99%)	184 (92%)	15 (8%)	2 (1%)	12	20
1	H	201/203 (99%)	181 (90%)	19 (10%)	1 (0%)	24	39

Continued on next page...

Continued from previous page...

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	I	201/203 (99%)	182 (90%)	17 (8%)	2 (1%)	12	20
1	J	201/203 (99%)	179 (89%)	20 (10%)	2 (1%)	12	20
1	K	201/203 (99%)	176 (88%)	24 (12%)	1 (0%)	24	39
1	L	201/203 (99%)	180 (90%)	17 (8%)	4 (2%)	6	10
1	M	201/203 (99%)	182 (90%)	17 (8%)	2 (1%)	12	20
1	N	201/203 (99%)	185 (92%)	15 (8%)	1 (0%)	24	39
All	All	2814/2842 (99%)	2544 (90%)	248 (9%)	22 (1%)	16	27

5 of 22 Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	J	129	ILE
1	M	130	ILE
1	C	130	ILE
1	F	136	ILE
1	G	130	ILE

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	167/167 (100%)	160 (96%)	7 (4%)	26	45
1	B	167/167 (100%)	161 (96%)	6 (4%)	31	51
1	C	167/167 (100%)	164 (98%)	3 (2%)	51	72
1	D	167/167 (100%)	161 (96%)	6 (4%)	31	51
1	E	167/167 (100%)	159 (95%)	8 (5%)	23	39
1	F	167/167 (100%)	161 (96%)	6 (4%)	31	51
1	G	167/167 (100%)	161 (96%)	6 (4%)	31	51
1	H	167/167 (100%)	160 (96%)	7 (4%)	26	45
1	I	167/167 (100%)	161 (96%)	6 (4%)	31	51
1	J	167/167 (100%)	159 (95%)	8 (5%)	23	39

Continued on next page...

Continued from previous page...

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	K	167/167 (100%)	163 (98%)	4 (2%)	43	65
1	L	167/167 (100%)	156 (93%)	11 (7%)	15	26
1	M	167/167 (100%)	157 (94%)	10 (6%)	17	30
1	N	167/167 (100%)	161 (96%)	6 (4%)	31	51
All	All	2338/2338 (100%)	2244 (96%)	94 (4%)	28	47

5 of 94 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	J	102	LEU
1	L	72	THR
1	J	126	SER
1	K	16	ARG
1	L	144	LEU

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 39 such sidechains are listed below:

Mol	Chain	Res	Type
1	J	42	ASN
1	M	47	GLN
1	J	123	HIS
1	L	47	GLN
1	N	34	GLN

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates ⓘ

There are no oligosaccharides in this entry.

5.6 Ligand geometry

Of 18 ligands modelled in this entry, 8 are monoatomic - leaving 10 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
4	GOL	N	301	-	5,5,5	0.97	0	5,5,5	1.05	0
2	UER	M	301	-	33,34,34	6.05	11 (33%)	41,52,52	3.32	16 (39%)
4	GOL	G	301	-	5,5,5	0.93	0	5,5,5	1.03	0
2	UER	G	302	-	33,34,34	6.05	11 (33%)	41,52,52	3.04	16 (39%)
2	UER	D	301	-	33,34,34	6.05	12 (36%)	41,52,52	2.87	15 (36%)
4	GOL	B	303	-	5,5,5	0.94	0	5,5,5	1.04	0
2	UER	A	301	-	33,34,34	6.02	11 (33%)	41,52,52	2.94	15 (36%)
4	GOL	C	301	-	5,5,5	0.92	0	5,5,5	1.04	0
2	UER	B	301	-	33,34,34	6.05	11 (33%)	41,52,52	2.72	13 (31%)
2	UER	A	302	-	33,34,34	6.04	12 (36%)	41,52,52	2.95	16 (39%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
4	GOL	N	301	-	-	2/4/4/4	-
2	UER	M	301	-	-	20/32/45/45	0/3/3/3
4	GOL	G	301	-	-	0/4/4/4	-
2	UER	G	302	-	-	18/32/45/45	0/3/3/3
2	UER	D	301	-	-	3/32/45/45	0/3/3/3
4	GOL	B	303	-	-	0/4/4/4	-
2	UER	A	301	-	-	15/32/45/45	0/3/3/3
4	GOL	C	301	-	-	0/4/4/4	-
2	UER	B	301	-	-	9/32/45/45	0/3/3/3
2	UER	A	302	-	-	12/32/45/45	0/3/3/3

The worst 5 of 68 bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	G	302	UER	O02-S01	21.51	1.65	1.44
2	B	301	UER	O03-S01	21.49	1.65	1.44
2	A	301	UER	O02-S01	21.49	1.65	1.44
2	D	301	UER	O02-S01	21.47	1.65	1.44
2	B	301	UER	O02-S01	21.46	1.65	1.44

The worst 5 of 91 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	M	301	UER	O03-S01-O02	-8.83	107.98	119.06
2	M	301	UER	S01-C03-N01	8.67	122.50	114.40
2	G	302	UER	O03-S01-O02	-8.53	108.35	119.06
2	D	301	UER	O03-S01-O02	-8.42	108.49	119.06
2	A	302	UER	O03-S01-O02	-7.98	109.04	119.06

There are no chirality outliers.

5 of 79 torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
2	A	301	UER	C04-C03-S01-O02
2	A	301	UER	N01-C03-S01-O02
2	A	301	UER	C09-C06-C07-N02
2	A	301	UER	C07-C06-S01-O02
2	A	301	UER	C07-C06-S01-O03


There are no ring outliers.


7 monomers are involved in 11 short contacts:


Mol	Chain	Res	Type	Clashes	Symm-Clashes
4	N	301	GOL	2	0
2	M	301	UER	1	0
4	G	301	GOL	3	0
2	G	302	UER	1	0
2	D	301	UER	1	0
2	A	301	UER	1	0
2	A	302	UER	2	0


The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier.


Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.














5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	203/203 (100%)	-1.81	0 100 100	21, 37, 101, 132	0
1	B	203/203 (100%)	-1.83	0 100 100	17, 34, 103, 132	0
1	C	203/203 (100%)	-1.79	0 100 100	21, 38, 100, 136	0
1	D	203/203 (100%)	-1.84	0 100 100	18, 34, 93, 120	0
1	E	203/203 (100%)	-1.83	0 100 100	17, 32, 95, 131	0
1	F	203/203 (100%)	-1.81	0 100 100	22, 33, 103, 132	0
1	G	203/203 (100%)	-1.82	0 100 100	16, 33, 93, 129	0
1	H	203/203 (100%)	-1.81	0 100 100	19, 36, 109, 131	0
1	I	203/203 (100%)	-1.80	0 100 100	20, 34, 99, 135	0
1	J	203/203 (100%)	-1.81	0 100 100	21, 37, 99, 114	0
1	K	203/203 (100%)	-1.81	0 100 100	20, 34, 97, 137	0
1	L	203/203 (100%)	-1.80	0 100 100	21, 32, 94, 134	0
1	M	203/203 (100%)	-1.83	0 100 100	14, 30, 96, 138	0
1	N	203/203 (100%)	-1.79	0 100 100	20, 35, 108, 133	0
All	All	2842/2842 (100%)	-1.81	0 100 100	14, 35, 103, 138	0

There are no RSRZ outliers to report.

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.4 Ligands ⓘ




In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
2	UER	A	301	32/32	0.98	0.04	75,122,133,135	0
2	UER	G	302	32/32	0.98	0.05	74,96,104,106	0
3	NA	D	302	1/1	0.98	0.11	64,64,64,64	0
3	NA	F	301	1/1	0.98	0.03	19,19,19,19	0
2	UER	A	302	32/32	0.99	0.05	65,80,93,94	0
2	UER	M	301	32/32	0.99	0.04	51,89,108,113	0
2	UER	B	301	32/32	0.99	0.03	43,97,121,130	0
3	NA	E	301	1/1	0.99	0.02	20,20,20,20	0
2	UER	D	301	32/32	0.99	0.04	59,74,88,91	0
3	NA	H	301	1/1	0.99	0.04	21,21,21,21	0
4	GOL	B	303	6/6	0.99	0.02	38,48,52,60	0
4	GOL	C	301	6/6	0.99	0.02	37,39,42,46	0
4	GOL	G	301	6/6	0.99	0.05	27,29,35,46	0
3	NA	N	302	1/1	1.00	0.02	15,15,15,15	0
3	NA	G	303	1/1	1.00	0.04	32,32,32,32	0
3	NA	B	302	1/1	1.00	0.01	41,41,41,41	0
3	NA	L	301	1/1	1.00	0.02	38,38,38,38	0
4	GOL	N	301	6/6	1.00	0.03	24,28,30,35	0




The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.

Electron density around UER A 301:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)




**Electron density around UER G 302:**

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)






Electron density around UER A 302:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)




**Electron density around UER M 301:**

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)






Electron density around UER B 301:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)

**Electron density around UER D 301:**

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)



6.5 Other polymers [i](#)

There are no such residues in this entry.