



wwPDB X-ray Structure Validation Summary Report ⓘ

Mar 10, 2026 – 07:15 AM UTC

PDB ID : 7DOC / pdb_00007doc
Title : Crystal structure of Zika NS2B-NS3 protease with compound 5
Authors : Quek, J.P.
Deposited on : 2020-12-14
Resolution : 1.90 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4-5-2 with Phenix2.0
Mogul : 2022.3.0, CSD as543be (2022)
Xtriage (Phenix) : 2.0
EDS : 3.0
Percentile statistics : 20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4 : 9.0.010 (Gargrove)
Density-Fitness : 1.0.12
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 1.90 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	7789 (1.90-1.90)
Clashscore	190562	8410 (1.90-1.90)
Ramachandran outliers	187476	8333 (1.90-1.90)
Sidechain outliers	187428	8333 (1.90-1.90)
RSRZ outliers	180081	7790 (1.90-1.90)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	39	<div> <div></div> <div>74%</div> <div>26%</div> </div>
1	G	39	<div> <div>3%</div> <div></div> <div>79%</div> <div>21%</div> </div>
2	B	153	<div> <div></div> <div>76%</div> <div>24%</div> </div>
3	C	38	<div> <div>3%</div> <div></div> <div>76%</div> <div>24%</div> </div>
4	D	152	<div> <div>%</div> <div></div> <div>70%</div> <div>30%</div> </div>

Continued on next page...

Continued from previous page...

Mol	Chain	Length	Quality of chain
5	E	39	<div><div></div><div>74%</div><div>26%</div></div>
6	F	158	<div><div></div><div>61%</div><div>35%</div><div></div><div>.</div></div>
7	H	153	<div><div>%</div><div></div><div>75%</div><div>25%</div></div>
8	I	5	<div><div></div><div>60%</div><div>40%</div></div>

2 Entry composition [i](#)

There are 9 unique types of molecules in this entry. The entry contains 5976 atoms, of which 48 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Core protein.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	39	Total	C	N	O	S	0	0	0
			289	179	47	62	1			
1	G	39	Total	C	N	O	S	0	0	0
			287	178	44	64	1			

- Molecule 2 is a protein called Core protein.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
2	B	153	Total	C	N	O	S	0	0	0
			1083	689	179	210	5			

- Molecule 3 is a protein called Genome polypeptide.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
3	C	38	Total	C	N	O	S	0	0	0
			274	171	43	59	1			

- Molecule 4 is a protein called Core protein.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
4	D	152	Total	C	N	O	S	0	0	0
			1066	678	175	208	5			

- Molecule 5 is a protein called Serine protease subunit NS2B.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
5	E	39	Total	C	N	O	S	0	0	0
			288	179	44	64	1			

- Molecule 6 is a protein called Core protein.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
6	F	152	Total	C	N	O	S	0	0	0
			1104	700	187	212	5			

- Molecule 7 is a protein called Genome polyprotein.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
7	H	153	Total	C	N	O	S	0	0	0
			1110	702	191	212	5			

- Molecule 8 is a protein called 3-PYRIDIN-4-YL-2,4-DIHYDRO-INDENO[1,2-C.]PYRAZO
LE.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
8	I	5	Total	C	H	N	O	S	0	0
			104	33	48	14	8	1		

- Molecule 9 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
9	A	12	Total	O	0	0
			12	12		
9	B	50	Total	O	0	0
			50	50		
9	C	15	Total	O	0	0
			15	15		
9	D	60	Total	O	0	0
			60	60		
9	E	25	Total	O	0	0
			25	25		
9	F	89	Total	O	0	0
			89	89		
9	G	32	Total	O	0	0
			32	32		
9	H	88	Total	O	0	0
			88	88		

3 Residue-property plots


These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: Core protein

Chain A: 




- Molecule 1: Core protein

Chain G: 




- Molecule 2: Core protein

Chain B: 



- Molecule 3: Genome polyprotein

Chain C: 



- Molecule 4: Core protein

Chain D: 






- Molecule 5: Serine protease subunit NS2B

Chain E: 74% 26%




- Molecule 6: Core protein

Chain F: 61% 35% .



- Molecule 7: Genome polyprotein

Chain H: 75% 25%



- Molecule 8: 3-PYRIDIN-4-YL-2,4-DIHYDRO-INDENO[1,2-C]PYRAZOLE

Chain I: 60% 40%



4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, α , β , γ	59.15Å 59.61Å 215.07Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	45.63 – 1.90 45.63 – 1.90	Depositor EDS
% Data completeness (in resolution range)	99.4 (45.63-1.90) 97.2 (45.63-1.90)	Depositor EDS
R_{merge}	0.13	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	34.93 (at 1.91Å)	Xtriage
Refinement program	PHENIX 1.16_3549	Depositor
R, R_{free}	0.209 , 0.229 0.210 , 0.228	Depositor DCC
R_{free} test set	2982 reflections (4.92%)	wwPDB-VP
Wilson B-factor (Å ²)	35.4	Xtriage
Anisotropy	0.545	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.35 , 57.2	EDS
L-test for twinning ²	$\langle L \rangle = 0.43$, $\langle L^2 \rangle = 0.25$	Xtriage
Estimated twinning fraction	0.084 for k,h,-l	Xtriage
Reported twinning fraction	0.150 for k,h,-l	Depositor
Outliers	1 of 60244 reflections (0.002%)	Xtriage
F_o, F_c correlation	0.96	EDS
Total number of atoms	5976	wwPDB-VP
Average B, all atoms (Å ²)	45.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The analyses of the Patterson function reveals a significant off-origin peak that is 21.97 % of the origin peak, indicating pseudo-translational symmetry. The chance of finding a peak of this or larger height randomly in a structure without pseudo-translational symmetry is equal to 6.2714e-03.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: HHC

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.24	0/293	0.60	0/399
1	G	0.07	0/291	0.24	0/397
2	B	0.15	0/1107	0.33	0/1517
3	C	0.13	0/278	0.26	0/380
4	D	0.09	0/1090	0.26	0/1494
5	E	0.13	0/292	0.31	0/398
6	F	0.19	0/1126	0.32	0/1534
7	H	0.14	0/1134	0.38	0/1546
8	I	3.90	6/33 (18.2%)	2.22	3/38 (7.9%)
All	All	0.33	6/5644 (0.1%)	0.37	3/7703 (0.0%)

The worst 5 of 6 bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
8	I	4	ARG	NE-CZ	11.28	1.45	1.33
8	I	4	ARG	CZ-NH2	9.28	1.45	1.33
8	I	3	LYS	C-N	8.51	1.45	1.33
8	I	4	ARG	C-N	8.50	1.45	1.33
8	I	2	GLY	C-N	8.33	1.45	1.33

All (3) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
8	I	3	LYS	N-CA-CB	-5.44	102.79	111.43
8	I	2	GLY	CA-C-N	-5.24	114.01	122.82
8	I	2	GLY	C-N-CA	-5.24	114.01	122.82

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts ⓘ

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	289	0	256	8	2
1	G	287	0	249	8	1
2	B	1083	0	1017	33	0
3	C	274	0	239	13	0
4	D	1066	0	991	41	1
5	E	288	0	251	15	0
6	F	1104	0	1051	55	0
7	H	1110	0	1067	40	0
8	I	56	48	42	7	0
9	A	12	0	0	0	1
9	B	50	0	0	6	1
9	C	15	0	0	1	0
9	D	60	0	0	4	0
9	E	25	0	0	4	0
9	F	89	0	0	10	1
9	G	32	0	0	1	0
9	H	88	0	0	5	1
All	All	5928	48	5163	186	5

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 17.

The worst 5 of 186 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
6:F:86:ASP:OD2	9:F:201:HOH:O	1.93	0.86
7:H:94:GLU:OE1	7:H:110:GLN:NE2	2.15	0.79
7:H:27:THR:CG2	7:H:34:THR:HB	2.13	0.78
4:D:116:PHE:HB2	4:D:123:ILE:HG23	1.66	0.77
7:H:27:THR:HG22	7:H:34:THR:HB	1.67	0.77

All (5) symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:58:ASP:OD1	1:A:70:ASN:OD1[3_555]	1.41	0.79
9:F:281:HOH:O	9:H:274:HOH:O[4_545]	1.97	0.23
9:A:109:HOH:O	9:B:243:HOH:O[3_555]	2.09	0.11
1:A:58:ASP:CG	1:A:70:ASN:OD1[3_555]	2.14	0.06
4:D:68:TYR:OH	1:G:80:GLU:O[3_455]	2.17	0.03

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	37/39 (95%)	35 (95%)	2 (5%)	0	100	100
1	G	37/39 (95%)	36 (97%)	1 (3%)	0	100	100
2	B	151/153 (99%)	145 (96%)	5 (3%)	1 (1%)	18	10
3	C	36/38 (95%)	35 (97%)	1 (3%)	0	100	100
4	D	150/152 (99%)	145 (97%)	5 (3%)	0	100	100
5	E	37/39 (95%)	36 (97%)	1 (3%)	0	100	100
6	F	146/158 (92%)	141 (97%)	5 (3%)	0	100	100
7	H	151/153 (99%)	143 (95%)	8 (5%)	0	100	100
8	I	3/5 (60%)	3 (100%)	0	0	100	100
All	All	748/776 (96%)	719 (96%)	28 (4%)	1 (0%)	48	40

All (1) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
2	B	30	LEU

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar

resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	29/33 (88%)	29 (100%)	0	100	100
1	G	29/33 (88%)	29 (100%)	0	100	100
2	B	106/120 (88%)	106 (100%)	0	100	100
3	C	27/32 (84%)	27 (100%)	0	100	100
4	D	103/119 (87%)	103 (100%)	0	100	100
5	E	29/33 (88%)	29 (100%)	0	100	100
6	F	110/124 (89%)	110 (100%)	0	100	100
7	H	111/120 (92%)	111 (100%)	0	100	100
8	I	3/3 (100%)	1 (33%)	2 (67%)	0	0
All	All	547/617 (89%)	545 (100%)	2 (0%)	84	87

All (2) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
8	I	3	LYS
8	I	4	ARG

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (1) such sidechains are listed below:

Mol	Chain	Res	Type
4	D	74	GLN

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates ⓘ

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

There are no ligands in this entry.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	39/39 (100%)	0.08	0 100 100	43, 55, 73, 77	0
1	G	39/39 (100%)	-0.12	1 (2%) 57 61	32, 40, 62, 71	0
2	B	153/153 (100%)	-0.10	0 100 100	33, 47, 63, 69	0
3	C	38/38 (100%)	0.16	1 (2%) 57 61	46, 55, 71, 91	0
4	D	152/152 (100%)	0.00	1 (0%) 84 86	32, 48, 72, 91	0
5	E	39/39 (100%)	-0.20	0 100 100	36, 47, 61, 74	0
6	F	152/158 (96%)	-0.47	0 100 100	28, 36, 54, 76	0
7	H	153/153 (100%)	-0.45	1 (0%) 84 86	25, 36, 62, 78	0
8	I	4/5 (80%)	-0.10	0 100 100	50, 55, 71, 83	0
All	All	769/776 (99%)	-0.21	4 (0%) 87 89	25, 44, 67, 91	0

All (4) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
3	C	78	LEU	3.0
1	G	64	ASP	2.5
4	D	164	ALA	2.5
7	H	18	THR	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.4 Ligands [i](#)

There are no ligands in this entry.

6.5 Other polymers [i](#)

There are no such residues in this entry.