



wwPDB X-ray Structure Validation Summary Report ⓘ

Mar 9, 2026 – 06:41 AM UTC

PDB ID : 2E2M / pdb_00002e2m
Title : Crystal structure of archaeal peroxiredoxin, thioredoxin peroxidase from *Aeropyrum pernix* K1 (sulfinic acid form)
Authors : Nakamura, T.; Yamamoto, T.; Abe, M.; Matsumura, H.; Hagihara, Y.; Goto, T.; Yamaguchi, T.; Inoue, T.
Deposited on : 2006-11-14
Resolution : 2.60 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4-5-2 with Phenix2.0
Mogul : 2022.3.0, CSD as543be (2022)
Xtriage (Phenix) : 2.0
EDS : 3.0
Percentile statistics : 20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4 : 9.0.010 (Gargrove)
Density-Fitness : 1.0.12
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.49


1 Overall quality at a glance i

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.60 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	4008 (2.60-2.60)
Clashscore	190562	4347 (2.60-2.60)
Ramachandran outliers	187476	4277 (2.60-2.60)
Sidechain outliers	187428	4277 (2.60-2.60)
RSRZ outliers	180081	4008 (2.60-2.60)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	250	<div style="display: flex; align-items: center;"> <div style="width: 2%; height: 10px; background-color: red; margin-right: 5px;"></div> <div style="width: 79%; height: 10px; background-color: green; margin-right: 5px;"></div> <div style="width: 15%; height: 10px; background-color: yellow; margin-right: 5px;"></div> <div style="width: 6%; height: 10px; background-color: grey;"></div> </div>
1	B	250	<div style="display: flex; align-items: center;"> <div style="width: 3%; height: 10px; background-color: red; margin-right: 5px;"></div> <div style="width: 83%; height: 10px; background-color: green; margin-right: 5px;"></div> <div style="width: 11%; height: 10px; background-color: yellow; margin-right: 5px;"></div> <div style="width: 5%; height: 10px; background-color: grey;"></div> </div>
1	C	250	<div style="display: flex; align-items: center;"> <div style="width: 4%; height: 10px; background-color: red; margin-right: 5px;"></div> <div style="width: 80%; height: 10px; background-color: green; margin-right: 5px;"></div> <div style="width: 16%; height: 10px; background-color: yellow; margin-right: 5px;"></div> <div style="width: 0%; height: 10px; background-color: grey;"></div> </div>
1	D	250	<div style="display: flex; align-items: center;"> <div style="width: 2%; height: 10px; background-color: red; margin-right: 5px;"></div> <div style="width: 79%; height: 10px; background-color: green; margin-right: 5px;"></div> <div style="width: 14%; height: 10px; background-color: yellow; margin-right: 5px;"></div> <div style="width: 7%; height: 10px; background-color: grey;"></div> </div>
1	E	250	<div style="display: flex; align-items: center;"> <div style="width: 3%; height: 10px; background-color: red; margin-right: 5px;"></div> <div style="width: 77%; height: 10px; background-color: green; margin-right: 5px;"></div> <div style="width: 17%; height: 10px; background-color: yellow; margin-right: 5px;"></div> <div style="width: 5%; height: 10px; background-color: grey;"></div> </div>

Continued on next page...

Continued from previous page...

Mol	Chain	Length	Quality of chain
1	F	250	2% 80% 14% • 5%
1	G	250	3% 78% 17% • 5%
1	H	250	2% 77% 18% • •
1	I	250	4% 81% 13% 5%
1	J	250	2% 77% 17% • 5%

2 Entry composition

There are 2 unique types of molecules in this entry. The entry contains 19750 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Probable peroxiredoxin.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	236	Total 1920	C 1236	N 337	O 341	S 6	0	0	0
1	B	237	Total 1928	C 1241	N 338	O 343	S 6	0	0	0
1	C	242	Total 1964	C 1261	N 345	O 352	S 6	0	0	0
1	D	233	Total 1897	C 1222	N 334	O 335	S 6	0	0	0
1	E	238	Total 1934	C 1244	N 339	O 345	S 6	0	0	0
1	F	238	Total 1934	C 1244	N 339	O 345	S 6	0	0	0
1	G	238	Total 1934	C 1244	N 339	O 345	S 6	0	0	0
1	H	241	Total 1954	C 1255	N 342	O 351	S 6	0	0	0
1	I	237	Total 1929	C 1241	N 338	O 344	S 6	0	0	0
1	J	238	Total 1934	C 1244	N 339	O 345	S 6	0	0	0

There are 10 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	207	SER	CYS	engineered mutation	UNP Q9Y9L0
B	207	SER	CYS	engineered mutation	UNP Q9Y9L0
C	207	SER	CYS	engineered mutation	UNP Q9Y9L0
D	207	SER	CYS	engineered mutation	UNP Q9Y9L0
E	207	SER	CYS	engineered mutation	UNP Q9Y9L0
F	207	SER	CYS	engineered mutation	UNP Q9Y9L0
G	207	SER	CYS	engineered mutation	UNP Q9Y9L0
H	207	SER	CYS	engineered mutation	UNP Q9Y9L0
I	207	SER	CYS	engineered mutation	UNP Q9Y9L0

Continued on next page...

Continued from previous page...

Chain	Residue	Modelled	Actual	Comment	Reference
J	207	SER	CYS	engineered mutation	UNP Q9Y9L0


- Molecule 2 is water.

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
2	A	43	Total O 43 43	0	0
2	B	62	Total O 62 62	0	0
2	C	47	Total O 47 47	0	0
2	D	57	Total O 57 57	0	0
2	E	37	Total O 37 37	0	0
2	F	40	Total O 40 40	0	0
2	G	47	Total O 47 47	0	0
2	H	32	Total O 32 32	0	0
2	I	34	Total O 34 34	0	0
2	J	23	Total O 23 23	0	0


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: Probable peroxiredoxin




- Molecule 1: Probable peroxiredoxin




- Molecule 1: Probable peroxiredoxin





- Molecule 1: Probable peroxiredoxin









- Molecule 1: Probable peroxiredoxin





- Molecule 1: Probable peroxiredoxin




- Molecule 1: Probable peroxiredoxin




- Molecule 1: Probable peroxiredoxin





- Molecule 1: Probable peroxiredoxin

Chain I:  4% 81% 13% 5%



- Molecule 1: Probable peroxiredoxin

Chain J:  2% 77% 17% 5%



4 Data and refinement statistics

Property	Value	Source
Space group	P 1	Depositor
Cell constants a, b, c, α , β , γ	76.24Å 103.13Å 104.68Å 105.88° 105.12° 92.63°	Depositor
Resolution (Å)	39.37 – 2.60 39.37 – 2.60	Depositor EDS
% Data completeness (in resolution range)	90.5 (39.37-2.60) 90.5 (39.37-2.60)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.71 (at 2.61Å)	Xtrriage
Refinement program	REFMAC 5.2.0019	Depositor
R, R_{free}	0.231 , 0.280 0.225 , 0.271	Depositor DCC
R_{free} test set	4086 reflections (4.52%)	wwPDB-VP
Wilson B-factor (Å ²)	35.2	Xtrriage
Anisotropy	0.221	Xtrriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.35 , 38.1	EDS
L-test for twinning ²	$\langle L \rangle = 0.48$, $\langle L^2 \rangle = 0.30$	Xtrriage
Estimated twinning fraction	No twinning to report.	Xtrriage
F_o, F_c correlation	0.91	EDS
Total number of atoms	19750	wwPDB-VP
Average B, all atoms (Å ²)	33.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 12.79% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: CSD

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# $ Z > 5$	RMSZ	# $ Z > 5$
1	A	0.43	0/1964	0.80	4/2669 (0.1%)
1	B	0.42	0/1972	0.76	0/2680
1	C	0.42	0/2010	0.79	0/2733
1	D	0.43	0/1941	0.78	0/2637
1	E	0.42	0/1978	0.78	2/2688 (0.1%)
1	F	0.42	0/1978	0.78	2/2688 (0.1%)
1	G	0.43	0/1978	0.78	0/2688
1	H	0.43	0/1998	0.79	3/2715 (0.1%)
1	I	0.43	0/1973	0.80	4/2681 (0.1%)
1	J	0.43	0/1978	0.80	2/2688 (0.1%)
All	All	0.43	0/19770	0.79	17/26867 (0.1%)

There are no bond length outliers.

The worst 5 of 17 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	J	188	PRO	CA-C-N	6.99	124.76	119.66
1	J	188	PRO	C-N-CA	6.99	124.76	119.66
1	H	188	PRO	CA-C-N	6.36	124.31	119.66
1	H	188	PRO	C-N-CA	6.36	124.31	119.66
1	A	188	PRO	CA-C-N	6.18	124.17	119.66

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen

atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1920	0	1913	21	0
1	B	1928	0	1919	18	0
1	C	1964	0	1948	20	0
1	D	1897	0	1892	18	0
1	E	1934	0	1924	28	0
1	F	1934	0	1924	22	0
1	G	1934	0	1924	29	0
1	H	1954	0	1940	28	0
1	I	1929	0	1919	18	0
1	J	1934	0	1924	25	0
2	A	43	0	0	0	0
2	B	62	0	0	0	0
2	C	47	0	0	0	0
2	D	57	0	0	0	0
2	E	37	0	0	0	0
2	F	40	0	0	0	0
2	G	47	0	0	0	0
2	H	32	0	0	0	0
2	I	34	0	0	0	0
2	J	23	0	0	0	0
All	All	19750	0	19227	190	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 5.

The worst 5 of 190 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:G:142:TYR:HB3	1:H:8:ILE:HD11	1.48	0.95
1:A:5:ILE:HG12	1:B:5:ILE:HG12	1.61	0.83
1:J:140:MET:HE2	1:J:142:TYR:OH	1.81	0.81
1:E:147:LEU:HD12	1:F:161:LEU:HD13	1.72	0.72
1:E:177:ASN:HD21	1:E:227:ARG:HE	1.40	0.69

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	231/250 (92%)	223 (96%)	6 (3%)	2 (1%)	14	30
1	B	232/250 (93%)	224 (97%)	6 (3%)	2 (1%)	14	30
1	C	239/250 (96%)	227 (95%)	11 (5%)	1 (0%)	30	51
1	D	228/250 (91%)	221 (97%)	5 (2%)	2 (1%)	14	30
1	E	233/250 (93%)	227 (97%)	4 (2%)	2 (1%)	14	30
1	F	233/250 (93%)	226 (97%)	5 (2%)	2 (1%)	14	30
1	G	233/250 (93%)	223 (96%)	8 (3%)	2 (1%)	14	30
1	H	236/250 (94%)	227 (96%)	6 (2%)	3 (1%)	9	21
1	I	232/250 (93%)	223 (96%)	7 (3%)	2 (1%)	14	30
1	J	233/250 (93%)	223 (96%)	9 (4%)	1 (0%)	30	51
All	All	2330/2500 (93%)	2244 (96%)	67 (3%)	19 (1%)	16	34

5 of 19 Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	122	THR
1	G	239	LYS
1	H	122	THR
1	I	239	LYS
1	J	122	THR

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	204/215 (95%)	198 (97%)	6 (3%)	37	65
1	B	204/215 (95%)	200 (98%)	4 (2%)	48	74
1	C	208/215 (97%)	200 (96%)	8 (4%)	29	56
1	D	201/215 (94%)	198 (98%)	3 (2%)	57	80
1	E	205/215 (95%)	201 (98%)	4 (2%)	48	74
1	F	205/215 (95%)	198 (97%)	7 (3%)	32	60
1	G	205/215 (95%)	201 (98%)	4 (2%)	48	74
1	H	207/215 (96%)	203 (98%)	4 (2%)	50	75
1	I	205/215 (95%)	201 (98%)	4 (2%)	48	74
1	J	205/215 (95%)	196 (96%)	9 (4%)	25	50
All	All	2049/2150 (95%)	1996 (97%)	53 (3%)	40	68

5 of 53 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	F	101	ILE
1	G	235	GLU
1	J	167	LEU
1	F	147	LEU
1	G	116	LEU

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 14 such sidechains are listed below:

Mol	Chain	Res	Type
1	C	65	GLN
1	C	195	GLN
1	J	29	HIS
1	E	195	GLN
1	G	29	HIS

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

10 non-standard protein/DNA/RNA residues are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
1	CSD	G	50	1	4,7,8	18.15	1 (25%)	1,8,10	3.66	1 (100%)
1	CSD	E	50	1	4,7,8	18.16	1 (25%)	1,8,10	2.15	1 (100%)
1	CSD	D	50	1	4,7,8	18.18	1 (25%)	1,8,10	3.21	1 (100%)
1	CSD	B	50	1	4,7,8	17.95	1 (25%)	1,8,10	2.17	1 (100%)
1	CSD	I	50	1	4,7,8	18.00	1 (25%)	1,8,10	0.41	0
1	CSD	J	50	1	4,7,8	18.01	1 (25%)	1,8,10	2.69	1 (100%)
1	CSD	A	50	1	4,7,8	18.03	1 (25%)	1,8,10	2.08	1 (100%)
1	CSD	H	50	1	4,7,8	17.83	1 (25%)	1,8,10	1.52	0
1	CSD	F	50	1	4,7,8	18.28	1 (25%)	1,8,10	1.21	0
1	CSD	C	50	1	4,7,8	18.04	1 (25%)	1,8,10	1.53	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
1	CSD	G	50	1	-	1/2/6/8	-
1	CSD	E	50	1	-	0/2/6/8	-
1	CSD	D	50	1	-	1/2/6/8	-
1	CSD	B	50	1	-	0/2/6/8	-
1	CSD	I	50	1	-	1/2/6/8	-
1	CSD	J	50	1	-	0/2/6/8	-
1	CSD	A	50	1	-	0/2/6/8	-
1	CSD	H	50	1	-	1/2/6/8	-
1	CSD	F	50	1	-	0/2/6/8	-
1	CSD	C	50	1	-	1/2/6/8	-

The worst 5 of 10 bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	F	50	CSD	OD1-SG	36.55	1.80	1.47

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	D	50	CSD	OD1-SG	36.34	1.80	1.47
1	E	50	CSD	OD1-SG	36.31	1.80	1.47
1	G	50	CSD	OD1-SG	36.28	1.80	1.47
1	C	50	CSD	OD1-SG	36.06	1.80	1.47

The worst 5 of 6 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	G	50	CSD	OD1-SG-CB	3.66	112.34	105.60
1	D	50	CSD	OD1-SG-CB	3.21	111.51	105.60
1	J	50	CSD	OD1-SG-CB	2.69	110.57	105.60
1	B	50	CSD	OD1-SG-CB	2.17	109.61	105.60
1	E	50	CSD	OD1-SG-CB	2.15	109.56	105.60

There are no chirality outliers.

All (5) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
1	C	50	CSD	CA-CB-SG-OD1
1	D	50	CSD	CA-CB-SG-OD1
1	G	50	CSD	CA-CB-SG-OD1
1	H	50	CSD	CA-CB-SG-OD1
1	I	50	CSD	CA-CB-SG-OD1

There are no ring outliers.

1 monomer is involved in 1 short contact:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
1	F	50	CSD	1	0

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

There are no ligands in this entry.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	235/250 (94%)	-0.02	4 (1%) 69 64	22, 29, 43, 48	0
1	B	236/250 (94%)	0.18	7 (2%) 52 47	22, 30, 44, 53	0
1	C	241/250 (96%)	0.21	10 (4%) 41 36	23, 32, 49, 53	0
1	D	232/250 (92%)	0.21	5 (2%) 62 57	23, 32, 44, 46	0
1	E	237/250 (94%)	0.36	8 (3%) 48 42	29, 37, 54, 58	0
1	F	237/250 (94%)	0.17	4 (1%) 69 64	26, 34, 52, 60	0
1	G	237/250 (94%)	0.24	8 (3%) 48 42	25, 32, 46, 53	0
1	H	240/250 (96%)	0.26	6 (2%) 58 52	25, 33, 49, 59	0
1	I	236/250 (94%)	0.22	10 (4%) 40 35	23, 32, 49, 55	0
1	J	237/250 (94%)	0.17	5 (2%) 63 58	24, 33, 47, 53	0
All	All	2368/2500 (94%)	0.20	67 (2%) 55 49	22, 33, 48, 60	0

The worst 5 of 67 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	B	121	ALA	4.6
1	J	245	ALA	4.5
1	H	118	ALA	4.3
1	H	120	SER	4.1
1	C	118	ALA	3.8

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled ‘Q< 0.9’ lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(\AA^2)	Q<0.9
1	CSD	I	50	8/9	0.93	0.10	32,32,33,34	0
1	CSD	F	50	8/9	0.94	0.07	27,27,28,28	0
1	CSD	D	50	8/9	0.95	0.09	28,28,30,31	0
1	CSD	A	50	8/9	0.95	0.08	25,25,26,27	0
1	CSD	H	50	8/9	0.95	0.10	28,29,30,30	0
1	CSD	C	50	8/9	0.95	0.09	30,30,31,31	0
1	CSD	J	50	8/9	0.95	0.09	26,26,27,27	0
1	CSD	G	50	8/9	0.96	0.07	28,28,30,30	0
1	CSD	E	50	8/9	0.96	0.08	29,29,30,31	0
1	CSD	B	50	8/9	0.97	0.06	24,24,24,24	0

6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.4 Ligands [i](#)

There are no ligands in this entry.

6.5 Other polymers [i](#)

There are no such residues in this entry.