



Full wwPDB X-ray Structure Validation Report ⓘ

Mar 18, 2026 – 01:34 AM UTC

PDB ID : 3E65 / pdb_00003e65
Title : Murine INOS dimer with HEME, pterin and inhibitor AR-C120011
Authors : Rosenfeld, R.J.; Garcin, E.D.; Getzoff, E.D.
Deposited on : 2008-08-14
Resolution : 2.05 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4-5-2 with Phenix2.0
Mogul : 2022.3.0, CSD as543be (2022)
Xtriage (Phenix) : 2.0
EDS : 3.0
Buster-report : wwPDB partial adaption of 1.1.7 (2018)
Percentile statistics : 20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4 : 9.0.010 (Gargrove)
Density-Fitness : 1.0.12
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.05 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	2260 (2.04-2.04)
Clashscore	190562	2333 (2.04-2.04)
Ramachandran outliers	187476	2318 (2.04-2.04)
Sidechain outliers	187428	2318 (2.04-2.04)
RSRZ outliers	180081	2260 (2.04-2.04)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	433	
1	B	433	

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
3	H4B	A	902	X	-	-	-
3	H4B	B	902	X	-	-	-

2 Entry composition [i](#)

There are 5 unique types of molecules in this entry. The entry contains 7139 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Nitric oxide synthase, inducible.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	417	Total	C	N	O	S	0	0	0
			3395	2177	584	614	20			
1	B	421	Total	C	N	O	S	0	0	0
			3421	2193	589	618	21			

- Molecule 2 is PROTOPORPHYRIN IX CONTAINING FE (CCD ID: HEM) (formula: $C_{34}H_{32}FeN_4O_4$).




Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
2	A	1	Total	C	Fe	N	O	0	0
			43	34	1	4	4		
2	B	1	Total	C	Fe	N	O	0	0
			43	34	1	4	4		

- Molecule 3 is 5,6,7,8-TETRAHYDROBIOPTERIN (CCD ID: H4B) (formula: $C_9H_{15}N_5O_3$).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
3	A	1	Total	C	N	O	0	0
			17	9	5	3		
3	B	1	Total	C	N	O	0	0
			17	9	5	3		

- Molecule 4 is 1-[4-(AMINOMETHYL)BENZOYL]-5'-FLUORO-1'H-SPIRO[PIPERIDINE-4,2'-QUINAZOLIN]-4'-AMINE (CCD ID: XXZ) (formula: $C_{20}H_{22}FN_5O$).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
4	A	1	Total	C	F	N	O	0
			27	20	1	5	1	

Continued on next page...

Continued from previous page...

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
4	B	1	Total	C	F	N	O	0	0
			27	20	1	5	1		


- Molecule 5 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
5	A	75	Total	O	0	0
			75	75		
5	B	74	Total	O	0	0
			74	74		

3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: Nitric oxide synthase, inducible



4 Data and refinement statistics

Property	Value	Source
Space group	P 61 2 2	Depositor
Cell constants a, b, c, α , β , γ	213.94Å 213.94Å 117.03Å 90.00° 90.00° 120.00°	Depositor
Resolution (Å)	19.98 – 2.05 19.98 – 2.05	Depositor EDS
% Data completeness (in resolution range)	95.5 (19.98-2.05) 95.3 (19.98-2.05)	Depositor EDS
R_{merge}	0.06	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	4.15 (at 2.05Å)	Xtriage
Refinement program	CNS	Depositor
R, R_{free}	0.249 , 0.271 0.241 , 0.259	Depositor DCC
R_{free} test set	4735 reflections (5.03%)	wwPDB-VP
Wilson B-factor (Å ²)	36.7	Xtriage
Anisotropy	0.595	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.35 , 28.7	EDS
L-test for twinning ²	$\langle L \rangle = 0.52$, $\langle L^2 \rangle = 0.36$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.94	EDS
Total number of atoms	7139	wwPDB-VP
Average B, all atoms (Å ²)	44.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 2.58% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: XXZ, H4B, HEM

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.45	0/3494	0.94	16/4751 (0.3%)
1	B	0.43	0/3521	0.91	10/4787 (0.2%)
All	All	0.44	0/7015	0.92	26/9538 (0.3%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	A	0	1

There are no bond length outliers.

All (26) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	B	459	VAL	CA-C-N	9.39	126.51	119.66
1	B	459	VAL	C-N-CA	9.39	126.51	119.66
1	A	102	PHE	N-CA-C	7.95	127.74	110.80
1	A	108	SER	CA-C-N	7.79	132.55	121.02
1	A	108	SER	C-N-CA	7.79	132.55	121.02
1	B	368	MET	N-CA-C	-7.27	97.36	109.07
1	A	368	MET	N-CA-C	-7.10	97.33	108.90
1	B	378	CYS	N-CA-C	6.31	121.14	113.38
1	A	122	GLY	N-CA-C	6.30	119.56	112.00
1	A	488	GLU	CA-C-N	5.93	126.08	119.32
1	A	488	GLU	C-N-CA	5.93	126.08	119.32
1	B	122	GLY	N-CA-C	5.84	119.01	112.00
1	A	153	ILE	N-CA-C	5.83	116.48	110.36
1	A	234	ILE	N-CA-C	5.79	116.89	109.30

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	358	PHE	CA-C-N	5.77	125.87	119.87
1	A	358	PHE	C-N-CA	5.77	125.87	119.87
1	B	369	GLY	N-CA-C	5.60	119.66	112.83
1	B	303	LEU	N-CA-C	5.55	118.00	109.07
1	A	110	LEU	CA-C-O	5.50	127.20	120.55
1	A	122	GLY	CA-C-N	5.22	125.69	120.52
1	A	122	GLY	C-N-CA	5.22	125.69	120.52
1	A	378	CYS	N-CA-C	5.12	120.38	113.72
1	B	234	ILE	N-CA-C	5.09	115.97	109.30
1	A	197	ARG	N-CA-C	5.07	118.96	112.87
1	B	386	LEU	N-CA-C	5.05	116.48	111.07
1	B	102	PHE	N-CA-C	5.03	114.79	108.45

There are no chirality outliers.

All (1) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	A	108	SER	Mainchain

5.2 Too-close contacts ⓘ

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	3395	0	3281	37	0
1	B	3421	0	3312	32	0
2	A	43	0	30	0	0
2	B	43	0	30	0	0
3	A	17	0	14	0	0
3	B	17	0	14	0	0
4	A	27	0	22	2	0
4	B	27	0	22	1	0
5	A	75	0	0	3	0
5	B	74	0	0	0	0
All	All	7139	0	6725	71	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 5.

All (71) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:366:TRP:HE1	1:A:428:MET:HE3	1.25	1.01
1:B:215:GLN:HE21	1:B:219:GLN:HE21	1.23	0.86
1:A:81:ILE:HD11	1:A:475:LEU:HD13	1.62	0.81
1:B:81:ILE:HD11	1:B:475:LEU:HD13	1.62	0.79
1:A:366:TRP:NE1	1:A:428:MET:HE3	2.00	0.77
1:A:215:GLN:HE21	1:A:219:GLN:HE21	1.33	0.76
1:A:83:ASN:ND2	1:A:85:GLY:H	1.85	0.74
1:B:215:GLN:HE21	1:B:219:GLN:NE2	1.89	0.68
1:B:194:CYS:HB3	1:B:197:ARG:HD2	1.75	0.68
1:A:83:ASN:HD22	1:A:85:GLY:H	1.43	0.64
1:A:153:ILE:O	1:A:157:LEU:HD23	2.02	0.60
1:A:108:SER:O	1:A:110:LEU:HD12	2.01	0.60
1:A:108:SER:O	1:A:110:LEU:CD1	2.50	0.60
1:A:366:TRP:HE1	1:A:428:MET:CE	2.10	0.60
1:B:144:TYR:O	1:B:147:SER:HB3	2.02	0.59
1:A:188:TRP:CE3	1:A:200:TRP:HA	2.40	0.57
1:A:465:SER:O	1:A:471:HIS:HE1	1.89	0.56
1:B:104:CYS:SG	1:B:109:CYS:HA	2.47	0.55
1:A:442:GLN:HE21	1:A:446:ARG:HH12	1.56	0.54
1:B:188:TRP:CE3	1:B:200:TRP:HA	2.42	0.54
1:A:77:GLN:O	1:A:96:HIS:HE1	1.90	0.54
1:A:428:MET:HE1	5:A:1320:HOH:O	2.07	0.53
1:B:101:ASP:CG	1:B:102:PHE:H	2.17	0.53
1:B:81:ILE:HD11	1:B:475:LEU:CD1	2.37	0.52
1:B:266:MET:HE2	1:B:272:ARG:CZ	2.40	0.51
1:A:209:ARG:O	1:A:242:PRO:HG3	2.12	0.49
1:A:246:ASP:OD1	1:A:249:HIS:HD2	1.95	0.49
1:A:83:ASN:HD22	1:A:83:ASN:C	2.21	0.49
1:A:469:VAL:HG13	1:A:474:MET:HE1	1.95	0.48
1:B:194:CYS:O	1:B:197:ARG:HD3	2.14	0.48
1:B:134:LEU:HB3	1:B:135:PRO:HD3	1.96	0.48
1:A:134:LEU:HB3	1:A:135:PRO:HD3	1.96	0.48
1:A:460:PRO:HG3	5:A:1444:HOH:O	2.14	0.47
1:B:246:ASP:OD1	1:B:249:HIS:HD2	1.98	0.47
4:B:907:XXZ:CE1	4:B:907:XXZ:H111	2.45	0.46
1:B:215:GLN:NE2	1:B:219:GLN:HE21	2.02	0.46
1:A:251:PHE:O	1:A:360:ALA:HB2	2.16	0.46
1:A:195:ILE:HD12	5:A:1443:HOH:O	2.15	0.46
1:B:253:LEU:HD22	1:B:347:ALA:HB1	1.99	0.45
1:B:137:ALA:O	1:B:141:ILE:HG12	2.16	0.45

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:77:GLN:O	1:B:96:HIS:HE1	2.00	0.45
1:A:81:ILE:HD11	1:A:475:LEU:CD1	2.40	0.44
1:B:215:GLN:O	1:B:219:GLN:HG3	2.18	0.44
1:A:243:GLN:HB3	1:A:358:PHE:CE2	2.52	0.44
1:B:252:ARG:NH2	1:B:489:PRO:HD3	2.33	0.43
1:B:132:GLU:O	1:B:135:PRO:HD2	2.19	0.43
1:B:103:THR:HG22	1:B:116:PRO:HB2	1.99	0.43
1:B:239:THR:O	1:B:361:CYS:HA	2.19	0.43
4:A:906:XXZ:CE1	4:A:906:XXZ:H111	2.48	0.42
1:B:190:ASN:O	1:B:192:PRO:HD3	2.18	0.42
1:B:480:SER:HA	1:B:481:PRO:C	2.44	0.42
1:B:188:TRP:CZ3	1:B:200:TRP:HA	2.54	0.42
1:B:243:GLN:HB3	1:B:358:PHE:CE2	2.54	0.42
1:A:83:ASN:HD22	1:A:85:GLY:N	2.13	0.42
1:A:188:TRP:CZ3	1:A:200:TRP:HA	2.54	0.42
1:A:254:TRP:HB2	1:A:302:VAL:HB	2.01	0.42
1:B:290:LYS:HA	1:B:291:PRO:HD2	1.96	0.42
1:A:388:GLU:O	1:A:392:ARG:HG3	2.19	0.42
1:A:190:ASN:O	1:A:192:PRO:HD3	2.20	0.41
1:A:382:ARG:NH2	4:A:906:XXZ:HD2	2.35	0.41
1:A:469:VAL:HG13	1:A:474:MET:CE	2.50	0.41
1:B:361:CYS:O	1:B:361:CYS:SG	2.79	0.41
1:B:110:LEU:HD12	1:B:110:LEU:N	2.35	0.41
1:B:254:TRP:HB2	1:B:302:VAL:HB	2.02	0.41
1:A:488:GLU:HA	1:A:489:PRO:HD2	1.87	0.41
1:B:103:THR:HG23	1:B:118:SER:OG	2.21	0.41
1:B:460:PRO:HG2	1:B:467:THR:HG21	2.03	0.41
1:A:189:ARG:HD2	1:A:200:TRP:CE3	2.56	0.41
1:A:446:ARG:HG3	1:A:446:ARG:HH21	1.85	0.41
1:A:239:THR:O	1:A:361:CYS:HA	2.21	0.40
1:A:252:ARG:NH2	1:A:489:PRO:HD3	2.36	0.40

There are no symmetry-related clashes.

5.3 Torsion angles ⓘ

5.3.1 Protein backbone ⓘ

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	413/433 (95%)	396 (96%)	16 (4%)	1 (0%)	43	37
1	B	419/433 (97%)	398 (95%)	20 (5%)	1 (0%)	43	37
All	All	832/866 (96%)	794 (95%)	36 (4%)	2 (0%)	43	37

All (2) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	B	105	LYS
1	A	100	SER

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	363/381 (95%)	359 (99%)	4 (1%)	65	68
1	B	366/381 (96%)	363 (99%)	3 (1%)	73	77
All	All	729/762 (96%)	722 (99%)	7 (1%)	68	72

All (7) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	83	ASN
1	A	103	THR
1	A	109	CYS
1	A	475	LEU
1	B	265	GLN
1	B	295	ARG
1	B	475	LEU

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (18) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	83	ASN
1	A	96	HIS
1	A	115	ASN
1	A	215	GLN
1	A	249	HIS
1	A	421	GLN
1	A	442	GLN
1	A	471	HIS
1	A	486	GLN
1	B	91	HIS
1	B	96	HIS
1	B	115	ASN
1	B	143	GLN
1	B	215	GLN
1	B	249	HIS
1	B	418	HIS
1	B	421	GLN
1	B	486	GLN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

6 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
4	XXZ	B	907	-	26,30,30	2.56	5 (19%)	37,44,44	1.70	9 (24%)
3	H4B	B	902	-	17,18,18	3.34	4 (23%)	14,26,26	1.98	5 (35%)
3	H4B	A	902	-	17,18,18	3.38	4 (23%)	14,26,26	1.97	5 (35%)
2	HEM	B	901	1	50,50,50	1.29	5 (10%)	67,82,82	1.01	4 (5%)
4	XXZ	A	906	-	26,30,30	2.64	5 (19%)	37,44,44	1.71	7 (18%)
2	HEM	A	901	1	50,50,50	1.41	7 (14%)	67,82,82	0.87	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
4	XXZ	B	907	-	-	0/10/36/36	0/3/4/4
3	H4B	B	902	-	1/1/3/5	2/8/17/17	0/2/2/2
3	H4B	A	902	-	1/1/3/5	2/8/17/17	0/2/2/2
2	HEM	B	901	1	-	4/14/54/54	-
4	XXZ	A	906	-	-	0/10/36/36	0/3/4/4
2	HEM	A	901	1	-	5/14/54/54	-

All (30) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
4	A	906	XXZ	C6-N1	11.17	1.56	1.39
4	B	907	XXZ	C6-N1	10.68	1.55	1.39
3	A	902	H4B	C7-C6	-8.80	1.43	1.52
3	B	902	H4B	C7-C6	-8.53	1.44	1.52
3	A	902	H4B	C6-N5	-7.06	1.33	1.47
3	B	902	H4B	C6-N5	-7.04	1.33	1.47
3	B	902	H4B	C7-N8	-6.99	1.39	1.46
3	A	902	H4B	C7-N8	-6.87	1.39	1.46
4	A	906	XXZ	C7-CG	4.05	1.52	1.45
4	B	907	XXZ	C7-CG	3.93	1.52	1.45
2	A	901	HEM	FE-NA	3.71	2.07	1.95
2	A	901	HEM	FE-ND	3.56	2.05	1.94
2	B	901	HEM	CAB-C3B	-3.39	1.38	1.47
2	A	901	HEM	CAB-C3B	-3.36	1.38	1.47
4	B	907	XXZ	CG-N20	2.93	1.37	1.29
4	A	906	XXZ	CG-N20	2.91	1.36	1.29
3	B	902	H4B	C4-N3	-2.87	1.33	1.38

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	B	901	HEM	FE-NA	2.79	2.04	1.95
2	B	901	HEM	FE-NB	2.78	2.03	1.94
2	B	901	HEM	FE-ND	2.75	2.03	1.94
3	A	902	H4B	C4-N3	-2.67	1.33	1.38
2	A	901	HEM	CAC-C3C	-2.65	1.40	1.47
2	A	901	HEM	FE-NB	2.64	2.03	1.94
2	B	901	HEM	CAC-C3C	-2.50	1.40	1.47
4	B	907	XXZ	CE1-CZ	2.48	1.43	1.39
2	A	901	HEM	FE-NC	2.46	2.03	1.95
4	B	907	XXZ	C15-N12	2.43	1.39	1.34
4	A	906	XXZ	CE1-CZ	2.26	1.42	1.39
2	A	901	HEM	CBC-CAC	2.13	1.40	1.30
4	A	906	XXZ	C15-N12	2.05	1.39	1.34

All (30) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
4	A	906	XXZ	C10-C5-N1	-5.24	102.26	110.47
4	B	907	XXZ	C10-C5-N1	-5.13	102.42	110.47
3	B	902	H4B	C2-N1-C8A	4.44	121.19	113.36
3	A	902	H4B	C2-N1-C8A	4.31	120.96	113.36
4	B	907	XXZ	C14-C5-C10	-3.93	106.83	109.99
4	A	906	XXZ	C14-C5-C10	-3.88	106.87	109.99
3	A	902	H4B	O4-C4-C4A	-3.77	118.17	127.26
3	B	902	H4B	O4-C4-C4A	-3.65	118.47	127.26
4	B	907	XXZ	C6-C7-CG	-2.89	114.40	117.41
3	A	902	H4B	C4A-C4-N3	2.84	119.92	112.13
3	B	902	H4B	C4A-C4-N3	2.77	119.73	112.13
4	B	907	XXZ	C13-C14-C5	-2.75	108.88	112.55
4	A	906	XXZ	CE1-CZ-C18	2.66	121.96	118.57
4	A	906	XXZ	C13-C14-C5	-2.62	109.05	112.55
2	B	901	HEM	CMD-C2D-C1D	2.58	129.07	125.03
4	B	907	XXZ	CE1-CZ-C18	2.53	121.79	118.57
3	B	902	H4B	C2-N3-C4	-2.48	120.61	125.11
4	A	906	XXZ	CD2-C8-CD1	2.48	121.92	118.23
3	A	902	H4B	C2-N3-C4	-2.46	120.66	125.11
4	A	906	XXZ	C6-C7-CG	-2.45	114.86	117.41
4	B	907	XXZ	CD2-C8-CD1	2.42	121.84	118.23
4	A	906	XXZ	C14-C13-N12	-2.24	106.64	110.82
2	B	901	HEM	CAA-C2A-C1A	2.22	129.28	124.94
4	B	907	XXZ	C7-CG-N21	2.19	123.48	119.46
2	B	901	HEM	CAB-C3B-C2B	-2.14	121.49	128.43

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
4	B	907	XXZ	C14-C13-N12	-2.06	106.97	110.82
3	A	902	H4B	C4-C4A-N5	2.05	121.85	116.27
4	B	907	XXZ	CE1-CD1-C8	-2.02	118.34	121.00
3	B	902	H4B	C4-C4A-N5	2.02	121.77	116.27
2	B	901	HEM	C4A-C3A-C2A	2.00	109.11	106.82

All (2) chirality outliers are listed below:

Mol	Chain	Res	Type	Atom
3	A	902	H4B	C6
3	B	902	H4B	C6

All (13) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
3	A	902	H4B	C7-C6-C9-O9
3	A	902	H4B	C7-C6-C9-C10
3	B	902	H4B	C7-C6-C9-O9
3	B	902	H4B	C7-C6-C9-C10
2	A	901	HEM	C2B-C3B-CAB-CBB
2	A	901	HEM	C4B-C3B-CAB-CBB
2	B	901	HEM	C4B-C3B-CAB-CBB
2	B	901	HEM	C2B-C3B-CAB-CBB
2	A	901	HEM	CAA-CBA-CGA-O1A
2	B	901	HEM	CAA-CBA-CGA-O1A
2	B	901	HEM	CAA-CBA-CGA-O2A
2	A	901	HEM	CAA-CBA-CGA-O2A
2	A	901	HEM	CAD-CBD-CGD-O2D


There are no ring outliers.



2 monomers are involved in 3 short contacts:


Mol	Chain	Res	Type	Clashes	Symm-Clashes
4	B	907	XXZ	1	0
4	A	906	XXZ	2	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be

highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.







5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	417/433 (96%)	0.78	38 (9%) 15 15	29, 43, 68, 88	0
1	B	421/433 (97%)	0.77	46 (10%) 10 10	27, 41, 67, 89	0
All	All	838/866 (96%)	0.78	84 (10%) 12 12	27, 42, 68, 89	0

All (84) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	B	102	PHE	9.0
1	B	99	THR	8.4
1	B	108	SER	8.1
1	A	102	PHE	8.1
1	A	103	THR	8.0
1	B	103	THR	7.0
1	B	107	LYS	6.0
1	A	110	LEU	5.8
1	B	106	SER	5.8
1	A	77	GLN	5.7
1	A	109	CYS	5.3
1	A	108	SER	5.2
1	A	497	ASN	5.1
1	B	497	ASN	4.8
1	A	100	SER	4.8
1	A	111	GLY	4.6
1	B	101	ASP	4.5
1	B	109	CYS	4.4
1	B	105	LYS	4.3
1	B	100	SER	4.2
1	A	101	ASP	4.0
1	B	130	LEU	3.8
1	B	110	LEU	3.5
1	B	151	ALA	3.4

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	B	111	GLY	3.3
1	B	98	ALA	3.2
1	B	494	ILE	3.2
1	A	81	ILE	3.2
1	B	131	GLU	3.2
1	B	104	CYS	3.2
1	B	77	GLN	3.1
1	A	99	THR	3.1
1	B	496	GLN	3.1
1	B	112	SER	3.0
1	A	146	GLY	3.0
1	A	495	TRP	3.0
1	B	267	PRO	2.9
1	B	148	PHE	2.8
1	A	165	LYS	2.8
1	A	124	ARG	2.8
1	B	450	GLY	2.8
1	B	78	TYR	2.7
1	A	80	ARG	2.7
1	B	292	ARG	2.7
1	B	81	ILE	2.6
1	B	495	TRP	2.6
1	A	169	THR	2.5
1	B	162	ALA	2.5
1	A	98	ALA	2.5
1	A	153	ILE	2.5
1	A	494	ILE	2.5
1	A	496	GLN	2.4
1	B	150	GLU	2.4
1	A	381	GLN	2.4
1	A	442	GLN	2.4
1	B	153	ILE	2.4
1	A	152	LYS	2.4
1	B	447	ALA	2.3
1	B	391	ARG	2.3
1	B	79	VAL	2.3
1	A	127	PRO	2.3
1	A	78	TYR	2.3
1	A	139	GLU	2.3
1	A	151	ALA	2.3
1	B	158	ALA	2.3
1	A	330	TYR	2.3

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	B	330	TYR	2.3
1	A	446	ARG	2.3
1	B	171	GLY	2.3
1	B	84	TRP	2.3
1	A	334	GLN	2.2
1	B	308	GLN	2.2
1	A	308	GLN	2.2
1	B	128	THR	2.2
1	B	91	HIS	2.2
1	B	380	THR	2.2
1	A	149	LYS	2.1
1	B	314	GLU	2.1
1	A	292	ARG	2.1
1	B	247	GLY	2.1
1	A	160	LEU	2.1
1	B	147	SER	2.1
1	A	79	VAL	2.0
1	A	84	TRP	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.


Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
3	H4B	A	902	17/17	0.94	0.06	30,33,36,36	0
4	XXZ	A	906	27/27	0.94	0.09	26,29,35,38	0
4	XXZ	B	907	27/27	0.94	0.09	28,31,34,36	0
3	H4B	B	902	17/17	0.96	0.06	24,30,33,35	0
2	HEM	A	901	43/43	0.97	0.08	28,31,34,39	0

Continued on next page...

Continued from previous page...




Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors($\text{\AA}^2$)	Q<0.9
2	HEM	B	901	43/43	0.98	0.07	25,29,32,39	0

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.






Electron density around XXZ B 907:


$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)



Electron density around HEM A 901:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)





6.5 Other polymers [i](#)

There are no such residues in this entry.