



wwPDB X-ray Structure Validation Summary Report ⓘ

Mar 14, 2026 – 11:03 AM UTC

PDB ID : 6EA4 / pdb_00006ea4
Title : ERAP2 bound to Aryl Sulfonamide Uncompetitive Inhibitor
Authors : Maben, Z.; Stern, L.J.
Deposited on : 2018-08-02
Resolution : 2.45 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4-5-2 with Phenix2.0
Mogul : 2022.3.0, CSD as543be (2022)
Xtriage (Phenix) : 2.0
EDS : 3.0
Buster-report : wwPDB partial adaption of 1.1.7 (2018)
Percentile statistics : 20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4 : 9.0.010 (Gargrove)
Density-Fitness : 1.0.12
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.45 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.








Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	1190 (2.46-2.46)
Clashscore	190562	1229 (2.46-2.46)
Ramachandran outliers	187476	1218 (2.46-2.46)
Sidechain outliers	187428	1218 (2.46-2.46)
RSRZ outliers	180081	1190 (2.46-2.46)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	911	
1	B	911	
2	C	5	
2	D	5	
2	G	5	

Continued on next page...

Continued from previous page...

Mol	Chain	Length	Quality of chain
3	E	2	 100%
3	F	2	 100%
3	I	2	 100%
4	H	3	 67%  33%

2 Entry composition i

There are 13 unique types of molecules in this entry. The entry contains 15097 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.


- Molecule 1 is a protein called Endoplasmic reticulum aminopeptidase 2.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	883	Total 7212	C 4653	N 1201	O 1328	S 30	0	7	0
1	B	869	Total 7054	C 4550	N 1169	O 1306	S 29	0	2	0

There are 8 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	392	ASN	LYS	variant	UNP Q6P179
A	961	ARG	-	expression tag	UNP Q6P179
A	962	HIS	-	expression tag	UNP Q6P179
A	963	HIS	-	expression tag	UNP Q6P179
B	392	ASN	LYS	variant	UNP Q6P179
B	961	ARG	-	expression tag	UNP Q6P179
B	962	HIS	-	expression tag	UNP Q6P179
B	963	HIS	-	expression tag	UNP Q6P179

- Molecule 2 is an oligosaccharide called alpha-D-mannopyranose-(1-3)-[alpha-D-mannopyranose-(1-6)]beta-D-mannopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose.




Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	Trace
			Total	C	N	O			
2	C	5	Total 61	C 34	N 2	O 25	0	0	0
2	D	5	Total 61	C 34	N 2	O 25	0	0	0
2	G	5	Total 61	C 34	N 2	O 25	0	0	0

- Molecule 3 is an oligosaccharide called 2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose.



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	Trace
			Total	C	N	O			
3	E	2	28	16	2	10	0	0	0
3	F	2	28	16	2	10	0	0	0
3	I	2	28	16	2	10	0	0	0

- Molecule 4 is an oligosaccharide called beta-D-mannopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose.



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	Trace
			Total	C	N	O			
4	H	3	39	22	2	15	0	0	0

- Molecule 5 is ZINC ION (CCD ID: ZN) (formula: Zn) (labeled as "Ligand of Interest" by depositor).


Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
			Total	Zn		
5	A	1	1	1	0	0
5	B	1	1	1	0	0

- Molecule 6 is 2-(N-MORPHOLINO)-ETHANESULFONIC ACID (CCD ID: MES) (formula: C₆H₁₃NO₄S).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
			Total	C	N	O	S		
6	A	1	Total	C	N	O	S	0	0
			12	6	1	4	1		
6	B	1	Total	C	N	O	S	0	0
			12	6	1	4	1		

- Molecule 7 is 4-methoxy-3-{{2-(piperidin-1-yl)-4-(trifluoromethyl)phenyl}sulfamoyl}benzoic acid (CCD ID: J2G) (formula: C₂₀H₂₁F₃N₂O₅S) (labeled as "Ligand of Interest" by depositor).




Mol	Chain	Residues	Atoms						ZeroOcc	AltConf
			Total	C	F	N	O	S		
7	A	1	Total	C	F	N	O	S	0	1
			62	40	6	4	10	2		

Continued on next page...

Continued from previous page...

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	
			Total	C	F	N	O			S
7	B	1	62	40	6	4	10	2	0	1

- Molecule 8 is LYSINE (CCD ID: LYS) (formula: C₆H₁₅N₂O₂) (labeled as "Ligand of Interest" by depositor).




Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
			Total	C	N	O		
8	A	1	10	6	2	2	0	0
8	A	1	10	6	2	2	0	0
8	B	1	10	6	2	2	0	0
8	B	1	10	6	2	2	0	0

- Molecule 9 is 2-acetamido-2-deoxy-beta-D-glucopyranose (CCD ID: NAG) (formula: C₈H₁₅NO₆).




Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
			Total	C	N	O		
9	A	1	Total 14	8	1	5	0	0
9	A	1	Total 14	8	1	5	0	0
9	A	1	Total 14	8	1	5	0	0
9	A	1	Total 14	8	1	5	0	0
9	B	1	Total 14	8	1	5	0	0
9	B	1	Total 14	8	1	5	0	0
9	B	1	Total 14	8	1	5	0	0

- Molecule 10 is IMIDAZOLE (CCD ID: IMD) (formula: C₃H₅N₂).




Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
10	A	1	Total	C	N	0	0
			5	3	2		
10	B	1	Total	C	N	0	0
			5	3	2		

- Molecule 11 is TRIETHYLENE GLYCOL (CCD ID: PGE) (formula: C₆H₁₄O₄).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
11	A	1	Total	C	O	0	0
			10	6	4		

- Molecule 12 is 1,2-ETHANEDIOL (CCD ID: EDO) (formula: C₂H₆O₂).



Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
12	A	1	Total C O 4 2 2	0	0
12	B	1	Total C O 4 2 2	0	0


- Molecule 13 is water.

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
13	A	175	Total O 175 175	0	0
13	B	34	Total O 34 34	0	0


3 Residue-property plots [i](#)


These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: Endoplasmic reticulum aminopeptidase 2



- Molecule 1: Endoplasmic reticulum aminopeptidase 2





- Molecule 2: alpha-D-mannopyranose-(1-3)-[alpha-D-mannopyranose-(1-6)]beta-D-mannopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose

Chain C: 60% 40%



- Molecule 2: alpha-D-mannopyranose-(1-3)-[alpha-D-mannopyranose-(1-6)]beta-D-mannopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose

Chain D: 60% 40%



- Molecule 2: alpha-D-mannopyranose-(1-3)-[alpha-D-mannopyranose-(1-6)]beta-D-mannopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose

Chain G: 60% 40%



- Molecule 3: 2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose

Chain E: 100%



- Molecule 3: 2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose

Chain F: 100%



- Molecule 3: 2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose

Chain I:  100%

MAG1
MAG2

- Molecule 4: beta-D-mannopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose

Chain H:  67% 33%

MAG1
MAG2
BOM3

4 Data and refinement statistics i

Property	Value	Source
Space group	P 1 21 1	Depositor
Cell constants a, b, c, α , β , γ	75.23Å 135.45Å 127.28Å 90.00° 90.11° 90.00°	Depositor
Resolution (Å)	29.29 – 2.45 29.29 – 2.45	Depositor EDS
% Data completeness (in resolution range)	97.1 (29.29-2.45) 87.1 (29.29-2.45)	Depositor EDS
R_{merge}	0.09	Depositor
R_{sym}	0.07	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.17 (at 2.45Å)	Xtrriage
Refinement program	PHENIX 1.20.1_4487	Depositor
R, R_{free}	0.208 , 0.236 0.208 , 0.234	Depositor DCC
R_{free} test set	2027 reflections (2.23%)	wwPDB-VP
Wilson B-factor (Å ²)	48.0	Xtrriage
Anisotropy	0.386	Xtrriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.30 , 48.6	EDS
L-test for twinning ²	$\langle L \rangle = 0.49$, $\langle L^2 \rangle = 0.32$	Xtrriage
Estimated twinning fraction	0.026 for h,-k,-l	Xtrriage
F_o, F_c correlation	0.94	EDS
Total number of atoms	15097	wwPDB-VP
Average B, all atoms (Å ²)	82.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 3.13% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: MES, EDO, BMA, IMD, PGE, J2G, MAN, NAG, ZN

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.08	0/7408	0.24	0/10041
1	B	0.07	0/7233	0.23	0/9808
All	All	0.07	0/14641	0.23	0/19849

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	7212	0	7151	51	0
1	B	7054	0	6957	74	0
2	C	61	0	52	0	0
2	D	61	0	52	0	0
2	G	61	0	52	0	0
3	E	28	0	25	0	0
3	F	28	0	25	0	0
3	I	28	0	25	0	0
4	H	39	0	34	1	0
5	A	1	0	0	0	0
5	B	1	0	0	0	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
6	A	12	0	12	3	0
6	B	12	0	12	2	0
7	A	62	0	0	1	0
7	B	62	0	0	0	0
8	A	20	0	24	2	0
8	B	20	0	24	2	0
9	A	56	0	52	0	0
9	B	42	0	39	0	0
10	A	5	0	5	0	0
10	B	5	0	5	2	0
11	A	10	0	14	2	0
12	A	4	0	6	0	0
12	B	4	0	6	0	0
13	A	175	0	0	2	0
13	B	34	0	0	1	0
All	All	15097	0	14572	129	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 4.

The worst 5 of 129 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:777:TRP:HE1	1:A:783:LYS:HD2	1.42	0.83
1:A:383:MET:HE2	1:A:389:ILE:HA	1.63	0.81
1:A:177[B]:GLU:HB3	1:A:203:GLN:HG2	1.67	0.76
1:B:177[B]:GLU:HB3	1:B:203:GLN:HG2	1.72	0.72
1:B:200:GLU:OE1	8:B:1004:LYS:N	2.23	0.71

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	886/911 (97%)	859 (97%)	27 (3%)	0	100	100
1	B	863/911 (95%)	835 (97%)	28 (3%)	0	100	100
All	All	1749/1822 (96%)	1694 (97%)	55 (3%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	792/819 (97%)	790 (100%)	2 (0%)	86	91
1	B	772/819 (94%)	766 (99%)	6 (1%)	73	83
All	All	1564/1638 (96%)	1556 (100%)	8 (0%)	81	87

5 of 8 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	B	779	GLU
1	B	755	LEU
1	B	598	ASN
1	B	567	LEU
1	B	650	ASN

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 15 such sidechains are listed below:

Mol	Chain	Res	Type
1	B	159	HIS
1	B	689	HIS
1	B	338	ASN
1	B	869	GLN
1	B	544	GLN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

24 monosaccharides are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
2	NAG	C	1	1,2	14,14,15	0.26	0	17,19,21	0.39	0
2	NAG	C	2	2	14,14,15	0.26	0	17,19,21	0.45	0
2	BMA	C	3	2	11,11,12	0.53	0	15,15,17	0.79	0
2	MAN	C	4	2	11,11,12	0.75	0	15,15,17	1.53	2 (13%)
2	MAN	C	5	2	11,11,12	0.56	0	15,15,17	1.00	2 (13%)
2	NAG	D	1	1,2	14,14,15	0.26	0	17,19,21	0.39	0
2	NAG	D	2	2	14,14,15	0.22	0	17,19,21	0.50	0
2	BMA	D	3	2	11,11,12	0.67	0	15,15,17	0.74	0
2	MAN	D	4	2	11,11,12	0.86	1 (9%)	15,15,17	1.56	2 (13%)
2	MAN	D	5	2	11,11,12	0.64	0	15,15,17	1.01	2 (13%)
3	NAG	E	1	1,3	14,14,15	0.23	0	17,19,21	0.43	0
3	NAG	E	2	3	14,14,15	0.24	0	17,19,21	0.42	0
3	NAG	F	1	1,3	14,14,15	0.20	0	17,19,21	0.40	0
3	NAG	F	2	3	14,14,15	0.23	0	17,19,21	0.43	0
2	NAG	G	1	1,2	14,14,15	0.26	0	17,19,21	0.39	0
2	NAG	G	2	2	14,14,15	0.30	0	17,19,21	0.46	0
2	BMA	G	3	2	11,11,12	0.59	0	15,15,17	0.66	0
2	MAN	G	4	2	11,11,12	0.90	1 (9%)	15,15,17	1.58	2 (13%)
2	MAN	G	5	2	11,11,12	0.65	0	15,15,17	1.02	2 (13%)
4	NAG	H	1	1,4	14,14,15	0.17	0	17,19,21	0.42	0
4	NAG	H	2	4	14,14,15	0.25	0	17,19,21	0.46	0

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
4	BMA	H	3	4	11,11,12	0.61	0	15,15,17	0.65	0
3	NAG	I	1	1,3	14,14,15	0.26	0	17,19,21	0.46	0
3	NAG	I	2	3	14,14,15	0.25	0	17,19,21	0.41	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	NAG	C	1	1,2	-	0/6/23/26	0/1/1/1
2	NAG	C	2	2	-	0/6/23/26	0/1/1/1
2	BMA	C	3	2	-	0/2/19/22	0/1/1/1
2	MAN	C	4	2	-	2/2/19/22	0/1/1/1
2	MAN	C	5	2	-	0/2/19/22	0/1/1/1
2	NAG	D	1	1,2	-	0/6/23/26	0/1/1/1
2	NAG	D	2	2	-	2/6/23/26	0/1/1/1
2	BMA	D	3	2	-	0/2/19/22	0/1/1/1
2	MAN	D	4	2	-	0/2/19/22	0/1/1/1
2	MAN	D	5	2	-	1/2/19/22	0/1/1/1
3	NAG	E	1	1,3	-	0/6/23/26	0/1/1/1
3	NAG	E	2	3	-	2/6/23/26	0/1/1/1
3	NAG	F	1	1,3	-	0/6/23/26	0/1/1/1
3	NAG	F	2	3	-	2/6/23/26	0/1/1/1
2	NAG	G	1	1,2	-	0/6/23/26	0/1/1/1
2	NAG	G	2	2	-	0/6/23/26	0/1/1/1
2	BMA	G	3	2	-	0/2/19/22	0/1/1/1
2	MAN	G	4	2	-	2/2/19/22	0/1/1/1
2	MAN	G	5	2	-	2/2/19/22	0/1/1/1
4	NAG	H	1	1,4	-	0/6/23/26	0/1/1/1
4	NAG	H	2	4	-	0/6/23/26	0/1/1/1
4	BMA	H	3	4	-	0/2/19/22	0/1/1/1
3	NAG	I	1	1,3	-	0/6/23/26	0/1/1/1
3	NAG	I	2	3	-	1/6/23/26	0/1/1/1

All (2) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	D	4	MAN	C1-C2	2.32	1.57	1.52
2	G	4	MAN	C1-C2	2.31	1.57	1.52

The worst 5 of 12 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	G	4	MAN	C1-O5-C5	5.14	119.07	112.19
2	D	4	MAN	C1-O5-C5	4.99	118.88	112.19
2	C	4	MAN	C1-O5-C5	4.85	118.68	112.19
2	G	5	MAN	C1-O5-C5	2.81	115.95	112.19
2	D	5	MAN	C1-O5-C5	2.71	115.82	112.19

There are no chirality outliers.

5 of 14 torsion outliers are listed below:


Mol	Chain	Res	Type	Atoms
2	G	4	MAN	C4-C5-C6-O6
2	G	4	MAN	O5-C5-C6-O6
3	F	2	NAG	O5-C5-C6-O6
3	F	2	NAG	C4-C5-C6-O6
2	D	5	MAN	O5-C5-C6-O6


There are no ring outliers.


1 monomer is involved in 1 short contact:


Mol	Chain	Res	Type	Clashes	Symm-Clashes
4	H	1	NAG	1	0


The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for oligosaccharide.

















5.6 Ligand geometry [i](#)

Of 24 ligands modelled in this entry, 2 are monoatomic - leaving 22 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
6	MES	B	1002	-	12,12,12	2.31	1 (8%)	15,16,16	1.87	3 (20%)
8	LYS	B	1010	-	8,9,9	0.85	1 (12%)	7,10,10	0.94	1 (14%)
12	EDO	B	1009	-	3,3,3	0.43	0	2,2,2	0.37	0
12	EDO	A	1012	-	3,3,3	0.43	0	2,2,2	0.38	0
9	NAG	A	1007	1	14,14,15	0.31	0	17,19,21	0.47	0
9	NAG	A	1008	1	14,14,15	0.33	0	17,19,21	0.47	0
11	PGE	A	1011	-	9,9,9	0.31	0	8,8,8	0.28	0
9	NAG	B	1007	1	14,14,15	0.25	0	17,19,21	0.48	0
9	NAG	B	1005	1	14,14,15	0.22	0	17,19,21	0.42	0

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
8	LYS	A	1004	5	8,9,9	0.85	1 (12%)	7,10,10	1.07	1 (14%)
9	NAG	A	1006	1	14,14,15	0.24	0	17,19,21	0.46	0
10	IMD	B	1008	-	5,5,5	0.63	0	5,5,5	0.44	0
8	LYS	B	1004	5	8,9,9	0.85	1 (12%)	7,10,10	1.08	1 (14%)
9	NAG	A	1005	1	14,14,15	0.24	0	17,19,21	0.44	0
7	J2G	A	1003[A]	-	33,33,33	0.94	2 (6%)	48,49,49	1.26	7 (14%)
8	LYS	A	1009	-	8,9,9	0.87	1 (12%)	7,10,10	0.95	1 (14%)
7	J2G	A	1003[B]	-	33,33,33	0.94	2 (6%)	48,49,49	1.26	4 (8%)
6	MES	A	1002	-	12,12,12	2.37	1 (8%)	15,16,16	1.73	5 (33%)
9	NAG	B	1006	1	14,14,15	0.24	0	17,19,21	0.44	0
7	J2G	B	1003[A]	-	33,33,33	0.93	2 (6%)	48,49,49	1.25	7 (14%)
10	IMD	A	1010	-	5,5,5	0.64	0	5,5,5	0.43	0
7	J2G	B	1003[B]	-	33,33,33	0.93	1 (3%)	48,49,49	1.26	4 (8%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. ^{1,2} means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
6	MES	B	1002	-	-	0/6/14/14	0/1/1/1
8	LYS	B	1010	-	-	2/9/9/9	-
12	EDO	B	1009	-	-	0/1/1/1	-
12	EDO	A	1012	-	-	0/1/1/1	-
9	NAG	A	1007	1	-	2/6/23/26	0/1/1/1
9	NAG	A	1008	1	-	0/6/23/26	0/1/1/1
11	PGE	A	1011	-	-	3/7/7/7	-
9	NAG	B	1007	1	-	4/6/23/26	0/1/1/1
9	NAG	B	1005	1	-	2/6/23/26	0/1/1/1
8	LYS	A	1004	5	-	4/9/9/9	-
9	NAG	A	1006	1	-	2/6/23/26	0/1/1/1
10	IMD	B	1008	-	-	-	0/1/1/1
8	LYS	B	1004	5	-	4/9/9/9	-
9	NAG	A	1005	1	-	2/6/23/26	0/1/1/1
7	J2G	A	1003[A]	-	-	18/27/35/35	0/3/3/3
8	LYS	A	1009	-	-	2/9/9/9	-
7	J2G	A	1003[B]	-	-	13/27/35/35	0/3/3/3
6	MES	A	1002	-	-	3/6/14/14	0/1/1/1

Continued on next page...

Continued from previous page...

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
9	NAG	B	1006	1	-	1/6/23/26	0/1/1/1
7	J2G	B	1003[A]	-	-	17/27/35/35	0/3/3/3
10	IMD	A	1010	-	-	-	0/1/1/1
7	J2G	B	1003[B]	-	-	13/27/35/35	0/3/3/3

The worst 5 of 13 bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
6	A	1002	MES	C8-S	-7.92	1.66	1.77
6	B	1002	MES	C8-S	-7.71	1.66	1.77
7	A	1003[A]	J2G	SAG-NAH	2.61	1.67	1.63
7	B	1003[A]	J2G	SAG-NAH	2.55	1.67	1.63
7	B	1003[B]	J2G	OAB-CAE	-2.37	1.33	1.37

The worst 5 of 34 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
6	B	1002	MES	C5-N4-C3	4.09	117.66	108.84
7	A	1003[B]	J2G	CAF-SAG-NAH	-3.86	102.76	107.30
7	B	1003[B]	J2G	CBC-NAZ-CBA	3.63	119.73	111.57
7	B	1003[B]	J2G	OAC-SAG-CAF	3.57	113.58	107.68
6	A	1002	MES	C5-N4-C3	3.49	116.37	108.84

There are no chirality outliers.

5 of 92 torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
6	A	1002	MES	C7-C8-S-O2S
6	A	1002	MES	C7-C8-S-O3S
7	A	1003[A]	J2G	CAE-CAF-SAG-NAH
7	A	1003[A]	J2G	CAE-CAF-SAG-OAC
7	A	1003[A]	J2G	CAE-CAF-SAG-OAP

There are no ring outliers.

8 monomers are involved in 13 short contacts:


Mol	Chain	Res	Type	Clashes	Symm-Clashes
6	B	1002	MES	2	0
11	A	1011	PGE	2	0
8	A	1004	LYS	1	0



Continued on next page...


Continued from previous page...


Mol	Chain	Res	Type	Clashes	Symm-Clashes
10	B	1008	IMD	2	0
8	B	1004	LYS	2	0
8	A	1009	LYS	1	0
7	A	1003[B]	J2G	1	0
6	A	1002	MES	3	0


The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.














5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	883/911 (96%)	0.06	12 (1%) 73 75	28, 59, 95, 142	7 (0%)
1	B	869/911 (95%)	0.81	75 (8%) 16 14	42, 98, 158, 198	2 (0%)
All	All	1752/1822 (96%)	0.43	87 (4%) 34 32	28, 73, 144, 198	9 (0%)

The worst 5 of 87 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	B	783	LYS	5.8
1	A	531	ASN	4.5
1	B	578	ARG	4.4
1	A	963	HIS	4.4
1	A	784	LEU	3.9

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled ‘Q < 0.9’ lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
2	MAN	D	4	11/12	0.43	0.16	141,143,147,148	0
2	BMA	D	3	11/12	0.47	0.15	114,120,135,140	0
2	BMA	G	3	11/12	0.52	0.16	104,109,118,123	0
4	BMA	H	3	11/12	0.52	0.14	136,139,143,143	0
2	MAN	G	5	11/12	0.53	0.18	72,96,106,108	0

Continued on next page...




Continued from previous page...

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(\AA^2)	Q<0.9
2	MAN	C	4	11/12	0.54	0.12	124,128,132,134	0
2	MAN	C	5	11/12	0.55	0.14	98,112,119,121	0
2	MAN	G	4	11/12	0.57	0.11	129,132,135,136	0
2	MAN	D	5	11/12	0.60	0.16	91,103,112,112	0
2	BMA	C	3	11/12	0.61	0.14	114,117,123,128	0
3	NAG	I	2	14/15	0.66	0.15	135,142,143,143	0
3	NAG	F	2	14/15	0.66	0.12	122,136,140,141	0
3	NAG	E	2	14/15	0.68	0.15	106,117,121,121	0
4	NAG	H	2	14/15	0.69	0.13	113,121,126,132	0
3	NAG	I	1	14/15	0.70	0.17	112,125,134,139	0
4	NAG	H	1	14/15	0.71	0.15	89,100,111,114	0
3	NAG	F	1	14/15	0.72	0.14	96,105,123,130	0
2	NAG	C	2	14/15	0.83	0.13	68,86,99,102	0
3	NAG	E	1	14/15	0.83	0.11	71,87,102,111	0
2	NAG	G	2	14/15	0.84	0.12	63,80,93,100	0
2	NAG	D	2	14/15	0.88	0.11	65,76,89,108	0
2	NAG	G	1	14/15	0.91	0.09	52,63,73,83	0
2	NAG	C	1	14/15	0.92	0.08	53,64,76,78	0
2	NAG	D	1	14/15	0.94	0.07	40,57,62,62	0

The following is a graphical depiction of the model fit to experimental electron density for oligosaccharide. Each fit is shown from different orientation to approximate a three-dimensional view.




Electron density around Chain C:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)






Electron density around Chain D:


$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)



Electron density around Chain G:




$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)









Electron density around Chain F:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)

**Electron density around Chain I:**

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)





6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q < 0.9' lists the number of atoms with occupancy less than 0.9.


Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
9	NAG	B	1006	14/15	0.54	0.15	169,174,181,182	0
9	NAG	B	1005	14/15	0.59	0.16	143,153,157,157	0
9	NAG	A	1008	14/15	0.61	0.15	107,127,134,144	0
9	NAG	B	1007	14/15	0.68	0.20	188,196,199,200	0
8	LYS	B	1010	10/10	0.69	0.25	83,90,93,93	10
9	NAG	A	1005	14/15	0.69	0.16	119,128,137,141	0
11	PGE	A	1011	10/10	0.69	0.18	55,66,72,82	10
9	NAG	A	1006	14/15	0.70	0.12	89,106,117,118	0
10	IMD	B	1008	5/5	0.72	0.27	95,97,103,111	5
9	NAG	A	1007	14/15	0.73	0.18	118,121,127,131	0
10	IMD	A	1010	5/5	0.74	0.25	54,59,62,64	5

Continued on next page...

Continued from previous page...



Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
7	J2G	A	1003[B]	31/31	0.75	0.23	50,58,62,63	31
7	J2G	A	1003[A]	31/31	0.75	0.23	50,59,63,64	31
8	LYS	A	1009	10/10	0.76	0.18	63,68,71,72	10
7	J2G	B	1003[A]	31/31	0.79	0.23	81,86,89,89	31
7	J2G	B	1003[B]	31/31	0.79	0.23	81,85,88,90	31
8	LYS	B	1004	10/10	0.81	0.17	82,85,88,90	0
12	EDO	B	1009	4/4	0.85	0.26	98,103,108,112	0
6	MES	B	1002	12/12	0.86	0.18	82,87,88,100	12
6	MES	A	1002	12/12	0.89	0.16	51,58,71,81	12
12	EDO	A	1012	4/4	0.91	0.15	64,67,68,77	0
8	LYS	A	1004	10/10	0.93	0.13	49,51,61,65	0
5	ZN	B	1001	1/1	0.98	0.05	73,73,73,73	0
5	ZN	A	1001	1/1	1.00	0.04	41,41,41,41	0

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.






Electron density around J2G A 1003 (B):

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)






Electron density around J2G A 1003 (A):

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)




**Electron density around LYS A 1009:**

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)






Electron density around J2G B 1003 (A):

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)






Electron density around J2G B 1003 (B):

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)






Electron density around LYS B 1004:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)




**Electron density around LYS A 1004:**


$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)



Electron density around ZN B 1001:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)





6.5 Other polymers ⓘ

There are no such residues in this entry.