



wwPDB X-ray Structure Validation Summary Report ⓘ

Mar 9, 2026 – 06:05 PM UTC

PDB ID : 1EWD / pdb_00001ewd
Title : FRUCTOSE 1,6-BISPHOSPHATE ALDOLASE FROM RABBIT MUSCLE
Authors : Maurady, A.; Sygusch, J.
Deposited on : 2000-04-25
Resolution : 2.46 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4-5-2 with Phenix2.0
Xtriage (Phenix) : 2.0
EDS : 3.0
Percentile statistics : 20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4 : 9.0.010 (Gargrove)
Density-Fitness : 1.0.12
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.46 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	1190 (2.46-2.46)
Clashscore	190562	1229 (2.46-2.46)
Ramachandran outliers	187476	1218 (2.46-2.46)
Sidechain outliers	187428	1218 (2.46-2.46)
RSRZ outliers	180081	1190 (2.46-2.46)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	363	2% 68% 28% ..
1	B	363	2% 68% 29% .
1	C	363	% 67% 31% .
1	D	363	% 69% 28% .

2 Entry composition

There are 2 unique types of molecules in this entry. The entry contains 13358 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called FRUCTOSE 1,6-BISPHOSPHATE ALDOLASE.


Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	363	Total	C	N	O	S	0	0	0
			2756	1730	488	526	12			
1	B	363	Total	C	N	O	S	0	0	0
			2756	1730	488	526	12			
1	C	363	Total	C	N	O	S	0	0	0
			2756	1730	488	526	12			
1	D	363	Total	C	N	O	S	0	0	0
			2756	1730	488	526	12			

There are 8 discrepancies between the modelled and reference sequences:


Chain	Residue	Modelled	Actual	Comment	Reference
A	107	MET	LYS	engineered mutation	UNP P00883
A	344	SER	PRO	conflict	UNP P00883
B	107	MET	LYS	engineered mutation	UNP P00883
B	344	SER	PRO	conflict	UNP P00883
C	107	MET	LYS	engineered mutation	UNP P00883
C	344	SER	PRO	conflict	UNP P00883
D	107	MET	LYS	engineered mutation	UNP P00883
D	344	SER	PRO	conflict	UNP P00883

- Molecule 2 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
2	A	544	Total	O	0	0
			544	544		
2	B	494	Total	O	0	0
			494	494		
2	C	676	Total	O	0	0
			676	676		
2	D	620	Total	O	0	0
			620	620		



● Molecule 1: FRUCTOSE 1,6-BISPHOSPHATE ALDOLASE



4 Data and refinement statistics

Property	Value	Source
Space group	P 1 21 1	Depositor
Cell constants a, b, c, α , β , γ	164.19Å 57.58Å 85.35Å 90.00° 102.66° 90.00°	Depositor
Resolution (Å)	10.00 – 2.46 10.00 – 2.46	Depositor EDS
% Data completeness (in resolution range)	8.0 (10.00-2.46) 66.9 (10.00-2.46)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	3.83 (at 2.48Å)	Xtriage
Refinement program	X-PLOR 3.1	Depositor
R, R_{free}	0.170 , 0.234 0.165 , 0.225	Depositor DCC
R_{free} test set	3151 reflections (8.10%)	wwPDB-VP
Wilson B-factor (Å ²)	11.0	Xtriage
Anisotropy	1.275	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.23 , 100.5	EDS
L-test for twinning ²	$\langle L \rangle = 0.47$, $\langle L^2 \rangle = 0.30$	Xtriage
Estimated twinning fraction	0.012 for -h-l,-k,l	Xtriage
F_o, F_c correlation	0.93	EDS
Total number of atoms	13358	wwPDB-VP
Average B, all atoms (Å ²)	19.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 5.16% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.40	0/2809	0.94	13/3805 (0.3%)
1	B	0.41	0/2809	0.96	9/3805 (0.2%)
1	C	0.41	0/2809	0.90	4/3805 (0.1%)
1	D	0.40	0/2809	0.92	5/3805 (0.1%)
All	All	0.40	0/11236	0.93	31/15220 (0.2%)

There are no bond length outliers.

The worst 5 of 31 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	362	ALA	N-CA-C	-7.21	103.04	112.41
1	A	191	LEU	N-CA-C	6.84	117.99	109.57
1	B	289	CYS	CA-C-N	6.63	127.16	119.47
1	B	289	CYS	C-N-CA	6.63	127.16	119.47
1	D	300	SER	N-CA-C	-6.53	96.89	110.80

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2756	0	2770	93	0
1	B	2756	0	2770	89	0
1	C	2756	0	2770	114	0
1	D	2756	0	2770	87	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
2	A	544	0	0	35	0
2	B	494	0	0	26	0
2	C	676	0	0	51	0
2	D	620	0	0	40	0
All	All	13358	0	11080	371	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 17.

The worst 5 of 371 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:C:80:HIS:HA	2:C:3465:HOH:O	1.76	0.84
1:A:94:PRO:HD3	2:A:1852:HOH:O	1.80	0.81
1:A:24:ALA:HB3	1:A:27:LYS:HD3	1.63	0.80
1:C:24:ALA:HB3	1:C:27:LYS:HD2	1.63	0.80
1:C:307:ALA:HA	2:C:3937:HOH:O	1.81	0.80

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	361/363 (99%)	331 (92%)	26 (7%)	4 (1%)	11	12
1	B	361/363 (99%)	332 (92%)	23 (6%)	6 (2%)	7	6
1	C	361/363 (99%)	325 (90%)	34 (9%)	2 (1%)	21	27
1	D	361/363 (99%)	337 (93%)	19 (5%)	5 (1%)	9	8
All	All	1444/1452 (99%)	1325 (92%)	102 (7%)	17 (1%)	10	11

5 of 17 Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	361	HIS
1	B	353	SER
1	B	354	GLU
1	C	345	SER
1	D	5	PRO

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	291/291 (100%)	276 (95%)	15 (5%)	21	30
1	B	291/291 (100%)	279 (96%)	12 (4%)	27	40
1	C	291/291 (100%)	279 (96%)	12 (4%)	27	40
1	D	291/291 (100%)	274 (94%)	17 (6%)	18	27
All	All	1164/1164 (100%)	1108 (95%)	56 (5%)	23	34

5 of 56 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	C	106	ILE
1	D	347	GLN
1	C	191	LEU
1	D	332	LEU
1	D	207	LYS

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 28 such sidechains are listed below:

Mol	Chain	Res	Type
1	C	156	HIS
1	D	334	ASN
1	C	237	HIS
1	D	202	GLN
1	C	220	HIS

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

There are no ligands in this entry.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	363/363 (100%)	-0.40	7 (1%) 66 67	2, 7, 60, 92	0
1	B	363/363 (100%)	-0.40	9 (2%) 58 61	2, 6, 55, 87	0
1	C	363/363 (100%)	-0.46	5 (1%) 73 75	2, 6, 48, 81	0
1	D	363/363 (100%)	-0.41	2 (0%) 85 87	2, 8, 51, 94	0
All	All	1452/1452 (100%)	-0.42	23 (1%) 70 73	2, 7, 55, 94	0

The worst 5 of 23 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	B	350	ALA	4.1
1	B	355	SER	3.9
1	A	359	SER	3.4
1	B	351	ALA	3.3
1	B	348	ALA	3.2

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.4 Ligands [i](#)

There are no ligands in this entry.

6.5 Other polymers [i](#)

There are no such residues in this entry.