



wwPDB X-ray Structure Validation Summary Report ⓘ

Mar 7, 2026 – 03:50 AM UTC

PDB ID : 2F2S / pdb_00002f2s
Title : Human mitochondrial acetoacetyl-CoA thiolase
Authors : Min, J.R.; Dombrowski, L.; Antoshenko, T.; Wu, H.; Loppnau, P.; Weigelt, J.; Sundstrom, M.; Arrowsmith, C.H.; Edwards, A.M.; Bochkarev, A.; Plotnikov, A.N.; Structural Genomics Consortium (SGC)
Deposited on : 2005-11-17
Resolution : 2.00 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4-5-2 with Phenix2.0
Mogul : 2022.3.0, CSD as543be (2022)
Xtriage (Phenix) : 2.0
EDS : 3.0
Buster-report : wwPDB partial adaption of 1.1.7 (2018)
Percentile statistics : 20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4 : 9.0.010 (Gargrove)
Density-Fitness : 1.0.12
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.00 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	10052 (2.00-2.00)
Clashscore	190562	11152 (2.00-2.00)
Ramachandran outliers	187476	11031 (2.00-2.00)
Sidechain outliers	187428	11029 (2.00-2.00)
RSRZ outliers	180081	10067 (2.00-2.00)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	406	 4% 69% 23% . . .
1	B	406	 75% 17% . 5%
1	C	406	 2% 71% 21% . .
1	D	406	 % 70% 21% . . 6%

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
1	SCY	A	126	-	-	X	-
1	SCY	B	126	-	-	X	-
1	SCY	C	126	-	-	X	-
1	SCY	D	126	-	-	X	-

2 Entry composition i

There are 4 unique types of molecules in this entry. The entry contains 12020 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Acetyl-CoA acetyltransferase, mitochondrial.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	389	2834	1783	486	544	21	0	0	0
1	B	386	2818	1773	484	540	21	0	0	0
1	C	388	2841	1789	487	544	21	0	0	0
1	D	382	2801	1765	478	537	21	0	0	0

There are 80 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	22	MET	-	cloning artifact	UNP P24752
A	23	GLY	-	cloning artifact	UNP P24752
A	24	SER	-	cloning artifact	UNP P24752
A	25	SER	-	cloning artifact	UNP P24752
A	26	HIS	-	expression tag	UNP P24752
A	27	HIS	-	expression tag	UNP P24752
A	28	HIS	-	expression tag	UNP P24752
A	29	HIS	-	expression tag	UNP P24752
A	30	HIS	-	expression tag	UNP P24752
A	31	HIS	-	expression tag	UNP P24752
A	32	SER	-	cloning artifact	UNP P24752
A	33	SER	-	cloning artifact	UNP P24752
A	34	GLY	-	cloning artifact	UNP P24752
A	35	LEU	-	cloning artifact	UNP P24752
A	36	VAL	-	cloning artifact	UNP P24752
A	37	PRO	-	cloning artifact	UNP P24752
A	38	ARG	-	cloning artifact	UNP P24752
A	39	GLY	-	cloning artifact	UNP P24752
A	40	SER	-	cloning artifact	UNP P24752
A	126	SCY	CYS	modified residue	UNP P24752
B	22	MET	-	cloning artifact	UNP P24752

Continued on next page...

Continued from previous page...

Chain	Residue	Modelled	Actual	Comment	Reference
B	23	GLY	-	cloning artifact	UNP P24752
B	24	SER	-	cloning artifact	UNP P24752
B	25	SER	-	cloning artifact	UNP P24752
B	26	HIS	-	expression tag	UNP P24752
B	27	HIS	-	expression tag	UNP P24752
B	28	HIS	-	expression tag	UNP P24752
B	29	HIS	-	expression tag	UNP P24752
B	30	HIS	-	expression tag	UNP P24752
B	31	HIS	-	expression tag	UNP P24752
B	32	SER	-	cloning artifact	UNP P24752
B	33	SER	-	cloning artifact	UNP P24752
B	34	GLY	-	cloning artifact	UNP P24752
B	35	LEU	-	cloning artifact	UNP P24752
B	36	VAL	-	cloning artifact	UNP P24752
B	37	PRO	-	cloning artifact	UNP P24752
B	38	ARG	-	cloning artifact	UNP P24752
B	39	GLY	-	cloning artifact	UNP P24752
B	40	SER	-	cloning artifact	UNP P24752
B	126	SCY	CYS	modified residue	UNP P24752
C	22	MET	-	cloning artifact	UNP P24752
C	23	GLY	-	cloning artifact	UNP P24752
C	24	SER	-	cloning artifact	UNP P24752
C	25	SER	-	cloning artifact	UNP P24752
C	26	HIS	-	expression tag	UNP P24752
C	27	HIS	-	expression tag	UNP P24752
C	28	HIS	-	expression tag	UNP P24752
C	29	HIS	-	expression tag	UNP P24752
C	30	HIS	-	expression tag	UNP P24752
C	31	HIS	-	expression tag	UNP P24752
C	32	SER	-	cloning artifact	UNP P24752
C	33	SER	-	cloning artifact	UNP P24752
C	34	GLY	-	cloning artifact	UNP P24752
C	35	LEU	-	cloning artifact	UNP P24752
C	36	VAL	-	cloning artifact	UNP P24752
C	37	PRO	-	cloning artifact	UNP P24752
C	38	ARG	-	cloning artifact	UNP P24752
C	39	GLY	-	cloning artifact	UNP P24752
C	40	SER	-	cloning artifact	UNP P24752
C	126	SCY	CYS	modified residue	UNP P24752
D	22	MET	-	cloning artifact	UNP P24752
D	23	GLY	-	cloning artifact	UNP P24752
D	24	SER	-	cloning artifact	UNP P24752

Continued on next page...


Continued from previous page...

Chain	Residue	Modelled	Actual	Comment	Reference
D	25	SER	-	cloning artifact	UNP P24752
D	26	HIS	-	expression tag	UNP P24752
D	27	HIS	-	expression tag	UNP P24752
D	28	HIS	-	expression tag	UNP P24752
D	29	HIS	-	expression tag	UNP P24752
D	30	HIS	-	expression tag	UNP P24752
D	31	HIS	-	expression tag	UNP P24752
D	32	SER	-	cloning artifact	UNP P24752
D	33	SER	-	cloning artifact	UNP P24752
D	34	GLY	-	cloning artifact	UNP P24752
D	35	LEU	-	cloning artifact	UNP P24752
D	36	VAL	-	cloning artifact	UNP P24752
D	37	PRO	-	cloning artifact	UNP P24752
D	38	ARG	-	cloning artifact	UNP P24752
D	39	GLY	-	cloning artifact	UNP P24752
D	40	SER	-	cloning artifact	UNP P24752
D	126	SCY	CYS	modified residue	UNP P24752

- Molecule 2 is CHLORIDE ION (CCD ID: CL) (formula: Cl).

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
2	A	1	Total Cl 1 1	0	0
2	B	2	Total Cl 2 2	0	0

- Molecule 3 is COENZYME A (CCD ID: COA) (formula: C₂₁H₃₆N₇O₁₆P₃S).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	
3	A	1	Total	C	N	O	P	S	0	0
			48	21	7	16	3	1		
3	B	1	Total	C	N	O	P	S	0	0
			48	21	7	16	3	1		
3	C	1	Total	C	N	O	P	S	0	0
			48	21	7	16	3	1		
3	D	1	Total	C	N	O	P	S	0	0
			48	21	7	16	3	1		


- Molecule 4 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	A	99	Total	O	0	0
			99	99		
4	B	172	Total	O	0	0
			172	172		
4	C	137	Total	O	0	0
			137	137		
4	D	123	Total	O	0	0
			123	123		


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: Acetyl-CoA acetyltransferase, mitochondrial



- Molecule 1: Acetyl-CoA acetyltransferase, mitochondrial



- Molecule 1: Acetyl-CoA acetyltransferase, mitochondrial



4 Data and refinement statistics

Property	Value	Source
Space group	P 1 21 1	Depositor
Cell constants a, b, c, α , β , γ	56.99Å 126.64Å 111.86Å 90.00° 98.64° 90.00°	Depositor
Resolution (Å)	33.80 – 2.00 33.80 – 2.00	Depositor EDS
% Data completeness (in resolution range)	98.9 (33.80-2.00) 98.9 (33.80-2.00)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.19 (at 2.00Å)	Xtrriage
Refinement program	REFMAC 5.2.0005	Depositor
R, R_{free}	0.199 , 0.257 0.202 , 0.196	Depositor DCC
R_{free} test set	5269 reflections (4.99%)	wwPDB-VP
Wilson B-factor (Å ²)	40.3	Xtrriage
Anisotropy	0.054	Xtrriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.34 , 37.4	EDS
L-test for twinning ²	$\langle L \rangle = 0.49$, $\langle L^2 \rangle = 0.32$	Xtrriage
Estimated twinning fraction	No twinning to report.	Xtrriage
F_o, F_c correlation	0.96	EDS
Total number of atoms	12020	wwPDB-VP
Average B, all atoms (Å ²)	43.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 6.50% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: SCY, CL, COA

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	1.00	1/2864 (0.0%)	1.18	6/3876 (0.2%)
1	B	1.26	9/2848 (0.3%)	1.25	14/3851 (0.4%)
1	C	1.17	6/2871 (0.2%)	1.22	10/3883 (0.3%)
1	D	1.05	1/2830 (0.0%)	1.18	11/3828 (0.3%)
All	All	1.12	17/11413 (0.1%)	1.21	41/15438 (0.3%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	A	0	1
1	B	0	1
1	D	0	1
All	All	0	3

The worst 5 of 17 bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	B	236	ILE	CA-CB	8.92	1.63	1.53
1	B	71	ALA	CA-CB	7.43	1.64	1.53
1	B	264	VAL	CA-CB	7.39	1.57	1.54
1	D	81	ILE	CA-CB	7.31	1.61	1.55
1	A	147	VAL	CA-CB	6.99	1.64	1.54

The worst 5 of 41 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	C	246	PRO	N-CA-CB	7.07	110.78	103.00

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	B	113	LEU	CA-C-N	-7.01	112.54	119.76
1	B	113	LEU	C-N-CA	-7.01	112.54	119.76
1	A	58	SER	N-CA-C	6.87	119.61	111.71
1	A	50	THR	CA-C-N	-6.83	113.27	120.03

There are no chirality outliers.

All (3) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	A	386	PRO	Peptide
1	B	125	VAL	Mainchain
1	D	425	GLN	Peptide

5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2834	0	2887	70	0
1	B	2818	0	2882	63	0
1	C	2841	0	2910	65	0
1	D	2801	0	2869	82	0
2	A	1	0	0	0	0
2	B	2	0	0	0	0
3	A	48	0	32	2	0
3	B	48	0	32	5	0
3	C	48	0	32	2	0
3	D	48	0	32	0	0
4	A	99	0	0	0	0
4	B	172	0	0	7	0
4	C	137	0	0	5	0
4	D	123	0	0	6	0
All	All	12020	0	11676	267	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 12.

The worst 5 of 267 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:125:VAL:C	1:A:126:SCY:H2	1.21	1.49
1:B:131:LYS:HG3	1:B:135:MET:CE	1.86	1.06
1:B:131:LYS:CG	1:B:135:MET:HE2	1.87	1.04
1:D:426:LYS:O	1:D:426:LYS:HD3	1.58	1.03
1:D:122:ILE:HG13	1:D:135:MET:HE1	1.41	1.01

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	384/406 (95%)	370 (96%)	11 (3%)	3 (1%)	16	11
1	B	381/406 (94%)	369 (97%)	10 (3%)	2 (0%)	24	21
1	C	383/406 (94%)	362 (94%)	18 (5%)	3 (1%)	16	11
1	D	377/406 (93%)	367 (97%)	9 (2%)	1 (0%)	36	35
All	All	1525/1624 (94%)	1468 (96%)	48 (3%)	9 (1%)	21	17

5 of 9 Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	125	VAL
1	A	340	VAL
1	C	203	LYS
1	A	273	LYS
1	C	242	VAL

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	296/317 (93%)	263 (89%)	33 (11%)	6	3
1	B	296/317 (93%)	285 (96%)	11 (4%)	30	30
1	C	299/317 (94%)	272 (91%)	27 (9%)	9	6
1	D	295/317 (93%)	269 (91%)	26 (9%)	9	6
All	All	1186/1268 (94%)	1089 (92%)	97 (8%)	10	7

5 of 97 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	C	327	ILE
1	D	76	ILE
1	C	336	VAL
1	C	369	ILE
1	D	125	VAL

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 23 such sidechains are listed below:

Mol	Chain	Res	Type
1	C	305	ASN
1	D	209	ASN
1	C	425	GLN
1	D	272	GLN
1	B	211	GLN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

4 non-standard protein/DNA/RNA residues are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the

expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
1	SCY	A	126	1	7,8,9	1.26	1 (14%)	4,9,11	0.41	0
1	SCY	C	126	1	7,8,9	0.97	0	4,9,11	1.17	0
1	SCY	B	126	1	7,8,9	0.99	0	4,9,11	0.82	0
1	SCY	D	126	1	7,8,9	0.92	0	4,9,11	0.77	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
1	SCY	A	126	1	-	3/5/7/9	-
1	SCY	C	126	1	-	2/5/7/9	-
1	SCY	B	126	1	-	3/5/7/9	-
1	SCY	D	126	1	-	2/5/7/9	-

All (1) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	126	SCY	CB-SG	-2.35	1.76	1.81

There are no bond angle outliers.

There are no chirality outliers.

5 of 10 torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
1	A	126	SCY	OCD-CD-SG-CB
1	A	126	SCY	CE-CD-SG-CB
1	B	126	SCY	OCD-CD-SG-CB
1	B	126	SCY	CE-CD-SG-CB
1	C	126	SCY	OCD-CD-SG-CB

There are no ring outliers.

4 monomers are involved in 37 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
1	A	126	SCY	10	0

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Clashes	Symm-Clashes
1	C	126	SCY	7	0
1	B	126	SCY	9	0
1	D	126	SCY	11	0

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

Of 7 ligands modelled in this entry, 3 are monoatomic - leaving 4 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
3	COA	C	1003	-	47,50,50	1.80	6 (12%)	69,75,75	1.52	9 (13%)
3	COA	D	1004	-	47,50,50	1.72	6 (12%)	69,75,75	1.52	9 (13%)
3	COA	A	1001	-	47,50,50	1.73	6 (12%)	69,75,75	1.61	10 (14%)
3	COA	B	1002	-	47,50,50	1.80	6 (12%)	69,75,75	1.68	12 (17%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	COA	C	1003	-	-	9/48/64/64	0/3/3/3
3	COA	D	1004	-	-	10/48/64/64	0/3/3/3
3	COA	A	1001	-	-	14/48/64/64	0/3/3/3
3	COA	B	1002	-	-	6/48/64/64	0/3/3/3

The worst 5 of 24 bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
3	A	1001	COA	O9P-C9P	9.32	1.41	1.23
3	C	1003	COA	O9P-C9P	9.12	1.40	1.23
3	B	1002	COA	O9P-C9P	9.02	1.40	1.23
3	D	1004	COA	O9P-C9P	9.00	1.40	1.23
3	B	1002	COA	P1A-O3A	4.42	1.64	1.59

The worst 5 of 40 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	C	1003	COA	N3A-C2A-N1A	-6.03	119.46	128.58
3	D	1004	COA	N3A-C2A-N1A	-5.48	120.29	128.58
3	A	1001	COA	N3A-C2A-N1A	-5.39	120.43	128.58
3	B	1002	COA	N3A-C2A-N1A	-5.26	120.62	128.58
3	A	1001	COA	C5A-C4A-N3A	-4.90	119.97	126.72

There are no chirality outliers.

5 of 39 torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
3	A	1001	COA	C5B-O5B-P1A-O1A
3	A	1001	COA	C5B-O5B-P1A-O2A
3	A	1001	COA	C5B-O5B-P1A-O3A
3	A	1001	COA	CDP-CBP-CCP-O6A
3	A	1001	COA	CEP-CBP-CCP-O6A


There are no ring outliers.


3 monomers are involved in 9 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
3	C	1003	COA	2	0
3	A	1001	COA	2	0
3	B	1002	COA	5	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier.

The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.





5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

The following chains have linkage breaks:

Mol	Chain	Number of breaks
1	A	1
1	D	1
1	C	1

All chain breaks are listed below:

Model	Chain	Residue-1	Atom-1	Residue-2	Atom-2	Distance (Å)
1	A	125:VAL	C	126:SCY	N	1.75
1	D	126:SCY	C	127:ALA	N	1.73
1	C	126:SCY	C	127:ALA	N	1.66

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	388/406 (95%)	0.53	15 (3%) 43 42	30, 48, 69, 80	0
1	B	385/406 (94%)	-0.13	0 100 100	21, 34, 49, 55	0
1	C	387/406 (95%)	0.08	7 (1%) 67 67	23, 38, 60, 71	0
1	D	381/406 (93%)	0.06	4 (1%) 79 79	30, 41, 58, 67	0
All	All	1541/1624 (94%)	0.14	26 (1%) 69 68	21, 41, 63, 80	0

The worst 5 of 26 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	32	SER	4.2
1	A	242	VAL	3.7
1	C	261	PHE	3.5
1	A	301	ALA	3.0
1	C	265	PRO	3.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled ‘Q < 0.9’ lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
1	SCY	A	126	9/10	0.83	0.15	43,46,53,54	0
1	SCY	C	126	9/10	0.87	0.12	31,33,48,56	0
1	SCY	B	126	9/10	0.92	0.11	25,28,42,51	0
1	SCY	D	126	9/10	0.95	0.08	32,34,48,50	0

6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.4 Ligands [i](#)




In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(\AA^2)	Q<0.9
3	COA	A	1001	48/48	0.81	0.14	71,98,105,106	0
3	COA	C	1003	48/48	0.84	0.12	69,81,93,93	0
3	COA	B	1002	48/48	0.85	0.12	44,62,71,73	0
3	COA	D	1004	48/48	0.86	0.11	57,67,78,79	0
2	CL	A	2003	1/1	0.97	0.06	42,42,42,42	0
2	CL	B	2001	1/1	0.99	0.03	29,29,29,29	0
2	CL	B	2002	1/1	0.99	0.03	34,34,34,34	0




The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.

Electron density around COA A 1001:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)




**Electron density around COA C 1003:**

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)






Electron density around COA B 1002:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)

**Electron density around COA D 1004:**

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)



6.5 Other polymers [i](#)

There are no such residues in this entry.