



Full wwPDB X-ray Structure Validation Report ⓘ

Mar 18, 2026 – 12:14 AM UTC

PDB ID : 6F3E / pdb_00006f3e
Title : IRAK4 IN COMPLEX WITH inhibitor
Authors : Xue, Y.; Degorce, S.L.; Robb, G.R.; Ferguson, A.D.
Deposited on : 2017-11-28
Resolution : 2.67 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4-5-2 with Phenix2.0
Mogul	:	2022.3.0, CSD as543be (2022)
Xtriage (Phenix)	:	2.0
EDS	:	3.0
Buster-report	:	wwPDB partial adaption of 1.1.7 (2018)
Percentile statistics	:	20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4	:	9.0.010 (Gargrove)
Density-Fitness	:	1.0.12
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.67 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	5070 (2.70-2.66)
Clashscore	190562	5409 (2.70-2.66)
Ramachandran outliers	187476	5324 (2.70-2.66)
Sidechain outliers	187428	5324 (2.70-2.66)
RSRZ outliers	180081	5070 (2.70-2.66)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	295	 5% 75% 16% • 8%
1	B	295	 6% 75% 19% • •

2 Entry composition [i](#)

There are 4 unique types of molecules in this entry. The entry contains 4474 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Interleukin-1 receptor-associated kinase 4.


Mol	Chain	Residues	Atoms						ZeroOcc	AltConf	Trace
1	A	272	Total	C	N	O	P	S	0	0	0
			2157	1361	363	417	2	14			
1	B	284	Total	C	N	O	P	S	0	0	0
			2245	1408	378	443	2	14			

- Molecule 2 is SULFATE ION (CCD ID: SO4) (formula: O₄S).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
2	A	1	Total	O	S	0	0
			5	4	1		

- Molecule 3 is 2-[(3 {R})-12-(4-morpholin-4-ylcyclohexyl)oxy-7-thia-9,11-diazatricyclo[6.4.0.0^{2,6}]dodeca-1(8),2(6),9,11-tetraen-3-yl]ethanamide (CCD ID: CJQ) (formula: C₂₁H₂₈N₄O₃S).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
3	A	1	Total	C	N	O	S	0	0
			29	21	4	3	1		
3	B	1	Total	C	N	O	S	0	0
			29	21	4	3	1		


- Molecule 4 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	A	5	Total	O	0	0
			5	5		
4	B	4	Total	O	0	0
			4	4		


3 Residue-property plots

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: Interleukin-1 receptor-associated kinase 4



- Molecule 1: Interleukin-1 receptor-associated kinase 4



4 Data and refinement statistics

Property	Value	Source
Space group	I 2 2 2	Depositor
Cell constants a, b, c, α , β , γ	87.18Å 109.37Å 141.54Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	68.17 – 2.67 68.17 – 2.67	Depositor EDS
% Data completeness (in resolution range)	99.5 (68.17-2.67) 99.8 (68.17-2.67)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	2.47 (at 2.69Å)	Xtriage
Refinement program	BUSTER 2.11.5 PACIOREK	Depositor
R, R_{free}	0.199 , 0.254 0.211 , 0.266	Depositor DCC
R_{free} test set	998 reflections (5.12%)	wwPDB-VP
Wilson B-factor (Å ²)	59.7	Xtriage
Anisotropy	0.508	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.36 , 81.5	EDS
L-test for twinning ²	$\langle L \rangle = 0.47$, $\langle L^2 \rangle = 0.30$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.95	EDS
Total number of atoms	4474	wwPDB-VP
Average B, all atoms (Å ²)	85.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The analyses of the Patterson function reveals a significant off-origin peak that is 22.38 % of the origin peak, indicating pseudo-translational symmetry. The chance of finding a peak of this or larger height randomly in a structure without pseudo-translational symmetry is equal to 5.7641e-03. The detected translational NCS is most likely also responsible for the elevated intensity ratio.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: TPO, SO4, CJQ

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.87	1/2170 (0.0%)	1.36	13/2920 (0.4%)
1	B	0.91	1/2259 (0.0%)	1.41	13/3042 (0.4%)
All	All	0.89	2/4429 (0.0%)	1.39	26/5962 (0.4%)

All (2) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	B	265	MET	SD-CE	-7.46	1.60	1.79
1	A	265	MET	SD-CE	-7.26	1.61	1.79

All (26) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	B	181	ASP	CA-CB-CG	8.13	120.73	112.60
1	B	172	GLU	CA-C-N	6.57	131.85	120.68
1	B	172	GLU	C-N-CA	6.57	131.85	120.68
1	B	193	GLY	CA-C-N	6.07	131.37	123.00
1	B	193	GLY	C-N-CA	6.07	131.37	123.00
1	A	272	ASP	CA-CB-CG	5.99	118.59	112.60
1	B	451	GLN	CA-C-N	5.97	128.60	120.54
1	B	451	GLN	C-N-CA	5.97	128.60	120.54
1	B	272	ASP	CA-CB-CG	5.84	118.44	112.60
1	A	451	GLN	CA-C-N	5.80	128.37	120.54
1	A	451	GLN	C-N-CA	5.80	128.37	120.54
1	A	320	ASP	CA-CB-CG	5.74	118.34	112.60
1	B	433	ALA	CA-C-N	5.44	127.88	120.54
1	B	433	ALA	C-N-CA	5.44	127.88	120.54
1	B	178	ASN	CA-CB-CG	5.41	118.01	112.60
1	A	242	HIS	CA-C-N	5.37	128.22	120.38
1	A	242	HIS	C-N-CA	5.37	128.22	120.38

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	B	320	ASP	CA-CB-CG	5.29	117.89	112.60
1	A	433	ALA	CA-C-N	5.23	127.59	120.54
1	A	433	ALA	C-N-CA	5.23	127.59	120.54
1	B	175	ASN	CA-CB-CG	5.20	117.80	112.60
1	A	412	ASP	CA-CB-CG	5.18	117.78	112.60
1	A	186	SER	CA-C-N	5.07	127.88	121.85
1	A	186	SER	C-N-CA	5.07	127.88	121.85
1	A	320	ASP	CA-C-N	5.03	127.52	120.28
1	A	320	ASP	C-N-CA	5.03	127.52	120.28

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2157	0	2142	18	0
1	B	2245	0	2212	31	0
2	A	5	0	0	0	0
3	A	29	0	0	0	0
3	B	29	0	0	0	0
4	A	5	0	0	0	0
4	B	4	0	0	0	0
All	All	4474	0	4354	47	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 5.

All (47) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:176:VAL:HG11	1:B:205:VAL:HG12	1.54	0.87
1:B:253:SER:HA	1:B:258:LEU:HD22	1.75	0.69
1:B:265:MET:HE2	1:B:326:LYS:HG3	1.76	0.68
1:B:181:ASP:HB3	1:B:190:ASN:HB2	1.76	0.67

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:265:MET:HE2	1:A:326:LYS:HG3	1.78	0.65
1:A:182:GLU:HA	1:A:191:LYS:HB2	1.81	0.62
1:B:265:MET:CE	1:B:326:LYS:HG3	2.31	0.61
1:A:265:MET:CE	1:A:326:LYS:HG3	2.31	0.60
1:B:246:VAL:HG11	1:B:318:LEU:HD12	1.87	0.56
1:B:172:GLU:O	1:B:176:VAL:HG13	2.06	0.56
1:B:287:MET:HE2	1:B:323:PHE:HB3	1.89	0.55
1:B:288:ARG:HB3	1:B:380:ILE:HG23	1.89	0.55
1:B:273:ARG:HG3	1:B:319:LEU:HD12	1.88	0.54
1:A:192:MET:HE3	1:A:200:VAL:HG12	1.90	0.54
1:A:246:VAL:HG11	1:A:318:LEU:HD12	1.89	0.54
1:B:173:LEU:HA	1:B:176:VAL:HG22	1.90	0.54
1:B:321:GLU:H	1:B:321:GLU:CD	2.16	0.54
1:A:288:ARG:HB3	1:A:380:ILE:HG23	1.91	0.52
1:A:273:ARG:HG3	1:A:319:LEU:HD12	1.91	0.52
1:B:251:PHE:HB3	1:B:260:LEU:HD23	1.94	0.50
1:B:169:SER:HB3	1:B:172:GLU:HG3	1.93	0.50
1:B:310:ARG:HD3	1:B:364:ILE:HG23	1.95	0.49
1:A:310:ARG:HD3	1:A:364:ILE:HG23	1.95	0.49
1:B:205:VAL:O	1:B:205:VAL:HG23	2.14	0.48
1:B:194:GLU:H	1:B:194:GLU:CD	2.23	0.47
1:A:284:SER:H	1:A:287:MET:HE3	1.79	0.47
1:A:237:MET:HE1	1:A:246:VAL:HG23	1.98	0.45
1:A:165:PHE:HB3	1:A:250:GLY:HA2	1.97	0.45
1:B:170:PHE:CE1	1:B:259:CYS:SG	3.10	0.44
1:B:415:ASP:HB3	1:B:418:MET:HE2	1.99	0.44
1:B:237:MET:HE1	1:B:246:VAL:HG23	1.98	0.44
1:A:415:ASP:HB3	1:A:418:MET:HE2	2.00	0.44
1:A:265:MET:HE1	1:A:320:ASP:HB3	2.00	0.43
1:B:222:THR:HG22	1:B:223:THR:H	1.82	0.43
1:A:240:CYS:HB3	1:A:301:PHE:HE2	1.83	0.43
1:B:252:SER:HB3	1:B:259:CYS:HB2	2.01	0.43
1:B:287:MET:CE	1:B:323:PHE:HB3	2.48	0.42
1:A:282:PRO:HG2	1:B:279:GLY:HA2	2.02	0.41
1:A:345:TPO:HB	1:A:364:ILE:HD11	2.03	0.41
1:B:192:MET:HE2	1:B:202:LYS:N	2.36	0.41
1:B:192:MET:SD	1:B:264:TYR:HE1	2.43	0.41
1:A:181:ASP:HB3	1:A:190:ASN:HB2	2.01	0.41
1:A:279:GLY:HA2	1:B:282:PRO:HG2	2.03	0.41
1:B:176:VAL:HG23	1:B:177:THR:HG23	2.03	0.41
1:B:394:GLN:HE21	1:B:394:GLN:HB3	1.71	0.40

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:164:ARG:HD2	1:B:165:PHE:H	1.85	0.40
1:B:287:MET:HE2	1:B:323:PHE:CB	2.51	0.40

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	263/295 (89%)	247 (94%)	16 (6%)	0	100	100
1	B	277/295 (94%)	260 (94%)	14 (5%)	3 (1%)	11	26
All	All	540/590 (92%)	507 (94%)	30 (6%)	3 (1%)	21	41

All (3) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	B	181	ASP
1	B	254	ASP
1	B	196	GLY

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	237/255 (93%)	223 (94%)	14 (6%)	18	38

Continued on next page...

Continued from previous page...

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	B	247/255 (97%)	232 (94%)	15 (6%)	17	37
All	All	484/510 (95%)	455 (94%)	29 (6%)	17	38

All (29) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	171	TYR
1	A	185	ILE
1	A	194	GLU
1	A	226	LEU
1	A	228	GLN
1	A	243	GLU
1	A	277	LEU
1	A	329	ASP
1	A	349	VAL
1	A	365	THR
1	A	394	GLN
1	A	404	GLU
1	A	427	GLU
1	A	455	GLN
1	B	164	ARG
1	B	174	LYS
1	B	194	GLU
1	B	199	VAL
1	B	226	LEU
1	B	254	ASP
1	B	256	ASP
1	B	277	LEU
1	B	329	ASP
1	B	334	ARG
1	B	349	VAL
1	B	394	GLN
1	B	404	GLU
1	B	427	GLU
1	B	455	GLN

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (13) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	229	GLN
1	A	306	HIS

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	A	390	HIS
1	A	394	GLN
1	A	438	HIS
1	A	442	ASN
1	A	451	GLN
1	B	229	GLN
1	B	241	GLN
1	B	297	ASN
1	B	394	GLN
1	B	442	ASN
1	B	451	GLN

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

4 non-standard protein/DNA/RNA residues are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
1	TPO	B	345	1	8,10,11	1.28	1 (12%)	10,14,16	1.23	1 (10%)
1	TPO	A	345	1	8,10,11	1.33	1 (12%)	10,14,16	1.28	2 (20%)
1	TPO	A	342	1	8,10,11	1.30	1 (12%)	10,14,16	1.39	2 (20%)
1	TPO	B	342	1	8,10,11	1.12	0	10,14,16	1.58	1 (10%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
1	TPO	B	345	1	-	2/9/11/13	-
1	TPO	A	345	1	-	2/9/11/13	-
1	TPO	A	342	1	-	2/9/11/13	-
1	TPO	B	342	1	-	4/9/11/13	-

All (3) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	342	TPO	P-OG1	-2.54	1.55	1.59
1	A	345	TPO	CB-CA	2.51	1.59	1.53
1	B	345	TPO	CB-CA	2.12	1.58	1.53

All (6) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	B	342	TPO	P-OG1-CB	-3.22	114.57	123.33
1	A	342	TPO	P-OG1-CB	-2.77	115.79	123.33
1	A	345	TPO	O2P-P-OG1	2.56	115.81	105.85
1	B	345	TPO	O2P-P-OG1	2.55	115.79	105.85
1	A	342	TPO	CG2-CB-CA	2.07	117.29	113.26
1	A	345	TPO	CG2-CB-CA	2.06	117.27	113.26

There are no chirality outliers.

All (10) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
1	A	345	TPO	N-CA-CB-OG1
1	A	345	TPO	O-C-CA-CB
1	B	342	TPO	N-CA-CB-OG1
1	B	342	TPO	C-CA-CB-CG2
1	B	342	TPO	O-C-CA-CB
1	B	345	TPO	N-CA-CB-OG1
1	B	345	TPO	O-C-CA-CB
1	A	342	TPO	C-CA-CB-CG2
1	A	342	TPO	O-C-CA-CB
1	B	342	TPO	N-CA-CB-CG2

There are no ring outliers.

1 monomer is involved in 1 short contact:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
1	A	345	TPO	1	0

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

3 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
3	CJQ	A	502	-	32,33,33	0.54	0	35,47,47	1.07	3 (8%)
2	SO4	A	501	-	4,4,4	0.33	0	6,6,6	0.28	0
3	CJQ	B	501	-	32,33,33	0.45	0	35,47,47	1.13	3 (8%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	CJQ	A	502	-	-	6/12/39/39	0/5/5/5
3	CJQ	B	501	-	-	5/12/39/39	0/5/5/5

There are no bond length outliers.

All (6) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	B	501	CJQ	O1-C11-C16	-3.53	100.08	108.31
3	B	501	CJQ	C20-N3-C14	3.44	121.53	112.79
3	A	502	CJQ	C5-S-C4	-2.77	89.60	90.87
3	A	502	CJQ	C20-N3-C14	2.64	119.50	112.79

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	B	501	CJQ	C5-S-C4	-2.38	89.78	90.87
3	A	502	CJQ	O1-C11-C12	-2.08	103.47	108.31

There are no chirality outliers.

All (11) torsion outliers are listed below:


Mol	Chain	Res	Type	Atoms
3	A	502	CJQ	C13-C14-N3-C20
3	A	502	CJQ	C13-C14-N3-C17
3	A	502	CJQ	C15-C14-N3-C20
3	A	502	CJQ	C15-C14-N3-C17
3	B	501	CJQ	C13-C14-N3-C17
3	B	501	CJQ	C13-C14-N3-C20
3	B	501	CJQ	C15-C14-N3-C17
3	B	501	CJQ	C15-C14-N3-C20
3	A	502	CJQ	O-C10-C9-C8
3	A	502	CJQ	N2-C10-C9-C8
3	B	501	CJQ	C7-C8-C9-C10

There are no ring outliers.


No monomer is involved in short contacts.

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.

Ligand CJQ A 502



Ligand CJQ B 501



5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	270/295 (91%)	0.27	16 (5%) 28 24	42, 75, 158, 175	0
1	B	282/295 (95%)	0.33	19 (6%) 24 20	42, 76, 125, 171	0
All	All	552/590 (93%)	0.30	35 (6%) 26 22	42, 75, 154, 175	0

All (35) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	B	259	CYS	3.7
1	B	336	SER	3.6
1	A	333	ALA	3.2
1	B	343	VAL	3.2
1	A	252	SER	3.1
1	B	169	SER	3.1
1	B	170	PHE	3.1
1	A	167	SER	2.9
1	A	186	SER	2.9
1	B	171	TYR	2.8
1	B	176	VAL	2.7
1	B	187	VAL	2.7
1	B	195	GLY	2.6
1	A	196	GLY	2.6
1	B	168	PHE	2.6
1	A	171	TYR	2.6
1	A	175	ASN	2.5
1	A	197	PHE	2.5
1	B	251	PHE	2.5
1	B	226	LEU	2.4
1	A	230	PHE	2.4
1	A	238	ALA	2.4
1	B	201	TYR	2.4
1	B	344	MET	2.3

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	A	258	LEU	2.3
1	B	346	SER	2.3
1	A	165	PHE	2.3
1	A	343	VAL	2.2
1	B	179	ASN	2.1
1	A	350	GLY	2.1
1	B	222	THR	2.1
1	B	258	LEU	2.0
1	A	207	ASN	2.0
1	B	255	GLY	2.0
1	A	179	ASN	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
1	TPO	A	342	11/12	0.60	0.17	151,153,157,159	0
1	TPO	B	342	11/12	0.61	0.17	167,175,183,187	0
1	TPO	B	345	11/12	0.62	0.16	159,165,171,174	0
1	TPO	A	345	11/12	0.80	0.13	142,149,153,157	0

6.3 Carbohydrates [i](#)


There are no oligosaccharides in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.




Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
2	SO4	A	501	5/5	0.85	0.17	90,92,93,95	0
3	CJQ	A	502	29/29	0.94	0.09	50,61,66,67	0
3	CJQ	B	501	29/29	0.96	0.08	43,47,55,56	0

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.



Electron density around CJQ B 501:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)



6.5 Other polymers [i](#)

There are no such residues in this entry.