



wwPDB X-ray Structure Validation Summary Report ⓘ

Mar 20, 2026 – 09:51 AM UTC

PDB ID : 3F79 / pdb_00003f79
Title : Structure of pseudo-centered cell crystal form of the C-terminal phosphatase domain of *P. aeruginosa* RssB
Authors : Levchenko, I.; Grant, R.A.; Sauer, R.T.; Baker, T.A.
Deposited on : 2008-11-07
Resolution : 2.80 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4-5-2 with Phenix2.0
Mogul : 2022.3.0, CSD as543be (2022)
Xtriage (Phenix) : 2.0
EDS : 3.0
Percentile statistics : 20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4 : 9.0.010 (Gargrove)
Density-Fitness : 1.0.12
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.80 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	3866 (2.80-2.80)
Clashscore	190562	4276 (2.80-2.80)
Ramachandran outliers	187476	4196 (2.80-2.80)
Sidechain outliers	187428	4198 (2.80-2.80)
RSRZ outliers	180081	3869 (2.80-2.80)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	255	41% 45% 13%
1	B	255	43% 45% 10%
1	C	255	37% 50% 10%
1	D	255	39% 49% 10%
1	E	255	51% 40% 7%

Continued on next page...

Continued from previous page...

Mol	Chain	Length	Quality of chain
1	F	255	 A horizontal bar chart showing the quality of the chain. The bar is divided into three segments: a green segment representing 53%, a yellow segment representing 38%, and a small grey segment representing 8%.

2 Entry composition

There are 2 unique types of molecules in this entry. The entry contains 10386 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Probable two-component response regulator.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	223	Total	C	N	O	Se	12	0	0
			1659	1065	276	312	6			
1	B	229	Total	C	N	O	Se	5	0	0
			1738	1111	295	326	6			
1	C	229	Total	C	N	O	Se	5	0	0
			1728	1112	286	325	5			
1	D	230	Total	C	N	O	Se	0	0	0
			1719	1105	284	324	6			
1	E	236	Total	C	N	O	Se	13	0	0
			1778	1133	302	337	6			
1	F	235	Total	C	N	O	Se	12	0	0
			1757	1125	300	327	5			


- Molecule 2 is MAGNESIUM ION (CCD ID: MG) (formula: Mg).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
2	B	1	Total	Mg	0	0
			1	1		
2	C	1	Total	Mg	0	0
			1	1		
2	D	2	Total	Mg	0	0
			2	2		
2	E	1	Total	Mg	0	0
			1	1		
2	F	2	Total	Mg	0	0
			2	2		


3 Residue-property plots

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: Probable two-component response regulator




- Molecule 1: Probable two-component response regulator




- Molecule 1: Probable two-component response regulator






- Molecule 1: Probable two-component response regulator

Chain D: 39% 49% 10%




- Molecule 1: Probable two-component response regulator

Chain E: 51% 40% 7%



- Molecule 1: Probable two-component response regulator

Chain F: 53% 38% 8%



R306	GLY	P309	V310	G311	L312	L325	S328	F329	S330	L331	S332	L333	F334	S335	D336	G337	I338	L339	D340	V341	L342	L354	P355	D366	Q370	V371	F372	GLY	LEU	ALA	ASN	LEU	ALA	GLU	MSF	P381	I384	A385	L386	L389	S390	R391	N392	L393	ALA
------	-----	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	-----	-----	-----	-----	-----	-----	-----	-----	------	------	------	------	------	------	------	------	------	-----

4 Data and refinement statistics

Property	Value	Source
Space group	P 31	Depositor
Cell constants a, b, c, α , β , γ	124.87Å 124.87Å 92.90Å 90.00° 90.00° 120.00°	Depositor
Resolution (Å)	15.00 – 2.80 15.00 – 2.80	Depositor EDS
% Data completeness (in resolution range)	98.4 (15.00-2.80) 99.5 (15.00-2.80)	Depositor EDS
R_{merge}	0.08	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	2.85 (at 2.81Å)	Xtriage
Refinement program	PHENIX	Depositor
R, R_{free}	0.243 , 0.297 0.243 , 0.299	Depositor DCC
R_{free} test set	1983 reflections (4.98%)	wwPDB-VP
Wilson B-factor (Å ²)	56.8	Xtriage
Anisotropy	0.597	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.27 , 73.1	EDS
L-test for twinning ²	$\langle L \rangle = 0.46$, $\langle L^2 \rangle = 0.28$	Xtriage
Estimated twinning fraction	0.047 for -h,-k,l 0.048 for h,-h-k,-l 0.477 for -k,-h,-l	Xtriage
Reported twinning fraction	0.498 for -k,-h,-l	Depositor
Outliers	0 of 39804 reflections	Xtriage
F_o, F_c correlation	0.97	EDS
Total number of atoms	10386	wwPDB-VP
Average B, all atoms (Å ²)	91.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The analyses of the Patterson function reveals a significant off-origin peak that is 48.93 % of the origin peak, indicating pseudo-translational symmetry. The chance of finding a peak of this or larger height randomly in a structure without pseudo-translational symmetry is equal to 8.0149e-05. The detected translational NCS is most likely also responsible for the elevated intensity ratio.*

¹ Intensities estimated from amplitudes.

² Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: MG

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.27	0/1682	0.73	1/2271 (0.0%)
1	B	0.29	0/1762	0.73	0/2371
1	C	0.28	0/1756	0.77	1/2374 (0.0%)
1	D	0.31	0/1740	0.74	1/2349 (0.0%)
1	E	0.28	0/1800	0.73	0/2429
1	F	0.29	0/1785	0.73	0/2413
All	All	0.29	0/10525	0.74	3/14207 (0.0%)

There are no bond length outliers.

All (3) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	379	GLU	CB-CA-C	-6.44	109.16	116.63
1	C	279	ASN	N-CA-C	-5.58	106.42	113.23
1	D	245	PHE	N-CA-C	-5.43	105.36	111.28

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1659	0	1613	130	0
1	B	1738	0	1706	167	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	C	1728	0	1687	199	0
1	D	1719	0	1680	192	0
1	E	1778	0	1758	123	0
1	F	1757	0	1713	124	0
2	B	1	0	0	0	0
2	C	1	0	0	0	0
2	D	2	0	0	0	0
2	E	1	0	0	0	0
2	F	2	0	0	0	0
All	All	10386	0	10157	860	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 42.

The worst 5 of 860 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:D:246:LYS:CB	1:D:249:GLU:HB3	1.35	1.51
1:D:246:LYS:HB2	1:D:249:GLU:CB	1.39	1.49
1:D:380:MSE:SE	1:D:381:PRO:HD2	1.85	1.26
1:B:356:GLU:HG3	1:B:357:GLN:NE2	1.56	1.20
1:D:175:GLU:O	1:D:391:ARG:NH2	1.73	1.19

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	209/255 (82%)	193 (92%)	16 (8%)	0	100	100
1	B	217/255 (85%)	202 (93%)	14 (6%)	1 (0%)	24	55

Continued on next page...

Continued from previous page...

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	C	219/255 (86%)	199 (91%)	20 (9%)	0	100	100
1	D	218/255 (86%)	202 (93%)	15 (7%)	1 (0%)	24	55
1	E	226/255 (89%)	211 (93%)	15 (7%)	0	100	100
1	F	227/255 (89%)	214 (94%)	13 (6%)	0	100	100
All	All	1316/1530 (86%)	1221 (93%)	93 (7%)	2 (0%)	43	72

All (2) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	B	310	VAL
1	D	337	GLY

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	173/206 (84%)	172 (99%)	1 (1%)	78	92
1	B	182/206 (88%)	178 (98%)	4 (2%)	45	78
1	C	181/206 (88%)	174 (96%)	7 (4%)	28	64
1	D	179/206 (87%)	172 (96%)	7 (4%)	28	64
1	E	189/206 (92%)	187 (99%)	2 (1%)	65	87
1	F	181/206 (88%)	179 (99%)	2 (1%)	65	87
All	All	1085/1236 (88%)	1062 (98%)	23 (2%)	47	79

5 of 23 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	D	289	LEU
1	D	366	ASP
1	D	348	LYS
1	D	376	ASN
1	C	162	VAL

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 19 such sidechains are listed below:

Mol	Chain	Res	Type
1	E	214	HIS
1	E	370	GLN
1	F	150	ASN
1	E	255	ASN
1	C	266	HIS

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

Of 7 ligands modelled in this entry, 7 are monoatomic - leaving 0 for Mogul analysis.

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues ⓘ

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	215/255 (84%)	-1.14	0 100 100	41, 92, 166, 244	0
1	B	222/255 (87%)	-1.15	0 100 100	42, 86, 129, 224	0
1	C	223/255 (87%)	-1.15	0 100 100	43, 91, 141, 234	0
1	D	224/255 (87%)	-1.17	0 100 100	36, 88, 150, 275	0
1	E	228/255 (89%)	-1.21	0 100 100	39, 86, 146, 192	0
1	F	228/255 (89%)	-1.28	0 100 100	23, 80, 132, 226	0
All	All	1340/1530 (87%)	-1.18	0 100 100	23, 88, 146, 275	0

There are no RSRZ outliers to report.

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled ‘Q< 0.9’ lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
2	MG	D	6	1/1	0.98	0.08	78,78,78,78	0

Continued on next page...

Continued from previous page...

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors($\text{\AA}^2$)	Q<0.9
2	MG	C	5	1/1	0.99	0.02	50,50,50,50	0
2	MG	B	1	1/1	0.99	0.02	61,61,61,61	0
2	MG	F	3	1/1	0.99	0.03	47,47,47,47	0
2	MG	F	4	1/1	0.99	0.06	94,94,94,94	0
2	MG	D	7	1/1	1.00	0.02	40,40,40,40	0
2	MG	E	2	1/1	1.00	0.02	39,39,39,39	0

6.5 Other polymers [i](#)

There are no such residues in this entry.