



## wwPDB EM Validation Summary Report ⓘ

Mar 24, 2026 – 02:00 PM UTC

PDB ID : 7FDC / pdb\_00007fdc  
EMDB ID : EMD-31540  
Title : CryoEM Structures of Reconstituted V-ATPase, state3  
Authors : Khan, M.M.; Lee, S.; Oot, R.A.; Couch-Cardel, S.; KIm, H.; Wilkens, S.; Roh, S.H.  
Deposited on : 2021-07-16  
Resolution : 6.60 Å(reported)

This is a wwPDB EM Validation Summary Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<https://www.wwpdb.org/validation/2017/EMValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:



EMDB validation analysis : 0.0.1.dev132  
MolProbity : 4-5-2 with Phenix2.0  
Percentile statistics : 20250101.v01 (using entries in the PDB archive January 1st 2025)  
EM percentile statistics : 202505.v01 (Using data in the EMDB archive up until May 2025)  
MapQ : 1.9.13  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : 2.49

# 1 Overall quality at a glance i

The following experimental techniques were used to determine the structure:  
*ELECTRON MICROSCOPY*

The reported resolution of this entry is 6.60 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



| Metric                | Whole archive<br>(#Entries) | EM structures<br>(#Entries) | Similar EM resolution<br>(#Entries, resolution range(Å)) |
|-----------------------|-----------------------------|-----------------------------|--|
| Clashscore            | 229148                      | 23984                       | -  |
| Ramachandran outliers | 224038                      | 23583                       | -  |
| Sidechain outliers    | 223484                      | 23102                       | -  |
| Q-score               | -                           | 25397                       | 531 ( 6.10 - 7.10 )                                      |

The table below summarises the geometric issues observed across the polymeric chains and their fit to the map. The red, orange, yellow and green segments of the bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the EM map (all-atom inclusion  $< 40\%$ ). The numeric value is given above the bar.

| Mol | Chain | Length | Quality of chain   |
|-----|-------|--------|--|
| 1   | A     | 617    | <p>39% (Poor fit), 67% (0 outliers), 29% (1 outlier), 3% (2 outliers), 0% (3+ outliers), 0% (Not modelled)</p> |
| 1   | C     | 617    | <p>22% (Poor fit), 64% (0 outliers), 33% (1 outlier), 3% (2 outliers), 0% (3+ outliers), 0% (Not modelled)</p> |
| 1   | E     | 617    | <p>43% (Poor fit), 68% (0 outliers), 28% (1 outlier), 3% (2 outliers), 0% (3+ outliers), 0% (Not modelled)</p> |
| 2   | B     | 517    | <p>30% (Poor fit), 65% (0 outliers), 26% (1 outlier), 9% (2 outliers), 0% (3+ outliers), 0% (Not modelled)</p> |



Continued on next page...

Continued from previous page...

| Mol | Chain | Length | Quality of chain |
|-----|-------|--------|------------------|
| 2   | D     | 517    |                  |
| 2   | F     | 517    |                  |
| 3   | G     | 233    |                  |
| 3   | I     | 233    |                  |
| 3   | K     | 233    |                  |
| 4   | H     | 122    |                  |
| 4   | J     | 122    |                  |
| 4   | L     | 122    |                  |
| 5   | M     | 256    |                  |
| 6   | N     | 118    |                  |
| 7   | O     | 392    |                  |
| 8   | P     | 469    |                  |
| 9   | Q     | 840    |                  |
| 10  | S     | 345    |                  |
| 11  | T     | 213    |                  |
| 12  | U     | 164    |                  |
| 13  | V     | 160    |                  |
| 13  | W     | 160    |                  |
| 13  | X     | 160    |                  |
| 13  | Y     | 160    |                  |
| 13  | Z     | 160    |                  |
| 13  | a     | 160    |                  |
| 13  | b     | 160    |                  |
| 13  | c     | 160    |                  |
| 14  | d     | 73     |                  |

Continued on next page...

*Continued from previous page...*

| Mol | Chain | Length | Quality of chain   |
|-----|-------|--------|--|
| 15  | e     | 265    |  |
| 16  | f     | 85     |  |

## 2 Entry composition [i](#)

There are 16 unique types of molecules in this entry. The entry contains 64497 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Yeast Vacuolar ATPase A subunit.

| Mol | Chain | Residues | Atoms |      |     |     |    | AltConf | Trace |
|-----|-------|----------|-------|------|-----|-----|----|---------|-------|
|     |       |          | Total | C    | N   | O   | S  |         |       |
| 1   | A     | 594      | Total | C    | N   | O   | S  | 0       | 0     |
|     |       |          | 4586  | 2909 | 761 | 896 | 20 |         |       |
| 1   | C     | 594      | Total | C    | N   | O   | S  | 0       | 0     |
|     |       |          | 4586  | 2909 | 761 | 896 | 20 |         |       |
| 1   | E     | 594      | Total | C    | N   | O   | S  | 0       | 0     |
|     |       |          | 4586  | 2909 | 761 | 896 | 20 |         |       |

- Molecule 2 is a protein called V-type proton ATPase subunit B.

| Mol | Chain | Residues | Atoms |      |     |     |    | AltConf | Trace |
|-----|-------|----------|-------|------|-----|-----|----|---------|-------|
|     |       |          | Total | C    | N   | O   | S  |         |       |
| 2   | B     | 468      | Total | C    | N   | O   | S  | 0       | 0     |
|     |       |          | 3681  | 2332 | 631 | 706 | 12 |         |       |
| 2   | D     | 470      | Total | C    | N   | O   | S  | 0       | 0     |
|     |       |          | 3699  | 2342 | 633 | 712 | 12 |         |       |
| 2   | F     | 471      | Total | C    | N   | O   | S  | 0       | 0     |
|     |       |          | 3706  | 2347 | 634 | 713 | 12 |         |       |

- Molecule 3 is a protein called V-type proton ATPase subunit E.

| Mol | Chain | Residues | Atoms |      |     |     |   | AltConf | Trace |
|-----|-------|----------|-------|------|-----|-----|---|---------|-------|
|     |       |          | Total | C    | N   | O   | S |         |       |
| 3   | G     | 225      | Total | C    | N   | O   | S | 0       | 0     |
|     |       |          | 1802  | 1131 | 309 | 357 | 5 |         |       |
| 3   | I     | 225      | Total | C    | N   | O   | S | 0       | 0     |
|     |       |          | 1802  | 1131 | 309 | 357 | 5 |         |       |
| 3   | K     | 225      | Total | C    | N   | O   | S | 0       | 0     |
|     |       |          | 1802  | 1131 | 309 | 357 | 5 |         |       |

- Molecule 4 is a protein called V-type proton ATPase subunit G.

| Mol | Chain | Residues | Atoms |     |     |     | AltConf | Trace |
|-----|-------|----------|-------|-----|-----|-----|---------|-------|
|     |       |          | Total | C   | N   | O   |         |       |
| 4   | H     | 111      | Total | C   | N   | O   | 0       | 0     |
|     |       |          | 871   | 546 | 153 | 172 |         |       |

*Continued on next page...*

*Continued from previous page...*

| Mol | Chain | Residues | Atoms |     |     |     | AltConf | Trace |
|-----|-------|----------|-------|-----|-----|-----|---------|-------|
| 4   | J     | 111      | Total | C   | N   | O   | 0       | 0     |
|     |       |          | 871   | 546 | 153 | 172 |         |       |
| 4   | L     | 111      | Total | C   | N   | O   | 0       | 0     |
|     |       |          | 871   | 546 | 153 | 172 |         |       |

- Molecule 5 is a protein called V-type proton ATPase subunit D.

| Mol | Chain | Residues | Atoms |      |     |     |   | AltConf | Trace |
|-----|-------|----------|-------|------|-----|-----|---|---------|-------|
| 5   | M     | 218      | Total | C    | N   | O   | S | 0       | 0     |
|     |       |          | 1756  | 1100 | 315 | 336 | 5 |         |       |

- Molecule 6 is a protein called V-type proton ATPase subunit F.

| Mol | Chain | Residues | Atoms |     |     |     | AltConf | Trace |
|-----|-------|----------|-------|-----|-----|-----|---------|-------|
| 6   | N     | 115      | Total | C   | N   | O   | 0       | 0     |
|     |       |          | 928   | 589 | 157 | 182 |         |       |

- Molecule 7 is a protein called V-type proton ATPase subunit C.

| Mol | Chain | Residues | Atoms |      |     |     |   | AltConf | Trace |
|-----|-------|----------|-------|------|-----|-----|---|---------|-------|
| 7   | O     | 392      | Total | C    | N   | O   | S | 0       | 0     |
|     |       |          | 3121  | 2005 | 516 | 595 | 5 |         |       |

- Molecule 8 is a protein called Fusion of yeast V-type proton ATPase subunit H(NT) and human V-type proton ATPase subunit H(CT).

| Mol | Chain | Residues | Atoms |      |     |     |    | AltConf | Trace |
|-----|-------|----------|-------|------|-----|-----|----|---------|-------|
| 8   | P     | 452      | Total | C    | N   | O   | S  | 0       | 0     |
|     |       |          | 3665  | 2338 | 632 | 682 | 13 |         |       |

- Molecule 9 is a protein called Yeast Vacuolar ATPase a subunit.

| Mol | Chain | Residues | Atoms |      |     |      |    | AltConf | Trace |
|-----|-------|----------|-------|------|-----|------|----|---------|-------|
| 9   | Q     | 747      | Total | C    | N   | O    | S  | 0       | 0     |
|     |       |          | 6069  | 3960 | 986 | 1088 | 35 |         |       |

- Molecule 10 is a protein called V-type proton ATPase subunit d.

| Mol | Chain | Residues | Atoms |      |     |     |    | AltConf | Trace |
|-----|-------|----------|-------|------|-----|-----|----|---------|-------|
| 10  | S     | 344      | Total | C    | N   | O   | S  | 0       | 0     |
|     |       |          | 2793  | 1774 | 453 | 553 | 13 |         |       |

- Molecule 11 is a protein called V-type proton ATPase subunit c''.

| Mol | Chain | Residues | Atoms |     |     |     |   | AltConf | Trace |
|-----|-------|----------|-------|-----|-----|-----|---|---------|-------|
|     |       |          | Total | C   | N   | O   | S |         |       |
| 11  | T     | 200      | 1492  | 995 | 231 | 259 | 7 | 0       | 0     |

- Molecule 12 is a protein called V-type proton ATPase subunit c'.

| Mol | Chain | Residues | Atoms |     |     |     |    | AltConf | Trace |
|-----|-------|----------|-------|-----|-----|-----|----|---------|-------|
|     |       |          | Total | C   | N   | O   | S  |         |       |
| 12  | U     | 157      | 1139  | 753 | 179 | 195 | 12 | 0       | 0     |

- Molecule 13 is a protein called V-type proton ATPase subunit c.

| Mol | Chain | Residues | Atoms |     |     |     |   | AltConf | Trace |
|-----|-------|----------|-------|-----|-----|-----|---|---------|-------|
|     |       |          | Total | C   | N   | O   | S |         |       |
| 13  | V     | 159      | 1140  | 751 | 182 | 199 | 8 | 0       | 0     |
| 13  | W     | 159      | 1140  | 751 | 182 | 199 | 8 | 0       | 0     |
| 13  | X     | 159      | 1140  | 751 | 182 | 199 | 8 | 0       | 0     |
| 13  | Y     | 160      | 1146  | 754 | 183 | 200 | 9 | 0       | 0     |
| 13  | Z     | 159      | 1140  | 751 | 182 | 199 | 8 | 0       | 0     |
| 13  | a     | 159      | 1140  | 751 | 182 | 199 | 8 | 0       | 0     |
| 13  | b     | 160      | 1146  | 754 | 183 | 200 | 9 | 0       | 0     |
| 13  | c     | 159      | 1140  | 751 | 182 | 199 | 8 | 0       | 0     |

- Molecule 14 is a protein called V-type proton ATPase subunit e.

| Mol | Chain | Residues | Atoms |     |    |    |   | AltConf | Trace |
|-----|-------|----------|-------|-----|----|----|---|---------|-------|
|     |       |          | Total | C   | N  | O  | S |         |       |
| 14  | d     | 69       | 553   | 369 | 91 | 86 | 7 | 0       | 0     |

- Molecule 15 is a protein called V0 assembly protein 1.

| Mol | Chain | Residues | Atoms |     |    |    |   | AltConf | Trace |
|-----|-------|----------|-------|-----|----|----|---|---------|-------|
|     |       |          | Total | C   | N  | O  | S |         |       |
| 15  | e     | 52       | 403   | 268 | 59 | 74 | 2 | 0       | 0     |


- Molecule 16 is a protein called Yeast Vacuolar ATPase f subunit.

| Mol | Chain | Residues | Atoms |     |    |     |   | AltConf | Trace |
|-----|-------|----------|-------|-----|----|-----|---|---------|-------|
|     |       |          | Total | C   | N  | O   | S |         |       |
| 16  | f     | 76       | 583   | 386 | 94 | 100 | 3 | 0       | 0     |

### 3 Residue-property plots


These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and atom inclusion in map density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red diamond above a residue indicates a poor fit to the EM map for this residue (all-atom inclusion < 40%). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: Yeast Vacuolar ATPase A subunit




- Molecule 1: Yeast Vacuolar ATPase A subunit






• Molecule 2: V-type proton ATPase subunit B





• Molecule 2: V-type proton ATPase subunit B









• Molecule 4: V-type proton ATPase subunit G



• Molecule 4: V-type proton ATPase subunit G




• Molecule 4: V-type proton ATPase subunit G




• Molecule 5: V-type proton ATPase subunit D







• Molecule 6: V-type proton ATPase subunit F




• Molecule 7: V-type proton ATPase subunit C




- Molecule 8: Fusion of yeast V-type proton ATPase subunit H(NT) and human V-type proton ATPase subunit H(CT)



- Molecule 9: Yeast Vacuolar ATPase a subunit







• Molecule 10: V-type proton ATPase subunit d




Chain S:






• Molecule 11: V-type proton ATPase subunit c'




• Molecule 12: V-type proton ATPase subunit c'




• Molecule 13: V-type proton ATPase subunit c








• Molecule 13: V-type proton ATPase subunit c




• Molecule 13: V-type proton ATPase subunit c






• Molecule 13: V-type proton ATPase subunit c











• Molecule 13: V-type proton ATPase subunit c



• Molecule 13: V-type proton ATPase subunit c



• Molecule 13: V-type proton ATPase subunit c



• Molecule 13: V-type proton ATPase subunit c





## 4 Experimental information

| Property                             | Value                                   | Source    |
|--------------------------------------|---|-----------|
| EM reconstruction method             | SINGLE PARTICLE                         | Depositor |
| Imposed symmetry                     | POINT, Not provided                     |           |
| Number of particles used             | 15741                                   | Depositor |
| Resolution determination method      | FSC 0.143 CUT-OFF                       | Depositor |
| CTF correction method                | PHASE FLIPPING AND AMPLITUDE CORRECTION | Depositor |
| Microscope                           | TFS KRIOS                               | Depositor |
| Voltage (kV)                         | 300                                     | Depositor |
| Electron dose ( $e^-/\text{\AA}^2$ ) | 50                                      | Depositor |
| Minimum defocus (nm)                 | Not provided                            |           |
| Maximum defocus (nm)                 | Not provided                            |           |
| Magnification                        | Not provided                            |           |
| Image detector                       | GATAN K2 SUMMIT (4k x 4k)               | Depositor |
| Maximum map value                    | 0.066                                   | Depositor |
| Minimum map value                    | -0.024                                  | Depositor |
| Average map value                    | 0.000                                   | Depositor |
| Map value standard deviation         | 0.004                                   | Depositor |
| Recommended contour level            | 0.02                                    | Depositor |
| Map size ( $\text{\AA}$ )            | 432.00003, 432.00003, 432.00003         | wwPDB     |
| Map dimensions                       | 400, 400, 400                           | wwPDB     |
| Map angles ( $^\circ$ )              | 90.0, 90.0, 90.0                        | wwPDB     |
| Pixel spacing ( $\text{\AA}$ )       | 1.08, 1.08, 1.08                        | Depositor |

## 5 Model quality [i](#)

### 5.1 Standard geometry [i](#)

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

| Mol | Chain | Bond lengths |         | Bond angles |                |
|-----|-------|--------------|---------|-------------|----------------|
|     |       | RMSZ         | # Z  >5 | RMSZ        | # Z  >5        |
| 1   | A     | 0.17         | 0/4685  | 0.32        | 0/6351         |
| 1   | C     | 0.17         | 0/4685  | 0.34        | 0/6351         |
| 1   | E     | 0.16         | 0/4685  | 0.31        | 0/6351         |
| 2   | B     | 0.17         | 0/3750  | 0.33        | 0/5078         |
| 2   | D     | 0.18         | 0/3768  | 0.34        | 0/5102         |
| 2   | F     | 0.17         | 0/3776  | 0.33        | 0/5114         |
| 3   | G     | 0.16         | 0/1817  | 0.33        | 0/2436         |
| 3   | I     | 0.16         | 0/1817  | 0.32        | 0/2436         |
| 3   | K     | 0.17         | 0/1817  | 0.33        | 0/2436         |
| 4   | H     | 0.14         | 0/876   | 0.31        | 0/1164         |
| 4   | J     | 0.14         | 0/876   | 0.29        | 0/1164         |
| 4   | L     | 0.16         | 0/876   | 0.34        | 0/1164         |
| 5   | M     | 0.16         | 0/1775  | 0.33        | 0/2381         |
| 6   | N     | 0.15         | 0/944   | 0.32        | 0/1277         |
| 7   | O     | 0.18         | 0/3184  | 0.43        | 0/4314         |
| 8   | P     | 0.17         | 0/3723  | 0.39        | 0/5028         |
| 9   | Q     | 0.18         | 0/6221  | 0.40        | 0/8421         |
| 10  | S     | 0.13         | 0/2852  | 0.34        | 0/3870         |
| 11  | T     | 0.18         | 0/1523  | 0.37        | 0/2068         |
| 12  | U     | 0.16         | 0/1162  | 0.38        | 0/1575         |
| 13  | V     | 0.15         | 0/1158  | 0.36        | 0/1574         |
| 13  | W     | 0.16         | 0/1158  | 0.38        | 0/1574         |
| 13  | X     | 0.18         | 0/1158  | 0.37        | 0/1574         |
| 13  | Y     | 0.17         | 0/1164  | 0.41        | 1/1582 (0.1%)  |
| 13  | Z     | 0.17         | 0/1158  | 0.39        | 0/1574         |
| 13  | a     | 0.18         | 0/1158  | 0.41        | 0/1574         |
| 13  | b     | 0.19         | 0/1164  | 0.39        | 0/1582         |
| 13  | c     | 0.21         | 0/1158  | 0.43        | 0/1574         |
| 14  | d     | 0.16         | 0/569   | 0.41        | 0/776          |
| 15  | e     | 0.12         | 0/409   | 0.35        | 0/557          |
| 16  | f     | 0.12         | 0/600   | 0.34        | 0/822          |
| All | All   | 0.17         | 0/65666 | 0.36        | 1/88844 (0.0%) |

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if

the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

| Mol | Chain | #Chirality outliers | #Planarity outliers |
|-----|-------|---------------------|---------------------|
| 7   | O     | 0                   | 1                   |
| 8   | P     | 0                   | 1                   |
| All | All   | 0                   | 2                   |

There are no bond length outliers.

All (1) bond angle outliers are listed below:

| Mol | Chain | Res | Type | Atoms  | Z     | Observed(°) | Ideal(°) |
|-----|-------|-----|------|--------|-------|-------------|----------|
| 13  | Y     | 54  | ILE  | N-CA-C | -6.26 | 106.34      | 111.91   |

There are no chirality outliers.

All (2) planarity outliers are listed below:

| Mol | Chain | Res | Type | Group   |
|-----|-------|-----|------|---------|
| 7   | O     | 320 | GLY  | Peptide |
| 8   | P     | 354 | THR  | Peptide |

## 5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

| Mol | Chain | Non-H | H(model) | H(added) | Clashes | Symm-Clashes |
|-----|-------|-------|----------|----------|---------|--------------|
| 1   | A     | 4586  | 0        | 4525     | 126     | 0            |
| 1   | C     | 4586  | 0        | 4525     | 159     | 0            |
| 1   | E     | 4586  | 0        | 4525     | 109     | 0            |
| 2   | B     | 3681  | 0        | 3681     | 103     | 0            |
| 2   | D     | 3699  | 0        | 3693     | 116     | 0            |
| 2   | F     | 3706  | 0        | 3700     | 101     | 0            |
| 3   | G     | 1802  | 0        | 1873     | 50      | 0            |
| 3   | I     | 1802  | 0        | 1873     | 46      | 0            |
| 3   | K     | 1802  | 0        | 1873     | 64      | 0            |
| 4   | H     | 871   | 0        | 921      | 28      | 0            |
| 4   | J     | 871   | 0        | 921      | 29      | 0            |
| 4   | L     | 871   | 0        | 921      | 34      | 0            |

*Continued on next page...*

Continued from previous page...

| Mol | Chain | Non-H | H(model) | H(added) | Clashes | Symm-Clashes |
|-----|-------|-------|----------|----------|---------|--------------|
| 5   | M     | 1756  | 0        | 1802     | 57      | 0            |
| 6   | N     | 928   | 0        | 926      | 28      | 0            |
| 7   | O     | 3121  | 0        | 3155     | 164     | 0            |
| 8   | P     | 3665  | 0        | 3771     | 178     | 0            |
| 9   | Q     | 6069  | 0        | 6041     | 285     | 0            |
| 10  | S     | 2793  | 0        | 2677     | 92      | 0            |
| 11  | T     | 1492  | 0        | 1562     | 89      | 0            |
| 12  | U     | 1139  | 0        | 1194     | 73      | 0            |
| 13  | V     | 1140  | 0        | 1214     | 62      | 0            |
| 13  | W     | 1140  | 0        | 1214     | 47      | 0            |
| 13  | X     | 1140  | 0        | 1214     | 62      | 0            |
| 13  | Y     | 1146  | 0        | 1219     | 67      | 0            |
| 13  | Z     | 1140  | 0        | 1214     | 67      | 0            |
| 13  | a     | 1140  | 0        | 1214     | 57      | 0            |
| 13  | b     | 1146  | 0        | 1219     | 63      | 0            |
| 13  | c     | 1140  | 0        | 1214     | 98      | 0            |
| 14  | d     | 553   | 0        | 581      | 32      | 0            |
| 15  | e     | 403   | 0        | 432      | 11      | 0            |
| 16  | f     | 583   | 0        | 576      | 13      | 0            |
| All | All   | 64497 | 0        | 65470    | 2249    | 0            |

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 17.

The worst 5 of 2249 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

| Atom-1          | Atom-2            | Interatomic distance (Å) | Clash overlap (Å) |
|-----------------|-------------------|--------------------------|-------------------|
| 13:c:59:MET:HE3 | 13:c:133:LEU:HD21 | 1.31                     | 1.09              |
| 1:A:238:ARG:NH2 | 1:A:524:ASP:O     | 1.97                     | 0.97              |
| 9:Q:87:ASP:OD1  | 9:Q:90:LYS:NZ     | 2.01                     | 0.93              |
| 4:J:2:SER:OG    | 9:Q:150:GLU:OE2   | 1.88                     | 0.90              |
| 5:M:64:ALA:O    | 5:M:67:SER:OG     | 1.90                     | 0.90              |

There are no symmetry-related clashes.

## 5.3 Torsion angles

### 5.3.1 Protein backbone

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all PDB entries followed by that with respect to all EM entries.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

| Mol | Chain | Analysed       | Favoured  | Allowed   | Outliers | Percentiles |     |
|-----|-------|----------------|-----------|-----------|----------|-------------|-----|
| 1   | A     | 592/617 (96%)  | 531 (90%) | 61 (10%)  | 0        | 100         | 100 |
| 1   | C     | 592/617 (96%)  | 547 (92%) | 45 (8%)   | 0        | 100         | 100 |
| 1   | E     | 592/617 (96%)  | 544 (92%) | 48 (8%)   | 0        | 100         | 100 |
| 2   | B     | 464/517 (90%)  | 409 (88%) | 55 (12%)  | 0        | 100         | 100 |
| 2   | D     | 466/517 (90%)  | 420 (90%) | 46 (10%)  | 0        | 100         | 100 |
| 2   | F     | 467/517 (90%)  | 429 (92%) | 38 (8%)   | 0        | 100         | 100 |
| 3   | G     | 223/233 (96%)  | 216 (97%) | 7 (3%)    | 0        | 100         | 100 |
| 3   | I     | 223/233 (96%)  | 210 (94%) | 13 (6%)   | 0        | 100         | 100 |
| 3   | K     | 223/233 (96%)  | 206 (92%) | 17 (8%)   | 0        | 100         | 100 |
| 4   | H     | 109/122 (89%)  | 101 (93%) | 8 (7%)    | 0        | 100         | 100 |
| 4   | J     | 109/122 (89%)  | 103 (94%) | 6 (6%)    | 0        | 100         | 100 |
| 4   | L     | 109/122 (89%)  | 104 (95%) | 5 (5%)    | 0        | 100         | 100 |
| 5   | M     | 216/256 (84%)  | 200 (93%) | 16 (7%)   | 0        | 100         | 100 |
| 6   | N     | 113/118 (96%)  | 99 (88%)  | 14 (12%)  | 0        | 100         | 100 |
| 7   | O     | 390/392 (100%) | 329 (84%) | 61 (16%)  | 0        | 100         | 100 |
| 8   | P     | 448/469 (96%)  | 381 (85%) | 67 (15%)  | 0        | 100         | 100 |
| 9   | Q     | 741/840 (88%)  | 622 (84%) | 119 (16%) | 0        | 100         | 100 |
| 10  | S     | 342/345 (99%)  | 313 (92%) | 29 (8%)   | 0        | 100         | 100 |
| 11  | T     | 198/213 (93%)  | 181 (91%) | 17 (9%)   | 0        | 100         | 100 |
| 12  | U     | 155/164 (94%)  | 146 (94%) | 9 (6%)    | 0        | 100         | 100 |
| 13  | V     | 157/160 (98%)  | 143 (91%) | 14 (9%)   | 0        | 100         | 100 |
| 13  | W     | 157/160 (98%)  | 153 (98%) | 4 (2%)    | 0        | 100         | 100 |
| 13  | X     | 157/160 (98%)  | 142 (90%) | 15 (10%)  | 0        | 100         | 100 |
| 13  | Y     | 158/160 (99%)  | 148 (94%) | 10 (6%)   | 0        | 100         | 100 |
| 13  | Z     | 157/160 (98%)  | 141 (90%) | 16 (10%)  | 0        | 100         | 100 |

*Continued on next page...*

Continued from previous page...

| Mol | Chain | Analysed        | Favoured   | Allowed   | Outliers | Percentiles |     |
|-----|-------|-----------------|------------|-----------|----------|-------------|-----|
| 13  | a     | 157/160 (98%)   | 147 (94%)  | 10 (6%)   | 0        | 100         | 100 |
| 13  | b     | 158/160 (99%)   | 144 (91%)  | 14 (9%)   | 0        | 100         | 100 |
| 13  | c     | 157/160 (98%)   | 141 (90%)  | 16 (10%)  | 0        | 100         | 100 |
| 14  | d     | 67/73 (92%)     | 61 (91%)   | 6 (9%)    | 0        | 100         | 100 |
| 15  | e     | 50/265 (19%)    | 47 (94%)   | 3 (6%)    | 0        | 100         | 100 |
| 16  | f     | 74/85 (87%)     | 61 (82%)   | 13 (18%)  | 0        | 100         | 100 |
| All | All   | 8221/8967 (92%) | 7419 (90%) | 802 (10%) | 0        | 100         | 100 |

There are no Ramachandran outliers to report.

### 5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all PDB entries followed by that with respect to all EM entries.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

| Mol | Chain | Analysed       | Rotameric  | Outliers | Percentiles |     |
|-----|-------|----------------|------------|----------|-------------|-----|
| 1   | A     | 498/516 (96%)  | 498 (100%) | 0        | 100         | 100 |
| 1   | C     | 498/516 (96%)  | 498 (100%) | 0        | 100         | 100 |
| 1   | E     | 498/516 (96%)  | 498 (100%) | 0        | 100         | 100 |
| 2   | B     | 400/444 (90%)  | 400 (100%) | 0        | 100         | 100 |
| 2   | D     | 402/444 (90%)  | 402 (100%) | 0        | 100         | 100 |
| 2   | F     | 403/444 (91%)  | 403 (100%) | 0        | 100         | 100 |
| 3   | G     | 202/208 (97%)  | 202 (100%) | 0        | 100         | 100 |
| 3   | I     | 202/208 (97%)  | 202 (100%) | 0        | 100         | 100 |
| 3   | K     | 202/208 (97%)  | 202 (100%) | 0        | 100         | 100 |
| 4   | H     | 92/102 (90%)   | 92 (100%)  | 0        | 100         | 100 |
| 4   | J     | 92/102 (90%)   | 92 (100%)  | 0        | 100         | 100 |
| 4   | L     | 92/102 (90%)   | 92 (100%)  | 0        | 100         | 100 |
| 5   | M     | 190/221 (86%)  | 190 (100%) | 0        | 100         | 100 |
| 6   | N     | 102/104 (98%)  | 102 (100%) | 0        | 100         | 100 |
| 7   | O     | 348/348 (100%) | 348 (100%) | 0        | 100         | 100 |

Continued on next page...

Continued from previous page...

| Mol | Chain | Analysed        | Rotameric   | Outliers | Percentiles |     |
|-----|-------|-----------------|-------------|----------|-------------|-----|
| 8   | P     | 418/432 (97%)   | 418 (100%)  | 0        | 100         | 100 |
| 9   | Q     | 657/728 (90%)   | 657 (100%)  | 0        | 100         | 100 |
| 10  | S     | 308/309 (100%)  | 308 (100%)  | 0        | 100         | 100 |
| 11  | T     | 156/168 (93%)   | 156 (100%)  | 0        | 100         | 100 |
| 12  | U     | 119/125 (95%)   | 119 (100%)  | 0        | 100         | 100 |
| 13  | V     | 118/119 (99%)   | 118 (100%)  | 0        | 100         | 100 |
| 13  | W     | 118/119 (99%)   | 118 (100%)  | 0        | 100         | 100 |
| 13  | X     | 118/119 (99%)   | 118 (100%)  | 0        | 100         | 100 |
| 13  | Y     | 119/119 (100%)  | 119 (100%)  | 0        | 100         | 100 |
| 13  | Z     | 118/119 (99%)   | 118 (100%)  | 0        | 100         | 100 |
| 13  | a     | 118/119 (99%)   | 118 (100%)  | 0        | 100         | 100 |
| 13  | b     | 119/119 (100%)  | 119 (100%)  | 0        | 100         | 100 |
| 13  | c     | 118/119 (99%)   | 118 (100%)  | 0        | 100         | 100 |
| 14  | d     | 62/65 (95%)     | 62 (100%)   | 0        | 100         | 100 |
| 15  | e     | 47/244 (19%)    | 47 (100%)   | 0        | 100         | 100 |
| 16  | f     | 63/72 (88%)     | 63 (100%)   | 0        | 100         | 100 |
| All | All   | 6997/7578 (92%) | 6997 (100%) | 0        | 100         | 100 |

There are no protein residues with a non-rotameric sidechain to report.

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 36 such sidechains are listed below:

| Mol | Chain | Res | Type |
|-----|-------|-----|------|
| 10  | S     | 7   | ASN  |
| 16  | f     | 72  | GLN  |
| 10  | S     | 333 | ASN  |
| 13  | a     | 53  | ASN  |
| 4   | J     | 34  | GLN  |

### 5.3.3 RNA

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

## 5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

## 5.6 Ligand geometry [i](#)

There are no ligands in this entry.

## 5.7 Other polymers [i](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.


## 6 Map visualisation [i](#)

This section contains visualisations of the EMDB entry EMD-31540. These allow visual inspection of the internal detail of the map and identification of artifacts.


Images derived from a raw map, generated by summing the deposited half-maps, are presented below the corresponding image components of the primary map to allow further visual inspection and comparison with those of the primary map.

### 6.1 Orthogonal projections [i](#)


#### 6.1.1 Primary map



X




Y




Z


#### 6.1.2 Raw map



X



Y




Z


The images above show the map projected in three orthogonal directions.

## 6.2 Central slices [i](#)


### 6.2.1 Primary map



X Index: 200




Y Index: 200




Z Index: 200


### 6.2.2 Raw map



X Index: 200



Y Index: 200




Z Index: 200


The images above show central slices of the map in three orthogonal directions.

## 6.3 Largest variance slices [i](#)


### 6.3.1 Primary map



X Index: 205




Y Index: 216




Z Index: 126


### 6.3.2 Raw map



X Index: 203



Y Index: 215




Z Index: 126


The images above show the largest variance slices of the map in three orthogonal directions.

## 6.4 Orthogonal standard-deviation projections (False-color) [i](#)


### 6.4.1 Primary map



X




Y




Z


### 6.4.2 Raw map



X



Y




Z

The images above show the map standard deviation projections with false color in three orthogonal directions. Minimum values are shown in green, max in blue, and dark to light orange shades represent small to large values respectively.


## 6.5 Orthogonal surface views [i](#)

### 6.5.1 Primary map



The images above show the 3D surface view of the map at the recommended contour level 0.02. These images, in conjunction with the slice images, may facilitate assessment of whether an appropriate contour level has been provided.

### 6.5.2 Raw map



These images show the 3D surface of the raw map. The raw map's contour level was selected so that its surface encloses the same volume as the primary map does at its recommended contour level.


## 6.6 Mask visualisation [i](#)

This section was not generated. No masks/segmentation were deposited.

## 7 Map analysis [i](#)


This section contains the results of statistical analysis of the map.

### 7.1 Map-value distribution [i](#)



The map-value distribution is plotted in 128 intervals along the x-axis. The y-axis is logarithmic. A spike in this graph at zero usually indicates that the volume has been masked.


## 7.2 Volume estimate [i](#)



The volume at the recommended contour level is 626 nm<sup>3</sup>; this corresponds to an approximate mass of 565 kDa.

The volume estimate graph shows how the enclosed volume varies with the contour level. The recommended contour level is shown as a vertical line and the intersection between the line and the curve gives the volume of the enclosed surface at the given level.

### 7.3 Rotationally averaged power spectrum [i](#)




\*Reported resolution corresponds to spatial frequency of 0.152 Å<sup>-1</sup>

## 8 Fourier-Shell correlation [i](#)

Fourier-Shell Correlation (FSC) is the most commonly used method to estimate the resolution of single-particle and subtomogram-averaged maps. The shape of the curve depends on the imposed symmetry, mask and whether or not the two 3D reconstructions used were processed from a common reference. The reported resolution is shown as a black line. A curve is displayed for the half-bit criterion in addition to lines showing the 0.143 gold standard cut-off and 0.5 cut-off.

### 8.1 FSC [i](#)



\*Reported resolution corresponds to spatial frequency of 0.152 Å<sup>-1</sup>

## 8.2 Resolution estimates [i](#)


| Resolution estimate (Å)   | Estimation criterion (FSC cut-off) |       |          |
|---------------------------|------------------------------------|-------|----------|
|                           | 0.143                              | 0.5   | Half-bit |
| Reported by author        | 6.60                               | -     | -        |
| Author-provided FSC curve | 6.60                               | 8.67  | 6.89     |
| Unmasked-calculated*      | 9.07                               | 18.69 | 9.55     |

\*Resolution estimate based on FSC curve calculated by comparison of deposited half-maps. The value from deposited half-maps intersecting FSC 0.143 CUT-OFF 9.07 differs from the reported value 6.6 by more than 10 %

## 9 Map-model fit [i](#)


This section contains information regarding the fit between EMDB map EMD-31540 and PDB model 7FDC. Per-residue inclusion information can be found in section 3 on page 9.

### 9.1 Map-model overlay [i](#)




The images above show the 3D surface view of the map at the recommended contour level 0.02 at 50% transparency in yellow overlaid with a ribbon representation of the model coloured in blue. These images allow for the visual assessment of the quality of fit between the atomic model and the map.

## 9.2 Q-score mapped to coordinate model [i](#)




The images above show the model with each residue coloured according its Q-score. This shows their resolvability in the map with higher Q-score values reflecting better resolvability. Please note: Q-score is calculating the resolvability of atoms, and thus high values are only expected at resolutions at which atoms can be resolved. Low Q-score values may therefore be expected for many entries.

## 9.3 Atom inclusion mapped to coordinate model [i](#)



The images above show the model with each residue coloured according to its atom inclusion. This shows to what extent they are inside the map at the recommended contour level (0.02).
































































## 9.4 Atom inclusion [i](#)



At the recommended contour level, 53% of all backbone atoms, 47% of all non-hydrogen atoms, are inside the map.

## 9.5 Map-model fit summary

The table lists the average atom inclusion at the recommended contour level (0.02) and Q-score for the entire model and for each chain.

| Chain | Atom inclusion   | Q-score   |
|-------|--|---|
| All   |  0.4670   |  0.0690    |
| A     |  0.5100   |  0.0370    |
| B     |  0.5700   |  0.0780    |
| C     |  0.6670   |  0.1330    |
| D     |  0.5900   |  0.0870    |
| E     |  0.5030   |  0.0320    |
| F     |  0.4930   |  0.0300    |
| G     |  0.3170   |  0.0200    |
| H     |  0.1460   |  0.0520    |
| I     |  0.6010   |  0.0720    |
| J     |  0.5620   |  0.0950    |
| K     |  0.6380   |  0.1250    |
| L     |  0.5790   |  0.1490    |
| M     |  0.5560   |  0.0960    |
| N     |  0.5350  |  0.1030   |
| O     |  0.3780 |  0.1050  |
| P     |  0.4100 |  0.1250  |
| Q     |  0.4640 |  0.0870  |
| S     |  0.4250 |  0.0720  |
| T     |  0.3460 |  0.0220  |
| U     |  0.3110 |  0.0500  |
| V     |  0.3180 |  0.0350  |
| W     |  0.3160 |  -0.0040 |
| X     |  0.4330 |  0.0430  |
| Y     |  0.5000 |  0.0990  |
| Z     |  0.3270 |  0.0460  |
| a     |  0.2130 |  -0.0100 |
| b     |  0.2070 |  0.0150  |
| c     |  0.3030 |  0.0070  |
| d     |  0.2750 |  -0.0120 |
| e     |  0.1600 |  0.0010  |
| f     |  0.2780 |  0.0350  |

