



# wwPDB X-ray Structure Validation Summary Report ⓘ

Mar 18, 2026 – 05:04 AM UTC

PDB ID : 7FF1 / pdb\_00007ff1  
Title : Structure of C34E136G/N36  
Authors : Yu, D.W.; Qin, B.  
Deposited on : 2021-07-22  
Resolution : 1.69 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4-5-2 with Phenix2.0  
Mogul : 2022.3.0, CSD as543be (2022)  
Xtriage (Phenix) : 2.0  
EDS : 3.0  
Percentile statistics : 20250101.v01 (using entries in the PDB archive January 1st 2025)  
CCP4 : 9.0.010 (Gargrove)  
Density-Fitness : 1.0.12  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : 2.49


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

*X-RAY DIFFRACTION*

The reported resolution of this entry is 1.69 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	180053	1054 (1.68-1.68)
Clashscore	190562	1078 (1.68-1.68)
Ramachandran outliers	187476	1068 (1.68-1.68)
Sidechain outliers	187428	1067 (1.68-1.68)
RSRZ outliers	180081	1055 (1.68-1.68)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	B	37	 70% 22% 8%
1	E	37	 73% 24% .
1	N	37	 76% 19% 5%
2	A	35	 80% 14% 6%
2	C	35	 83% 17%

*Continued on next page...*

Continued from previous page...

Mol	Chain	Length	Quality of chain
2	D	35	<div><div></div><div>80%</div><div>11%</div><div>9%</div></div>

## 2 Entry composition

There are 3 unique types of molecules in this entry. The entry contains 1900 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called gp41 N36.

Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	Trace
1	N	37	Total	C	N	O	3	0	0
			293	186	56	51			
1	B	37	Total	C	N	O	3	0	0
			293	186	56	51			
1	E	37	Total	C	N	O	3	0	0
			293	186	56	51			

There are 3 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
N	545	ACE	-	acetylation	UNP C7F357
B	545	ACE	-	acetylation	UNP C7F357
E	545	ACE	-	acetylation	UNP C7F357

- Molecule 2 is a protein called gp41 C34E136G.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
2	C	35	Total	C	N	O	S	3	0	0
			296	183	50	62	1			
2	A	35	Total	C	N	O	S	3	0	0
			296	183	50	62	1			
2	D	35	Total	C	N	O	S	3	0	0
			296	183	50	62	1			

- Molecule 3 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	N	29	Total	O	0	0
			29	29		
3	C	16	Total	O	0	0
			16	16		

*Continued on next page...*


*Continued from previous page...*


Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	B	16	Total 16	O 16	0	0
3	A	27	Total 27	O 27	0	0
3	E	19	Total 19	O 19	0	0
3	D	26	Total 26	O 26	0	0

### 3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: gp41 N36

Chain N:  76% 19% 5%




- Molecule 1: gp41 N36

Chain B:  70% 22% 8%




- Molecule 1: gp41 N36

Chain E:  73% 24% 3%





- Molecule 2: gp41 C34E136G

Chain C:  83% 17%





- Molecule 2: gp41 C34E136G

Chain A:  80% 14% 6%



- Molecule 2: gp41 C34E136G



## 4 Data and refinement statistics

Property	Value	Source
Space group	C 1 2 1	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	88.09Å 50.39Å 56.91Å 90.00° 90.13° 90.00°	Depositor
Resolution (Å)	43.74 – 1.69 43.74 – 1.69	Depositor EDS
% Data completeness (in resolution range)	99.7 (43.74-1.69) 97.3 (43.74-1.69)	Depositor EDS
$R_{merge}$	0.04	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	1.19 (at 1.69Å)	Xtriage
Refinement program	PHENIX 1.10.1_2155	Depositor
R, $R_{free}$	0.216 , 0.224 0.219 , 0.225	Depositor DCC
$R_{free}$ test set	1272 reflections (4.52%)	wwPDB-VP
Wilson B-factor (Å <sup>2</sup> )	36.4	Xtriage
Anisotropy	0.249	Xtriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.32 , 37.1	EDS
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.47$ , $\langle L^2 \rangle = 0.30$	Xtriage
Estimated twinning fraction	0.046 for -1/2*h-3/2*k,-1/2*h+1/2*k,-l 0.054 for -1/2*h+3/2*k,1/2*h+1/2*k,-l 0.227 for 1/2*h-3/2*k,-1/2*h-1/2*k,-l 0.240 for 1/2*h+3/2*k,1/2*h-1/2*k,-l 0.049 for -h,-k,l	Xtriage
$F_o, F_c$ correlation	0.97	EDS
Total number of atoms	1900	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	44.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 15.38% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.



## 5 Model quality [i](#)

### 5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: ACE

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# $ Z  > 5$	RMSZ	# $ Z  > 5$
1	B	2.04	7/293 (2.4%)	0.83	0/396
1	E	2.01	2/293 (0.7%)	0.89	2/396 (0.5%)
1	N	1.80	4/293 (1.4%)	0.87	0/396
2	A	2.02	5/299 (1.7%)	0.95	1/404 (0.2%)
2	C	1.81	2/299 (0.7%)	0.83	0/404
2	D	1.88	7/299 (2.3%)	0.89	0/404
All	All	1.93	27/1776 (1.5%)	0.88	3/2400 (0.1%)

The worst 5 of 27 bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	A	627	ACE	C-N	-17.46	0.98	1.33
1	E	545	ACE	C-N	-14.99	1.03	1.33
1	B	579	ARG	NE-CZ	-10.82	1.21	1.33
1	B	579	ARG	CZ-NH1	-8.61	1.20	1.32
2	C	648	GLU	CD-OE1	-8.35	1.09	1.25

All (3) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	E	580	ILE	CA-C-N	-6.14	110.64	121.70
1	E	580	ILE	C-N-CA	-6.14	110.64	121.70
2	A	654	GLU	CB-CG-CD	5.76	122.40	112.60

There are no chirality outliers.

There are no planarity outliers.

## 5.2 Too-close contacts ⓘ

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	B	293	0	313	37	0
1	E	293	0	312	34	0
1	N	293	0	313	28	0
2	A	296	0	271	4	0
2	C	296	0	272	7	1
2	D	296	0	271	10	0
3	A	27	0	0	2	0
3	B	16	0	0	5	0
3	C	16	0	0	2	1
3	D	26	0	0	3	1
3	E	19	0	0	3	1
3	N	29	0	0	6	0
All	All	1900	0	1752	82	2

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 24.

The worst 5 of 82 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:580:ILE:CD1	1:E:580:ILE:HD11	1.18	1.60
1:B:580:ILE:HD13	1:E:580:ILE:CD1	1.29	1.57
2:C:628:TRP:CE2	1:E:577:GLN:NE2	1.98	1.30
1:N:580:ILE:CD1	1:B:580:ILE:HD11	1.63	1.28
1:B:580:ILE:O	1:B:581:LEU:HD23	1.18	1.24

All (2) symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
2:C:629:MET:SD	3:D:725:HOH:O[4_556]	1.88	0.32
3:C:709:HOH:O	3:E:614:HOH:O[1_554]	2.19	0.01

## 5.3 Torsion angles [i](#)

### 5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	B	35/37 (95%)	33 (94%)	1 (3%)	1 (3%)	3	0
1	E	35/37 (95%)	35 (100%)	0	0	100	100
1	N	35/37 (95%)	34 (97%)	0	1 (3%)	3	0
2	A	33/35 (94%)	33 (100%)	0	0	100	100
2	C	33/35 (94%)	33 (100%)	0	0	100	100
2	D	33/35 (94%)	33 (100%)	0	0	100	100
All	All	204/216 (94%)	201 (98%)	1 (0%)	2 (1%)	12	2

All (2) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	B	580	ILE
1	N	580	ILE

### 5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	B	31/31 (100%)	30 (97%)	1 (3%)	34	9
1	E	31/31 (100%)	31 (100%)	0	100	100
1	N	31/31 (100%)	31 (100%)	0	100	100
2	A	33/33 (100%)	33 (100%)	0	100	100
2	C	33/33 (100%)	33 (100%)	0	100	100

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
2	D	33/33 (100%)	33 (100%)	0	100	100
All	All	192/192 (100%)	191 (100%)	1 (0%)	81	72

All (1) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	B	565	LEU

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (4) such sidechains are listed below:

Mol	Chain	Res	Type
1	N	564	HIS
1	E	553	ASN
1	E	564	HIS
1	E	577	GLN

### 5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

## 5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

## 5.6 Ligand geometry [i](#)

There are no ligands in this entry.

## 5.7 Other polymers [i](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues

The following chains have linkage breaks:

Mol	Chain	Number of breaks
1	E	1
2	A	1

All chain breaks are listed below:

Model	Chain	Residue-1	Atom-1	Residue-2	Atom-2	Distance (Å)
1	E	545:ACE	C	546:SER	N	1.03
1	A	627:ACE	C	628:TRP	N	0.98

## 6 Fit of model and data [i](#)

### 6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	B	36/37 (97%)	-0.79	0 100 100	32, 40, 69, 79	0
1	E	36/37 (97%)	-0.75	0 100 100	29, 37, 60, 70	0
1	N	36/37 (97%)	-0.77	0 100 100	29, 38, 61, 76	0
2	A	34/35 (97%)	-0.87	0 100 100	31, 43, 58, 65	0
2	C	34/35 (97%)	-0.82	0 100 100	32, 44, 57, 66	0
2	D	34/35 (97%)	-0.87	0 100 100	34, 42, 59, 61	0
All	All	210/216 (97%)	-0.81	0 100 100	29, 41, 61, 79	0

There are no RSRZ outliers to report.

### 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

### 6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

### 6.4 Ligands [i](#)

There are no ligands in this entry.

### 6.5 Other polymers [i](#)

There are no such residues in this entry.