



Full wwPDB X-ray Structure Validation Report ⓘ

Mar 8, 2026 – 03:42 AM UTC

PDB ID : 7FTY / pdb_00007fty
Title : Crystal Structure of human cyclic GMP-AMP synthase in complex with 2-[(4-phenylphenyl)methylamino]-5-propyl-4H-[1,2,4]triazolo[1,5-a]pyrimidin-7-one :2,2,2-trifluoroacetic acid
Authors : Leibrock, L.; Benz, J.; Groebke-Zbinden, K.; Rudolph, M.G.
Deposited on : 2023-02-08
Resolution : 2.30 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4-5-2 with Phenix2.0
Mogul	:	2022.3.0, CSD as543be (2022)
Xtriage (Phenix)	:	2.0
EDS	:	3.0
Buster-report	:	wwPDB partial adaption of 1.1.7 (2018)
Percentile statistics	:	20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4	:	9.0.010 (Gargrove)
Density-Fitness	:	1.0.12
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.30 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	6319 (2.30-2.30)
Clashscore	190562	6919 (2.30-2.30)
Ramachandran outliers	187476	6854 (2.30-2.30)
Sidechain outliers	187428	6854 (2.30-2.30)
RSRZ outliers	180081	6325 (2.30-2.30)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	362	<div> <div>14%</div> <div>79%</div> <div>17%</div> <div>..</div> </div>

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.


- Molecule 1 is a protein called Cyclic GMP-AMP synthase.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	352	Total	C	N	O	S	0	0	0
			2896	1853	502	526	15			

- Molecule 2 is ZINC ION (CCD ID: ZN) (formula: Zn).

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
2	A	1	Total Zn 1 1	0	0

- Molecule 3 is (8R)-2-([(1,1'-biphenyl]-4-yl)methyl]amino}-5-propyl[1,2,4]triazolo[1,5-a]pyrimidin-7(4H)-one (CCD ID: YQZ) (formula: C₂₁H₂₁N₅O) (labeled as "Ligand of Interest" by depositor).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
3	A	1	Total	C	N	O	0	0
			27	21	5	1		


- Molecule 4 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	A	33	Total 33	O 33	0	0

3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

• Molecule 1: Cyclic GMP-AMP synthase



4 Data and refinement statistics

Property	Value	Source
Space group	P 2 ₁ 2 ₁ 2 ₁	Depositor
Cell constants a, b, c, α , β , γ	45.06Å 59.24Å 164.24Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	82.12 – 2.30 82.12 – 2.30	Depositor EDS
% Data completeness (in resolution range)	85.5 (82.12-2.30) 82.8 (82.12-2.30)	Depositor EDS
R_{merge}	0.04	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	0.87 (at 2.29Å)	Xtriage
Refinement program	BUSTER 2.11.7	Depositor
R, R_{free}	0.230 , 0.290 0.236 , 0.292	Depositor DCC
R_{free} test set	888 reflections (4.35%)	wwPDB-VP
Wilson B-factor (Å ²)	69.1	Xtriage
Anisotropy	0.039	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.28 , 67.6	EDS
L-test for twinning ²	$\langle L \rangle = 0.47$, $\langle L^2 \rangle = 0.30$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.94	EDS
Total number of atoms	2957	wwPDB-VP
Average B, all atoms (Å ²)	93.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 6.72% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: ZN, YQZ

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.85	0/2952	1.39	15/3956 (0.4%)

There are no bond length outliers.

All (15) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed($^{\circ}$)	Ideal($^{\circ}$)
1	A	214	TYR	N-CA-C	-7.75	104.34	113.88
1	A	392	LYS	N-CA-C	-7.30	102.71	111.69
1	A	175	SER	CA-C-N	6.69	129.14	120.44
1	A	175	SER	C-N-CA	6.69	129.14	120.44
1	A	456	ASP	CA-CB-CG	5.74	118.34	112.60
1	A	462	LEU	CA-C-N	5.52	127.61	120.44
1	A	462	LEU	C-N-CA	5.52	127.61	120.44
1	A	181	THR	CA-C-N	5.43	127.88	120.54
1	A	181	THR	C-N-CA	5.43	127.88	120.54
1	A	200	ASP	CA-C-N	5.29	128.76	120.82
1	A	200	ASP	C-N-CA	5.29	128.76	120.82
1	A	441	ALA	CA-C-N	5.25	127.62	120.54
1	A	441	ALA	C-N-CA	5.25	127.62	120.54
1	A	279	LYS	CA-C-N	5.18	127.18	120.44
1	A	279	LYS	C-N-CA	5.18	127.18	120.44

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen

atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2896	0	2946	24	0
2	A	1	0	0	0	0
3	A	27	0	0	0	0
4	A	33	0	0	0	0
All	All	2957	0	2946	24	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 4.

All (24) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:256:ASN:HB3	1:A:257:PRO:HD3	1.72	0.71
1:A:192:HIS:O	1:A:196:ARG:HG2	2.02	0.59
1:A:305:SER:HB2	1:A:321:THR:HG23	1.86	0.57
1:A:262:LEU:HD22	1:A:276:MET:HE2	1.86	0.56
1:A:324:LEU:O	1:A:358:TYR:HA	2.06	0.55
1:A:168:VAL:HG21	1:A:519:PHE:HB3	1.93	0.50
1:A:245:THR:OG1	1:A:248:TYR:HB2	2.11	0.50
1:A:240:GLU:HB2	1:A:252:LYS:HE2	1.96	0.47
1:A:277:LEU:HD21	1:A:361:PRO:HG2	1.96	0.47
1:A:352:LEU:O	1:A:355:LYS:HB2	2.15	0.47
1:A:420:LEU:HD22	1:A:507:GLN:HB3	1.97	0.46
1:A:164:LYS:O	1:A:168:VAL:HG23	2.16	0.45
1:A:168:VAL:CG2	1:A:519:PHE:HB3	2.46	0.45
1:A:305:SER:HA	1:A:306:PRO:C	2.42	0.45
1:A:459:ASP:O	1:A:460:LEU:C	2.59	0.44
1:A:334:THR:HA	1:A:484:PHE:O	2.17	0.44
1:A:344:LEU:HD11	1:A:386:ILE:HG23	1.99	0.43
1:A:234:VAL:HB	1:A:237:ILE:HD11	2.01	0.43
1:A:221:SER:O	1:A:223:PRO:HD3	2.18	0.43
1:A:237:ILE:HD12	1:A:326:SER:HB2	2.00	0.42
1:A:359:LEU:HB3	1:A:375:TRP:HB3	2.01	0.42
1:A:475:LEU:HD21	1:A:504:LEU:HD23	2.02	0.42
1:A:168:VAL:O	1:A:172:LEU:HG	2.20	0.41
1:A:209:LEU:HD13	1:A:229:MET:SD	2.61	0.41

There are no symmetry-related clashes.

5.3 Torsion angles

5.3.1 Protein backbone

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	346/362 (96%)	325 (94%)	17 (5%)	4 (1%)	10	12

All (4) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	258	LYS
1	A	255	ARG
1	A	256	ASN
1	A	345	SER

5.3.2 Protein sidechains

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	325/334 (97%)	308 (95%)	17 (5%)	21	31

All (17) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	163	SER
1	A	176	ARG
1	A	194	LEU
1	A	201	SER
1	A	209	LEU
1	A	220	ILE
1	A	232	LEU

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	A	240	GLU
1	A	260	ASN
1	A	279	LYS
1	A	297	ILE
1	A	438	VAL
1	A	457	ARG
1	A	462	LEU
1	A	497	ASP
1	A	500	SER
1	A	509	GLU

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (9) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	192	HIS
1	A	210	ASN
1	A	217	HIS
1	A	260	ASN
1	A	264	GLN
1	A	363	HIS
1	A	449	ASN
1	A	451	GLN
1	A	514	ASN

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates ⓘ

There are no oligosaccharides in this entry.

5.6 Ligand geometry ⓘ

Of 2 ligands modelled in this entry, 1 is monoatomic - leaving 1 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
3	YQZ	A	602	-	29,30,30	2.12	8 (27%)	35,41,41	0.89	2 (5%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	YQZ	A	602	-	-	3/12/12/12	0/4/4/4

All (8) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
3	A	602	YQZ	C23-N17	6.55	1.42	1.35
3	A	602	YQZ	C15-N16	4.76	1.39	1.33
3	A	602	YQZ	C15-N24	3.40	1.43	1.36
3	A	602	YQZ	C13-N14	2.60	1.51	1.46
3	A	602	YQZ	C20-C18	2.60	1.49	1.43
3	A	602	YQZ	C3-C2	2.26	1.43	1.38
3	A	602	YQZ	C12-C7	2.13	1.43	1.39
3	A	602	YQZ	C6-C5	2.09	1.43	1.39

All (2) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	A	602	YQZ	C18-N17-N16	2.67	128.91	123.58
3	A	602	YQZ	C20-C21-N22	2.22	121.38	119.54

There are no chirality outliers.

All (3) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
3	A	602	YQZ	N16-C15-N14-C13

Continued on next page...


Continued from previous page...

Mol	Chain	Res	Type	Atoms
3	A	602	YQZ	N24-C15-N14-C13
3	A	602	YQZ	C2-C13-N14-C15

There are no ring outliers.

No monomer is involved in short contacts.

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.



5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data ⓘ

6.1 Protein, DNA and RNA chains ⓘ

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	352/362 (97%)	0.93	52 (14%) 5 6	55, 92, 137, 176	0

All (52) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	519	PHE	5.9
1	A	202	ALA	5.8
1	A	246	ARG	5.0
1	A	308	VAL	4.8
1	A	291	ILE	4.7
1	A	203	PHE	4.7
1	A	310	LEU	4.4
1	A	262	LEU	4.2
1	A	276	MET	4.0
1	A	296	VAL	3.8
1	A	322	LEU	3.8
1	A	471	PHE	3.6
1	A	179	ILE	3.5
1	A	503	PHE	3.5
1	A	371	GLN	3.4
1	A	201	SER	3.3
1	A	214	TYR	3.2
1	A	204	ARG	3.1
1	A	307	ALA	3.1
1	A	265	PHE	3.0
1	A	516	PHE	2.9
1	A	232	LEU	2.9
1	A	324	LEU	2.9
1	A	318	VAL	2.8
1	A	518	VAL	2.8
1	A	365	LYS	2.8
1	A	487	GLU	2.7

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	A	312	ILE	2.7
1	A	402	GLU	2.7
1	A	416	LEU	2.7
1	A	298	MET	2.6
1	A	472	LEU	2.6
1	A	257	PRO	2.6
1	A	297	ILE	2.6
1	A	415	TYR	2.5
1	A	162	ALA	2.5
1	A	303	GLY	2.5
1	A	438	VAL	2.4
1	A	263	SER	2.4
1	A	468	VAL	2.3
1	A	359	LEU	2.3
1	A	403	LYS	2.3
1	A	286	GLU	2.2
1	A	161	GLY	2.2
1	A	320	ILE	2.2
1	A	215	TYR	2.1
1	A	430	LEU	2.1
1	A	508	ILE	2.1
1	A	443	PHE	2.1
1	A	266	LEU	2.0
1	A	316	ILE	2.0
1	A	288	ILE	2.0

6.2 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates ⓘ


There are no oligosaccharides in this entry.

6.4 Ligands ⓘ

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(\AA^2)	Q<0.9
3	YQZ	A	602	27/27	0.93	0.11	56,66,82,82	0
2	ZN	A	601	1/1	0.99	0.06	63,63,63,63	0

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.



6.5 Other polymers ⓘ

There are no such residues in this entry.