Full wwPDB X-ray Structure Validation Report (i)

Aug 18, 2026 – 08:35 pm BST

PDB ID	:	6FW2 / pdb_00006fw2
Title	:	Crystal Structure of human mARC1
Authors	:	Kubitza, C.; Scheidig, A.
Deposited on	:	2018-03-05
Resolution	:	1.78 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at *validation@mail.wwpdb.org*
A user guide is available at
https://www.wwpdb.org/validation/2017/XrayValidationReportHelp
with specific help available everywhere you see the (i) symbol.

The types of validation reports are described at
http://www.wwpdb.org/validation/2017/FAQs#types.

The following versions of software and data (see references (i)) were used in the production of this report:

MolProbity	:	4-5-2 with Phenix2.0
Mogul	:	1.8.4, CSD as541be (2020)
Xtriage (Phenix)	:	2.0
EDS	:	3.0
Buster-report	:	wwPDB partial adaption of 1.1.7 (2018)
Percentile statistics	:	20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4	:	9.0.015 (Gargrove)
Density-Fitness	:	1.0.12
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.50

1 Overall quality at a glance (i)

The following experimental techniques were used to determine the structure:
X-RAY DIFFRACTION

The reported resolution of this entry is 1.78 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	1365 (1.78-1.78)
Clashscore	190562	1395 (1.78-1.78)
Ramachandran outliers	187476	1382 (1.78-1.78)
Sidechain outliers	187428	1382 (1.78-1.78)
RSRZ outliers	180081	1365 (1.78-1.78)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for >=3, 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions <=5% The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	459	5% 82% 12%

2 Entry composition (i)

There are 6 unique types of molecules in this entry. The entry contains 4072 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Mitochondrial amidoxime-reducing component 1,Endolysin,Mitochondrial amidoxime-reducing component 1.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	444	3562	2242	631	666	23	0	8	0

There are 19 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	48	MET	-	initiating methionine	UNP Q5VT66
A	49	ARG	-	expression tag	UNP Q5VT66
A	50	GLY	-	expression tag	UNP Q5VT66
A	51	SER	-	expression tag	UNP Q5VT66
A	52	MET	-	expression tag	UNP Q5VT66
A	1160	ALA	ALA	linker	UNP D9IEF7
A	1161	TYR	TYR	linker	UNP D9IEF7
A	165	ALA	THR	variant	UNP Q5VT66
A	337	GLN	-	expression tag	UNP Q5VT66
A	338	GLY	-	expression tag	UNP Q5VT66
A	339	THR	-	expression tag	UNP Q5VT66
A	340	VAL	-	expression tag	UNP Q5VT66
A	341	ASP	-	expression tag	UNP Q5VT66
A	342	HIS	-	expression tag	UNP Q5VT66
A	343	HIS	-	expression tag	UNP Q5VT66
A	344	HIS	-	expression tag	UNP Q5VT66
A	345	HIS	-	expression tag	UNP Q5VT66
A	346	HIS	-	expression tag	UNP Q5VT66
A	347	HIS	-	expression tag	UNP Q5VT66

- Molecule 2 is 2-[3-(2-HYDROXY-1,1-DIHYDROXYMETHYL-ETHYLAMINO)-PROPYLAMINO]-2-HYDROXYMETHYL-PROPANE-1,3-DIOL (CCD ID: B3P) (formula: $C_{11}H_{26}N_2O_6$).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
2	A	1	Total	C	N	O	0	0
			19	11	2	6		

- Molecule 3 is MOLYBDATE ION (CCD ID: MOO) (formula: MoO_4).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
3	A	1	Total	Mo	O		0	0
			5	1	4			
3	A	1	Total	Mo	O		0	0
			5	1	4			
3	A	1	Total	Mo	O		0	0
			5	1	4			

Continued on next page...

Continued from previous page...

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
3	A	1	Total Mo O 5 1 4	0	0

- Molecule 4 is PHOSPHATE ION (CCD ID: PO4) (formula: O_4P).



Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
4	A	1	Total O P 5 4 1	0	0

- Molecule 5 is (MOLYBDOPTERIN-S,S)-DIOXO-THIO-MOLYBDENUM(IV) (CCD ID: MTV) (formula: $C_{10}H_{13}MoN_5O_8PS_2$).



Mol	Chain	Residues	Atoms							ZeroOcc	AltConf
5	A	1	Total	C	Mo	N	O	P	S	0	0
			27	10	1	5	8	1	2		

- Molecule 6 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
6	A	439	Total	O	0	0
			439	439		

3 Residue-property plots (i)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density (RSRZ > 2). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

• Molecule 1: Mitochondrial amidoxime-reducing component 1,Endolysin,Mitochondrial amidoxime-reducing component 1



4 Data and refinement statistics (i)

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, α, β, γ	61.13Å 74.82Å 110.73Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	43.57 – 1.78 43.57 – 1.78	Depositor EDS
% Data completeness (in resolution range)	91.5 (43.57-1.78) 91.8 (43.57-1.78)	Depositor EDS
R_{merge}	0.08	Depositor
R_{sym}	(Not available)	Depositor
$< I/\sigma(I) >$ [1]	2.13 (at 1.78Å)	Xtriage
Refinement program	REFMAC	Depositor
R, R_{free}	0.169 , 0.208 0.169 , 0.208	Depositor DCC
R_{free} test set	2288 reflections (4.61%)	wwPDB-VP
Wilson B-factor (Å^2)	22.2	Xtriage
Anisotropy	0.101	Xtriage
Bulk solvent k_{sol}(e/Å^3), B_{sol}(Å^2)	0.37 , 44.8	EDS
L-test for twinning[2]	$< \|L\| > = 0.49$, $< L^2 > = 0.32$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o,F_c correlation	0.96	EDS
Total number of atoms	4072	wwPDB-VP
Average B, all atoms (Å^2)	29.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 5.02% of the height of the origin peak. No significant pseudotranslation is detected.*

[1] Intensities estimated from amplitudes.

[2] Theoretical values of $< |L| >$, $< L^2 >$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: MTV, PO4, MOO, B3P

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	#$\|Z\|$ >5	RMSZ	#$\|Z\|$ >5
1	A	1.53	20/3629 (0.6%)	1.38	21/4912 (0.4%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand.A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	A	0	1

All (20) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	158	GLY	N-CA	7.94	1.52	1.45
1	A	57	THR	CA-C	7.24	1.61	1.52
1	A	1127	ASP	N-CA	6.99	1.54	1.46
1	A	1161	TYR	C-N	6.80	1.42	1.33
1	A	241	ILE	N-CA	6.75	1.54	1.46
1	A	1149	VAL	N-CA	6.26	1.53	1.46
1	A	137	PRO	C-O	-6.00	1.17	1.23
1	A	138	ILE	C-O	-5.98	1.17	1.24
1	A	169	ILE	CA-C	5.68	1.59	1.52
1	A	241	ILE	C-O	-5.54	1.18	1.24
1	A	1016	LYS	CA-C	-5.46	1.46	1.52
1	A	92	ARG	CZ-NH2	5.39	1.40	1.33
1	A	72	VAL	N-CA	-5.37	1.42	1.46
1	A	136	LEU	C-O	-5.33	1.18	1.24
1	A	1046	LEU	N-CA	5.15	1.52	1.46
1	A	1075	VAL	N-CA	5.15	1.52	1.46
1	A	329	GLY	N-CA	5.13	1.52	1.45
1	A	115	ILE	N-CA	5.13	1.52	1.46

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	148	LYS	CA-CB	5.06	1.59	1.53
1	A	1129	ALA	CA-C	5.01	1.59	1.52

All (21) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(^{o})	Ideal(^{o})
1	A	1161	TYR	O-C-N	-9.00	110.62	122.59
1	A	1108	GLU	N-CA-C	7.66	119.27	111.07
1	A	288	LYS	N-CA-C	7.46	122.34	113.23
1	A	122	ASP	N-CA-C	7.40	119.42	111.36
1	A	1150	ILE	N-CA-C	7.22	117.36	110.42
1	A	289	GLU	C-N-CD	6.60	135.13	120.60
1	A	1108	GLU	N-CA-CB	-6.49	100.60	110.01
1	A	322	ASN	CA-C-N	6.02	125.99	120.03
1	A	322	ASN	C-N-CA	6.02	125.99	120.03
1	A	289	GLU	N-CA-C	-5.75	102.37	110.31
1	A	1152	THR	N-CA-C	-5.65	105.04	111.14
1	A	202	LYS	N-CA-C	5.56	119.91	113.18
1	A	121	GLY	CA-C-N	5.56	128.18	120.29
1	A	121	GLY	C-N-CA	5.56	128.18	120.29
1	A	311	SER	N-CA-C	-5.54	97.56	109.81
1	A	225	ASN	N-CA-C	5.53	118.06	111.71
1	A	128	ALA	O-C-N	-5.28	116.34	122.09
1	A	1149	VAL	N-CA-C	-5.22	105.44	110.72
1	A	301	ASP	CA-C-N	5.16	126.30	119.84
1	A	301	ASP	C-N-CA	5.16	126.30	119.84
1	A	1145	ARG	NE-CZ-NH1	-5.02	116.48	121.50

There are no chirality outliers.

All (1) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	A	128	ALA	Mainchain

5.2 Too-close contacts (i)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	3562	0	3589	40	0
2	A	19	0	26	0	0
3	A	20	0	0	1	0
4	A	5	0	0	0	0
5	A	27	0	7	1	0
6	A	439	0	0	11	1
All	All	4072	0	3622	42	1

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 6.

All (42) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:232:VAL:HG11	1:A:290:PRO:HG3	1.54	0.90
1:A:302:PRO:HA	1:A:305:ARG:HG3	1.74	0.70
3:A:1206:MOO:O1	6:A:1301:HOH:O	2.08	0.69
1:A:188:ARG:NH2	6:A:1305:HOH:O	2.25	0.68
1:A:275:LEU:HD13	6:A:1536:HOH:O	1.95	0.65
1:A:275:LEU:CD1	6:A:1536:HOH:O	2.44	0.65
5:A:1209:MTV:MOM1	5:A:1209:MTV:OM1	1.71	0.61
1:A:1076:ARG:HD3	6:A:1332:HOH:O	2.01	0.59
1:A:102[A]:ASN:OD1	6:A:1302:HOH:O	2.17	0.58
1:A:121:GLY:O	1:A:138:ILE:HD11	2.04	0.57
1:A:1080:ARG:CB	1:A:1080:ARG:HH11	2.18	0.56
1:A:1043:LYS:NZ	6:A:1306:HOH:O	2.38	0.56
1:A:1080:ARG:HH11	1:A:1080:ARG:HB2	1.72	0.55
1:A:277:THR:HG21	1:A:288:LYS:HA	1.88	0.54
1:A:217[B]:SER:CB	1:A:251:GLU:OE1	2.56	0.53
1:A:1061:ASP:OD2	1:A:287:ARG:NH2	2.43	0.51
1:A:122:ASP:O	1:A:138:ILE:HD12	2.11	0.50
1:A:217[B]:SER:HB2	1:A:251:GLU:OE1	2.12	0.49
1:A:1080:ARG:HB2	1:A:1080:ARG:NH1	2.26	0.49
1:A:217[A]:SER:OG	1:A:251:GLU:OE1	2.30	0.49
1:A:136:LEU:HB3	1:A:137:PRO:HD2	1.95	0.48
1:A:198:LEU:HD11	1:A:307:LEU:HD11	1.96	0.47
1:A:122:ASP:OD1	1:A:122:ASP:N	2.43	0.47
1:A:106:ALA:HB3	1:A:275:LEU:HD13	1.96	0.47
1:A:121:GLY:O	1:A:138:ILE:CD1	2.63	0.47
1:A:238:ARG:N	1:A:239:PRO:CD	2.77	0.47
1:A:217[B]:SER:HB3	1:A:251:GLU:OE1	2.15	0.46
1:A:107:ARG:HG3	1:A:275:LEU:HD21	1.98	0.46

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:126:LEU:HD12	1:A:134:LEU:HD23	1.98	0.45
1:A:152:HIS:NE2	1:A:209[A]:ASP:OD1	2.47	0.45
1:A:275:LEU:HD12	6:A:1536:HOH:O	2.15	0.44
1:A:148:LYS:NZ	6:A:1312:HOH:O	2.50	0.44
1:A:302:PRO:HA	1:A:305:ARG:CG	2.47	0.43
1:A:288:LYS:HE3	1:A:288:LYS:HB2	1.78	0.43
1:A:235:THR:HG22	6:A:1570:HOH:O	2.18	0.43
1:A:302:PRO:HA	1:A:305:ARG:HD2	2.00	0.43
1:A:251:GLU:HA	1:A:254:TRP:CE3	2.54	0.42
1:A:53:GLN:HE21	1:A:334:LEU:HD23	1.85	0.42
1:A:1083:LYS:HB3	1:A:1083:LYS:HE2	1.75	0.42
1:A:261:ASP:HB2	6:A:1314:HOH:O	2.19	0.41
1:A:251:GLU:HA	1:A:254:TRP:CD2	2.56	0.41
1:A:1146:ALA:O	1:A:1150:ILE:HG13	2.22	0.40

All (1) symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
6:A:1434:HOH:O	6:A:1584:HOH:O[4_545]	2.19	0.01

5.3 Torsion angles (i)

5.3.1 Protein backbone (i)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
1	A	450/459 (98%)	435 (97%)	15 (3%)	0	100 100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles
1	A	393/398 (99%)	390 (99%)	3 (1%)	73 63

All (3) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	122	ASP
1	A	1080	ARG
1	A	296	SER

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (3) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	53	GLN
1	A	1069	GLN
1	A	1132	ASN

5.3.3 RNA

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates

There are no oligosaccharides in this entry.

5.6 Ligand geometry

7 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\#\|Z\| > 2$	Counts	RMSZ	$\#\|Z\| > 2$
5	MTV	A	1209	1	21,30,30	1.96	5 (23%)	20,48,48	3.15	9 (45%)
3	MOO	A	1207	-	2,4,4	4.50	2 (100%)	-		
2	B3P	A	1203	-	18,18,18	1.23	1 (5%)	21,23,23	2.11	5 (23%)
3	MOO	A	1205	-	2,4,4	5.53	2 (100%)	-		
3	MOO	A	1206	-	2,4,4	5.73	2 (100%)	-		
4	PO4	A	1208	-	4,4,4	0.73	0	6,6,6	1.36	1 (16%)
3	MOO	A	1204	-	2,4,4	6.12	2 (100%)	-		

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
5	MTV	A	1209	1	-	0/6/44/44	0/4/4/4
2	B3P	A	1203	-	-	1/28/28/28	-

All (14) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
3	A	1204	MOO	O2-MO	6.38	1.90	1.75
3	A	1206	MOO	O1-MO	6.22	1.89	1.75
3	A	1204	MOO	O1-MO	5.84	1.88	1.75
3	A	1207	MOO	O1-MO	5.64	1.88	1.75
3	A	1205	MOO	O1-MO	5.60	1.88	1.75
3	A	1205	MOO	O2-MO	5.45	1.87	1.75
3	A	1206	MOO	O2-MO	5.20	1.87	1.75
5	A	1209	MTV	C7-C6	4.73	1.57	1.53
5	A	1209	MTV	C9-C10	4.65	1.46	1.38
2	A	1203	B3P	C11-C8	3.11	1.57	1.53
5	A	1209	MTV	O3'-C7	2.98	1.48	1.43
3	A	1207	MOO	O2-MO	2.94	1.82	1.75
5	A	1209	MTV	C7-N8	2.46	1.49	1.45

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
5	A	1209	MTV	C2-N2	2.08	1.39	1.34

All (15) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(^{o})	Ideal(^{o})
5	A	1209	MTV	C2'-C1'-S1'	8.74	125.11	120.15
5	A	1209	MTV	C1'-C2'-S2'	6.70	123.95	120.15
2	A	1203	B3P	C7-C4-C6	-4.91	99.66	110.04
2	A	1203	B3P	O5-C6-C4	-4.66	102.20	111.63
5	A	1209	MTV	C2-N1-C10	3.75	120.19	113.43
2	A	1203	B3P	O3-C11-C8	-3.55	104.45	111.63
5	A	1209	MTV	C2-N3-C4	-3.35	118.99	125.10
5	A	1209	MTV	C9-C4-N3	3.25	122.00	112.31
5	A	1209	MTV	O4-C4-C9	-2.95	120.48	127.24
4	A	1208	PO4	O3-P-O2	2.88	117.20	107.97
5	A	1209	MTV	O3P-P-O2P	2.86	118.57	107.64
5	A	1209	MTV	O3'-C7-N8	-2.78	105.71	108.57
5	A	1209	MTV	C4-C9-N5	2.70	124.00	116.76
2	A	1203	B3P	C3-N1-C4	2.28	119.31	116.08
2	A	1203	B3P	C7-C4-C5	2.03	114.34	110.04

There are no chirality outliers.

All (1) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
2	A	1203	B3P	C9-C8-N2-C2

There are no ring outliers.

2 monomers are involved in 2 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
5	A	1209	MTV	1	0
3	A	1206	MOO	1	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less then 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring

in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.

Ligand MTV A 1209



Bond lengths



Bond angles



Torsions



Rings



5.7 Other polymers

There are no such residues in this entry.

5.8 Polymer linkage issues

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

In the following table, the column labelled '#RSRZ> 2' contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled 'Q< 0.9' lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å^2)	Q<0.9
1	A	444/459 (96%)	0.05	24 (5%) 31 37	8, 23, 61, 126	8 (1%)

All (24) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	307	LEU	6.2
1	A	273[A]	CYS	5.5
1	A	274	ILE	4.7
1	A	302	PRO	3.8
1	A	297	TYR	3.4
1	A	308	TYR	3.3
1	A	301	ASP	3.1
1	A	310	LYS	3.0
1	A	300	CYS	3.0
1	A	312	PRO	3.0
1	A	122	ASP	2.9
1	A	249	TYR	2.8
1	A	275	LEU	2.8
1	A	313	LEU	2.7
1	A	248	VAL	2.6
1	A	311	SER	2.5
1	A	299	GLN	2.4
1	A	335	LEU	2.3
1	A	309	GLY	2.3
1	A	303	SER	2.2
1	A	294	LEU	2.2
1	A	52	MET	2.1
1	A	336	GLY	2.0
1	A	306	LYS	2.0

6.2 Non-standard residues in protein, DNA, RNA chains (i)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates (i)

There are no oligosaccharides in this entry.

6.4 Ligands (i)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å^2)	Q<0.9
4	PO4	A	1208	5/5	0.91	0.09	47,49,64,73	0
3	MOO	A	1206	5/5	0.94	0.12	47,50,61,67	5
2	B3P	A	1203	19/19	0.95	0.06	21,25,28,33	0
3	MOO	A	1204	5/5	0.96	0.10	38,41,50,54	5
5	MTV	A	1209	27/27	0.98	0.05	12,17,29,32	3
3	MOO	A	1205	5/5	0.99	0.07	31,32,34,35	5
3	MOO	A	1207	5/5	0.99	0.07	21,27,30,32	5

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.

Electron density around B3P A 1203:

$2mF_o$-DF_c (at 0.7 rmsd) in gray
mF_o-DF_c (at 3 rmsd) in purple (negative) and green (positive)





Electron density around MTV A 1209:

$2mF_o$-DF_c (at 0.7 rmsd) in gray
mF_o-DF_c (at 3 rmsd) in purple (negative) and green (positive)



6.5 Other polymers

There are no such residues in this entry.