



wwPDB X-ray Structure Validation Summary Report ⓘ

Mar 9, 2026 – 05:39 PM UTC

PDB ID : 2GN6 / pdb_00002gn6
Title : Crystal structure of UDP-GlcNAc inverting 4,6-dehydratase in complex with NADP and UDP-GlcNAc
Authors : Ishiyama, N.; Creuzenet, C.; Lam, J.S.; Berghuis, A.M.
Deposited on : 2006-04-09
Resolution : 2.70 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4-5-2 with Phenix2.0
Mogul	:	2022.3.0, CSD as543be (2022)
Xtriage (Phenix)	:	2.0
EDS	:	3.0
Buster-report	:	wwPDB partial adaption of 1.1.7 (2018)
Percentile statistics	:	20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4	:	9.0.010 (Gargrove)
Density-Fitness	:	1.0.12
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.70 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	3538 (2.70-2.70)
Clashscore	190562	3843 (2.70-2.70)
Ramachandran outliers	187476	3778 (2.70-2.70)
Sidechain outliers	187428	3778 (2.70-2.70)
RSRZ outliers	180081	3538 (2.70-2.70)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	344	17% 48% 44% . .
1	B	344	10% 53% 38% . 5%

2 Entry composition

There are 5 unique types of molecules in this entry. The entry contains 5517 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.


- Molecule 1 is a protein called UDP-GlcNAc C6 dehydratase.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	329	Total	C	N	O	S	0	0	0
			2590	1656	440	479	15			
1	B	327	Total	C	N	O	S	0	0	0
			2573	1647	436	475	15			

There are 22 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	-10	MET	-	cloning artifact	UNP O25511
A	-9	HIS	-	expression tag	UNP O25511
A	-8	HIS	-	expression tag	UNP O25511
A	-7	HIS	-	expression tag	UNP O25511
A	-6	HIS	-	expression tag	UNP O25511
A	-5	HIS	-	expression tag	UNP O25511
A	-4	HIS	-	expression tag	UNP O25511
A	-3	GLY	-	cloning artifact	UNP O25511
A	-2	SER	-	cloning artifact	UNP O25511
A	-1	MET	-	cloning artifact	UNP O25511
A	0	SER	-	cloning artifact	UNP O25511
B	-10	MET	-	cloning artifact	UNP O25511
B	-9	HIS	-	expression tag	UNP O25511
B	-8	HIS	-	expression tag	UNP O25511
B	-7	HIS	-	expression tag	UNP O25511
B	-6	HIS	-	expression tag	UNP O25511
B	-5	HIS	-	expression tag	UNP O25511
B	-4	HIS	-	expression tag	UNP O25511
B	-3	GLY	-	cloning artifact	UNP O25511
B	-2	SER	-	cloning artifact	UNP O25511
B	-1	MET	-	cloning artifact	UNP O25511
B	0	SER	-	cloning artifact	UNP O25511

- Molecule 2 is NADP NICOTINAMIDE-ADENINE-DINUCLEOTIDE PHOSPHATE (CCD ID: NAP) (formula: C₂₁H₂₈N₇O₁₇P₃).



-
- The chemical structure of UD1 is a complex molecule. It features a pyrimidine ring (C2-C6, N1, N3) with a carbonyl group at C2 and a nitrogen at N1. This ring is connected via a nitrogen atom (N1) to a sugar moiety (C1B, C2B, C3B, C4B, C5B, C6B). The sugar moiety is further connected to a phosphate group (P1, O1A, O1B, O1C, O1D, O1E, O1F, O1G, O1H, O1I, O1J, O1K, O1L, O1M, O1N, O1O, O1P, O1Q, O1R, O1S, O1T, O1U, O1V, O1W, O1X, O1Y, O1Z). The phosphate group is connected to a pyrophosphate group (P2, O2A, O2B, O2C, O2D, O2E, O2F, O2G, O2H, O2I, O2J, O2K, O2L, O2M, O2N, O2O, O2P, O2Q, O2R, O2S, O2T, O2U, O2V, O2W, O2X, O2Y, O2Z). The pyrophosphate group is connected to a sugar moiety (C3F, C4F, C5F, C6F, C7F, C8F, C9F, C10F, C11F, C12F, C13F, C14F, C15F, C16F, C17F, C18F, C19F, C20F, C21F, C22F, C23F, C24F, C25F, C26F, C27F, C28F, C29F, C30F, C31F, C32F, C33F, C34F, C35F, C36F, C37F, C38F, C39F, C40F, C41F, C42F, C43F, C44F, C45F, C46F, C47F, C48F, C49F, C50F, C51F, C52F, C53F, C54F, C55F, C56F, C57F, C58F, C59F, C60F, C61F, C62F, C63F, C64F, C65F, C66F, C67F, C68F, C69F, C70F, C71F, C72F, C73F, C74F, C75F, C76F, C77F, C78F, C79F, C80F, C81F, C82F, C83F, C84F, C85F, C86F, C87F, C88F, C89F, C90F, C91F, C92F, C93F, C94F, C95F, C96F, C97F, C98F, C99F, C100F, C101F, C102F, C103F, C104F, C105F, C106F, C107F, C108F, C109F, C110F, C111F, C112F, C113F, C114F, C115F, C116F, C117F, C118F, C119F, C120F, C121F, C122F, C123F, C124F, C125F, C126F, C127F, C128F, C129F, C130F, C131F, C132F, C133F, C134F, C135F, C136F, C137F, C138F, C139F, C140F, C141F, C142F, C143F, C144F, C145F, C146F, C147F, C148F, C149F, C150F, C151F, C152F, C153F, C154F, C155F, C156F, C157F, C158F, C159F, C160F, C161F, C162F, C163F, C164F, C165F, C166F, C167F, C168F, C169F, C170F, C171F, C172F, C173F, C174F, C175F, C176F, C177F, C178F, C179F, C180F, C181F, C182F, C183F, C184F, C185F, C186F, C187F, C188F, C189F, C190F, C191F, C192F, C193F, C194F, C195F, C196F, C197F, C198F, C199F, C200F, C201F, C202F, C203F, C204F, C205F, C206F, C207F, C208F, C209F, C210F, C211F, C212F, C213F, C214F, C215F, C216F, C217F, C218F, C219F, C220F, C221F, C222F, C223F, C224F, C225F, C226F, C227F, C228F, C229F, C230F, C231F, C232F, C233F, C234F, C235F, C236F, C237F, C238F, C239F, C240F, C241F, C242F, C243F, C244F, C245F, C246F, C247F, C248F, C249F, C250F, C251F, C252F, C253F, C254F, C255F, C256F, C257F, C258F, C259F, C260F, C261F, C262F, C263F, C264F, C265F, C266F, C267F, C268F, C269F, C270F, C271F, C272F, C273F, C274F, C275F, C276F, C277F, C278F, C279F, C280F, C281F, C282F, C283F, C284F, C285F, C286F, C287F, C288F, C289F, C290F, C291F, C292F, C293F, C294F, C295F, C296F, C297F, C298F, C299F, C300F, C301F, C302F, C303F, C304F, C305F, C306F, C307F, C308F, C309F, C310F, C311F, C312F, C313F, C314F, C315F, C316F, C317F, C318F, C319F, C320F, C321F, C322F, C323F, C324F, C325F, C326F, C327F, C328F, C329F, C330F, C331F, C332F, C333F, C334F, C335F, C336F, C337F, C338F, C339F, C340F, C341F, C342F, C343F, C344F, C345F, C346F, C347F, C348F, C349F, C350F, C351F, C352F, C353F, C354F, C355F, C356F, C357F, C358F, C359F, C360F, C361F, C362F, C363F, C364F, C365F, C366F, C367F, C368F, C369F, C370F, C371F, C372F, C373F, C374F, C375F, C376F, C377F, C378F, C379F, C380F, C381F, C382F, C383F, C384F, C385F, C386F, C387F, C388F, C389F, C390F, C391F, C392F, C393F, C394F, C395F, C396F, C397F, C398F, C399F, C400F, C401F, C402F, C403F, C404F, C405F, C406F, C407F, C408F, C409F, C410F, C411F, C412F, C413F, C414F, C415F, C416F, C417F, C418F, C419F, C420F, C421F, C422F, C423F, C424F, C425F, C426F, C427F, C428F, C429F, C430F, C431F, C432F, C433F, C434F, C435F, C436F, C437F, C438F, C439F, C440F, C441F, C442F, C443F, C444F, C445F, C446F, C447F, C448F, C449F, C450F, C451F, C452F, C453F, C454F, C455F, C456F, C457F, C458F, C459F, C460F, C461F, C462F, C463F, C464F, C465F, C466F, C467F, C468F, C469F, C470F, C471F, C472F, C473F, C474F, C475F, C476F, C477F, C478F, C479F, C480F, C481F, C482F, C483F, C484F, C485F, C486F, C487F, C488F, C489F, C490F, C491F, C492F, C493F, C494F, C495F, C496F, C497F, C498F, C499F, C500F, C501F, C502F, C503F, C504F, C505F, C506F, C507F, C508F, C509F, C510F, C511F, C512F, C513F, C514F, C515F, C516F, C517F, C518F, C519F, C520F, C521F, C522F, C523F, C524F, C525F, C526F, C527F, C528F, C529F, C530F, C531F, C532F, C533F, C534F, C535F, C536F, C537F, C538F, C539F, C540F, C541F, C542F, C543F, C544F, C545F, C546F, C547F, C548F, C549F, C550F, C551F, C552F, C553F, C554F, C555F, C556F, C557F, C558F, C559F, C560F, C561F, C562F, C563F, C564F, C565F, C566F, C567F, C568F, C569F, C570F, C571F, C572F, C573F, C574F, C575F, C576F, C577F, C578F, C579F, C580F, C581F, C582F, C583F, C584F, C585F, C586F, C587F, C588F, C589F, C590F, C591F, C592F, C593F, C594F, C595F, C596F, C597F, C598F, C599F, C600F, C601F, C602F, C603F, C604F, C605F, C606F, C607F, C608F, C609F, C610F, C611F, C612F, C613F, C614F, C615F, C616F, C617F, C618F, C619F, C620F, C621F, C622F, C623F, C624F, C625F, C626F, C627F, C628F, C629F, C630F, C631F, C632F, C633F, C634F, C635F, C636F, C637F, C638F, C639F, C640F,


Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
3	A	1	Total	C	N	O	P	14	0
			39	17	3	17	2		



Continued from previous page...

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
3	B	1	Total	C	N	O	P	14	0
			39	17	3	17	2		

- Molecule 4 is 2-(N-MORPHOLINO)-ETHANESULFONIC ACID (CCD ID: MES) (formula: $C_6H_{13}NO_4S$).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
4	B	1	Total	C	N	O	S	0	0
			12	6	1	4	1		

- Molecule 5 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
5	A	77	Total	O	0	0
			77	77		
5	B	91	Total	O	0	0
			91	91		

4 Data and refinement statistics

Property	Value	Source
Space group	P 63	Depositor
Cell constants a, b, c, α , β , γ	109.85Å 109.85Å 107.98Å 90.00° 90.00° 120.00°	Depositor
Resolution (Å)	50.00 – 2.70 50.00 – 2.70	Depositor EDS
% Data completeness (in resolution range)	97.6 (50.00-2.70) 97.7 (50.00-2.70)	Depositor EDS
R_{merge}	0.10	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	4.43 (at 2.69Å)	Xtriage
Refinement program	CNS	Depositor
R, R_{free}	0.204 , 0.269 0.203 , 0.267	Depositor DCC
R_{free} test set	2005 reflections (9.86%)	wwPDB-VP
Wilson B-factor (Å ²)	22.0	Xtriage
Anisotropy	0.125	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.36 , 57.6	EDS
L-test for twinning ²	$\langle L \rangle = 0.50$, $\langle L^2 \rangle = 0.33$	Xtriage
Estimated twinning fraction	0.045 for h,-h-k,-l	Xtriage
F_o, F_c correlation	0.92	EDS
Total number of atoms	5517	wwPDB-VP
Average B, all atoms (Å ²)	32.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 5.07% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: MES, UD1, NAP

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.43	0/2641	0.96	13/3568 (0.4%)
1	B	0.43	0/2624	0.98	10/3545 (0.3%)
All	All	0.43	0/5265	0.97	23/7113 (0.3%)

There are no bond length outliers.

The worst 5 of 23 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	B	225	HIS	N-CA-C	-8.56	102.97	113.50
1	B	224	MET	N-CA-C	7.73	121.44	110.23
1	B	45	ASP	N-CA-C	7.50	120.62	109.59
1	A	8	LEU	N-CA-C	-7.13	104.19	113.17
1	B	44	ARG	N-CA-C	7.01	119.96	111.40

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2590	0	2641	181	0
1	B	2573	0	2627	142	0
2	A	48	0	25	4	0
2	B	48	0	25	3	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
3	A	39	0	25	1	0
3	B	39	0	25	0	0
4	B	12	0	13	0	0
5	A	77	0	0	4	0
5	B	91	0	0	0	0
All	All	5517	0	5381	323	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 30.

The worst 5 of 323 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:198:ILE:HG22	1:A:199:THR:H	1.31	0.92
1:A:223:ARG:HA	1:A:301:LYS:HB2	1.56	0.88
1:A:176:GLY:HA2	1:A:183:PRO:HD3	1.54	0.87
1:B:8:LEU:HD22	1:B:13:ILE:HD11	1.56	0.86
1:A:186:LYS:HG2	1:A:246:LEU:HD22	1.58	0.84

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	327/344 (95%)	283 (86%)	36 (11%)	8 (2%)	4	12
1	B	325/344 (94%)	298 (92%)	20 (6%)	7 (2%)	5	14
All	All	652/688 (95%)	581 (89%)	56 (9%)	15 (2%)	5	13

5 of 15 Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	162	SER
1	A	178	ARG
1	A	247	ALA
1	A	308	LYS
1	B	254	ILE

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	290/304 (95%)	280 (97%)	10 (3%)	32	62
1	B	288/304 (95%)	279 (97%)	9 (3%)	35	65
All	All	578/608 (95%)	559 (97%)	19 (3%)	33	63

5 of 19 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	B	173	ASN
1	B	240	THR
1	B	307	GLN
1	B	210	LEU
1	A	244	LYS

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 21 such sidechains are listed below:

Mol	Chain	Res	Type
1	B	136	ASN
1	B	274	HIS
1	B	322	GLN
1	B	286	GLN
1	B	191	ASN

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

5 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
3	UD1	A	335	-	40,41,41	1.22	5 (12%)	59,62,62	1.30	3 (5%)
3	UD1	B	335	-	40,41,41	1.19	3 (7%)	59,62,62	1.32	3 (5%)
2	NAP	B	334	-	50,52,52	1.75	11 (22%)	71,80,80	1.86	13 (18%)
4	MES	B	336	-	12,12,12	0.92	1 (8%)	15,16,16	0.95	0
2	NAP	A	334	-	50,52,52	1.72	11 (22%)	71,80,80	1.88	14 (19%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	UD1	A	335	-	-	11/26/63/63	0/3/3/3
3	UD1	B	335	-	-	8/26/63/63	0/3/3/3
2	NAP	B	334	-	-	2/35/67/67	0/5/5/5
4	MES	B	336	-	-	2/6/14/14	0/1/1/1
2	NAP	A	334	-	-	3/35/67/67	0/5/5/5

The worst 5 of 31 bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	A	334	NAP	C2N-N1N	4.66	1.40	1.35
2	B	334	NAP	C2N-N1N	4.30	1.39	1.35
2	B	334	NAP	P2B-O1X	3.74	1.62	1.50
2	A	334	NAP	C2A-N1A	3.48	1.40	1.33
2	B	334	NAP	C2A-N1A	3.42	1.40	1.33

The worst 5 of 33 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	A	334	NAP	C4A-N9A-C8A	7.27	113.37	105.74
2	B	334	NAP	C4A-N9A-C8A	7.01	113.10	105.74
2	A	334	NAP	C6N-N1N-C2N	6.92	127.76	121.88
2	B	334	NAP	C6N-N1N-C2N	6.75	127.61	121.88
3	B	335	UD1	O5'-C1'-O1'	-6.69	102.61	111.36

There are no chirality outliers.

5 of 26 torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
3	A	335	UD1	C8'-C7'-N2'-C2'
3	A	335	UD1	O7'-C7'-N2'-C2'
3	A	335	UD1	C1'-O1'-PB-O3A
3	A	335	UD1	O4B-C4B-C5B-O5B
3	A	335	UD1	C5B-O5B-PA-O1A


There are no ring outliers.


3 monomers are involved in 8 short contacts:


Mol	Chain	Res	Type	Clashes	Symm-Clashes
3	A	335	UD1	1	0
2	B	334	NAP	3	0
2	A	334	NAP	4	0


The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier.

The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.









5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	329/344 (95%)	0.70	60 (18%) 3 3	5, 26, 80, 80	0
1	B	327/344 (95%)	0.32	34 (10%) 11 10	5, 22, 76, 80	0
All	All	656/688 (95%)	0.51	94 (14%) 6 5	5, 24, 80, 80	0

The worst 5 of 94 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	196	ILE	8.8
1	A	247	ALA	7.4
1	A	248	PRO	6.5
1	A	245	ALA	6.3
1	A	246	LEU	6.1

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)


There are no oligosaccharides in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled ‘Q< 0.9’ lists the number of atoms with occupancy less than 0.9.




Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors($\text{\AA}^2$)	Q<0.9
4	MES	B	336	12/12	0.70	0.33	78,80,80,80	0
3	UD1	A	335	39/39	0.79	0.20	75,79,80,80	14
3	UD1	B	335	39/39	0.85	0.16	56,61,72,72	14
2	NAP	B	334	48/48	0.96	0.10	13,24,49,51	0
2	NAP	A	334	48/48	0.97	0.09	7,22,40,42	0

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.







Electron density around UD1 B 335:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)

**Electron density around NAP B 334:**

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)





6.5 Other polymers [i](#)

There are no such residues in this entry.