



wwPDB X-ray Structure Validation Summary Report ⓘ

Aug 5, 2026 – 03:12 PM EDT

PDB ID : 3H3X / pdb_00003h3x
Title : Structure of the V74M large subunit mutant of NI-FE hydrogenase in an oxidized state
Authors : Volbeda, A.; Martinez, N.; Martin, L.; Fontecilla-Camps, J.C.
Deposited on : 2009-04-17
Resolution : 2.70 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4-5-2 with Phenix2.0
Mogul	:	2022.3.0, CSD as543be (2022)
Xtriage (Phenix)	:	2.0
EDS	:	3.0
Percentile statistics	:	20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4	:	9.0.015 (Gargrove)
Density-Fitness	:	1.0.12
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.50


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.70 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	3538 (2.70-2.70)
Clashscore	190562	3843 (2.70-2.70)
Ramachandran outliers	187476	3778 (2.70-2.70)
Sidechain outliers	187428	3778 (2.70-2.70)
RSRZ outliers	180081	3538 (2.70-2.70)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	264	
1	B	264	
1	C	264	
2	Q	549	
2	R	549	

Continued on next page...

Continued from previous page...

Mol	Chain	Length	Quality of chain
2	S	549	 A horizontal bar chart showing the quality of the chain. The bar is divided into two segments: a green segment representing 85% and a yellow segment representing 14%. The bar ends with a small black dot.

2 Entry composition

There are 9 unique types of molecules in this entry. The entry contains 19100 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Periplasmic [NiFe] hydrogenase small subunit.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	262	Total	C	N	O	S	0	0	0
			1973	1256	330	372	15			
1	B	262	Total	C	N	O	S	0	0	0
			1974	1257	330	372	15			
1	C	260	Total	C	N	O	S	0	0	0
			1960	1248	327	370	15			


- Molecule 2 is a protein called Periplasmic [NiFe] hydrogenase large subunit.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
2	Q	544	Total	C	N	O	S	0	0	0
			4163	2652	723	765	23			
2	R	545	Total	C	N	O	S	0	0	0
			4167	2654	724	766	23			
2	S	544	Total	C	N	O	S	0	0	0
			4163	2652	723	765	23			

There are 3 discrepancies between the modelled and reference sequences:


Chain	Residue	Modelled	Actual	Comment	Reference
Q	74	MET	VAL	engineered mutation	UNP P18188
R	74	MET	VAL	engineered mutation	UNP P18188
S	74	MET	VAL	engineered mutation	UNP P18188

- Molecule 3 is IRON/SULFUR CLUSTER (CCD ID: SF4) (formula: Fe₄S₄).




Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
3	A	1	Total	Fe	S	0	0
			8	4	4		
3	A	1	Total	Fe	S	0	0
			8	4	4		
3	B	1	Total	Fe	S	0	0
			8	4	4		
3	B	1	Total	Fe	S	0	0
			8	4	4		
3	C	1	Total	Fe	S	0	0
			8	4	4		
3	C	1	Total	Fe	S	0	0
			8	4	4		

- Molecule 4 is FE3-S4 CLUSTER (CCD ID: F3S) (formula: Fe_3S_4).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
4	A	1	Total	Fe	S	0	0
			7	3	4		
4	B	1	Total	Fe	S	0	0
			7	3	4		
4	C	1	Total	Fe	S	0	0
			7	3	4		

- Molecule 5 is GLYCEROL (CCD ID: GOL) (formula: C₃H₈O₃).




Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
5	A	1	Total	C	O	0	0
			6	3	3		

Continued on next page...

Continued from previous page...

Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
5	Q	1	Total	C	O	0	0
			6	3	3		
5	Q	1	Total	C	O	0	0
			6	3	3		
5	R	1	Total	C	O	0	0
			6	3	3		
5	R	1	Total	C	O	0	0
			6	3	3		
5	R	1	Total	C	O	0	0
			6	3	3		
5	S	1	Total	C	O	0	0
			6	3	3		

- Molecule 6 is CARBONMONOXIDE-(DICYANO) IRON (CCD ID: FCO) (formula: $\text{C}_3\text{FeN}_2\text{O}$).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
6	Q	1	Total	C	Fe	N	O	0	0
			7	3	1	2	1		
6	R	1	Total	C	Fe	N	O	0	0
			7	3	1	2	1		
6	S	1	Total	C	Fe	N	O	0	0
			7	3	1	2	1		

- Molecule 7 is NICKEL (II) ION (CCD ID: NI) (formula: Ni).

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
7	Q	1	Total Ni 1 1	0	0
7	R	1	Total Ni 1 1	0	0
7	S	1	Total Ni 1 1	0	0

- Molecule 8 is MAGNESIUM ION (CCD ID: MG) (formula: Mg).

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
8	Q	1	Total Mg 1 1	0	0
8	R	1	Total Mg 1 1	0	0
8	S	1	Total Mg 1 1	0	0


- Molecule 9 is water.

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
9	A	68	Total O 68 68	0	0
9	Q	144	Total O 144 144	0	0
9	B	58	Total O 58 58	0	0
9	R	133	Total O 133 133	0	0
9	C	65	Total O 65 65	0	0
9	S	94	Total O 94 94	0	0


3 Residue-property plots

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: Periplasmic [NiFe] hydrogenase small subunit




- Molecule 1: Periplasmic [NiFe] hydrogenase small subunit



- Molecule 1: Periplasmic [NiFe] hydrogenase small subunit




- Molecule 2: Periplasmic [NiFe] hydrogenase large subunit






• Molecule 2: Periplasmic [NiFe] hydrogenase large subunit

Chain R: 86% 13%



• Molecule 2: Periplasmic [NiFe] hydrogenase large subunit

Chain S: 85% 14%



4 Data and refinement statistics

Property	Value	Source
Space group	P 1 21 1	Depositor
Cell constants a, b, c, α , β , γ	62.40Å 99.20Å 182.30Å 90.00° 92.80° 90.00°	Depositor
Resolution (Å)	20.00 – 2.70 20.00 – 2.70	Depositor EDS
% Data completeness (in resolution range)	98.7 (20.00-2.70) 85.5 (20.00-2.70)	Depositor EDS
R_{merge}	0.05	Depositor
R_{sym}	0.05	Depositor
$\langle I/\sigma(I) \rangle$ ¹	4.78 (at 2.71Å)	Xtriage
Refinement program	REFMAC 5.2.0019	Depositor
R, R_{free}	0.203 , 0.247 0.205 , 0.247	Depositor DCC
R_{free} test set	2935 reflections (4.87%)	wwPDB-VP
Wilson B-factor (Å ²)	38.9	Xtriage
Anisotropy	0.249	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.37 , 46.8	EDS
L-test for twinning ²	$\langle L \rangle = 0.40$, $\langle L^2 \rangle = 0.22$	Xtriage
Estimated twinning fraction	0.065 for h,-k,-l	Xtriage
F_o, F_c correlation	0.94	EDS
Total number of atoms	19100	wwPDB-VP
Average B, all atoms (Å ²)	36.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 3.58% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: GOL, MG, FCO, F3S, NI, SF4

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.91	1/2027 (0.0%)	0.99	2/2759 (0.1%)
1	B	0.75	0/2028	0.91	1/2761 (0.0%)
1	C	0.85	0/2014	0.95	3/2743 (0.1%)
2	Q	0.92	2/4268 (0.0%)	1.00	3/5792 (0.1%)
2	R	0.80	0/4272	0.97	3/5798 (0.1%)
2	S	0.75	0/4268	0.93	3/5792 (0.1%)
All	All	0.83	3/18877 (0.0%)	0.96	15/25645 (0.1%)

All (3) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	Q	267	ILE	CA-CB	8.34	1.58	1.54
2	Q	515	VAL	CA-CB	6.34	1.62	1.54
1	A	78	LEU	C-O	-5.23	1.20	1.25

The worst 5 of 15 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	Q	267	ILE	N-CA-CB	7.10	115.99	110.45
2	S	462	MET	CA-C-N	-5.52	114.65	120.66
2	S	462	MET	C-N-CA	-5.52	114.65	120.66
1	C	120	VAL	N-CA-C	5.49	116.22	110.62
2	S	239	ASN	N-CA-C	5.48	118.10	109.39

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts ⓘ

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1973	0	1911	21	0
1	B	1974	0	1915	17	0
1	C	1960	0	1897	27	0
2	Q	4163	0	4136	42	0
2	R	4167	0	4136	46	0
2	S	4163	0	4136	44	0
3	A	16	0	0	0	0
3	B	16	0	0	0	0
3	C	16	0	0	0	0
4	A	7	0	0	0	0
4	B	7	0	0	0	0
4	C	7	0	0	0	0
5	A	6	0	8	0	0
5	Q	12	0	16	2	0
5	R	18	0	24	2	0
5	S	6	0	8	2	0
6	Q	7	0	0	1	0
6	R	7	0	0	0	0
6	S	7	0	0	0	0
7	Q	1	0	0	0	0
7	R	1	0	0	0	0
7	S	1	0	0	0	0
8	Q	1	0	0	0	0
8	R	1	0	0	0	0
8	S	1	0	0	0	0
9	A	68	0	0	3	0
9	B	58	0	0	4	0
9	C	65	0	0	3	0
9	Q	144	0	0	1	0
9	R	133	0	0	5	0
9	S	94	0	0	3	0
All	All	19100	0	18187	185	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 5.

The worst 5 of 185 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
2:S:104:ASN:HD22	5:S:561:GOL:H32	1.40	0.86
2:Q:89:ASP:O	2:Q:92:LYS:HE2	1.86	0.75
2:Q:320:ASP:HB2	2:R:6:PRO:HD3	1.70	0.74
2:Q:185:LEU:HB2	5:Q:562:GOL:H12	1.72	0.69
2:S:319:ARG:HG2	2:S:419:HIS:CE1	2.27	0.69

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	260/264 (98%)	253 (97%)	6 (2%)	1 (0%)	30	54
1	B	260/264 (98%)	252 (97%)	7 (3%)	1 (0%)	30	54
1	C	258/264 (98%)	250 (97%)	7 (3%)	1 (0%)	30	54
2	Q	542/549 (99%)	522 (96%)	20 (4%)	0	100	100
2	R	543/549 (99%)	519 (96%)	23 (4%)	1 (0%)	43	68
2	S	542/549 (99%)	520 (96%)	22 (4%)	0	100	100
All	All	2405/2439 (99%)	2316 (96%)	85 (4%)	4 (0%)	43	68

All (4) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	B	231	LEU
1	C	231	LEU
1	A	231	LEU
2	R	6	PRO

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	208/210 (99%)	206 (99%)	2 (1%)	68	86
1	B	208/210 (99%)	203 (98%)	5 (2%)	43	72
1	C	207/210 (99%)	203 (98%)	4 (2%)	50	77
2	Q	434/439 (99%)	424 (98%)	10 (2%)	44	73
2	R	434/439 (99%)	424 (98%)	10 (2%)	44	73
2	S	434/439 (99%)	429 (99%)	5 (1%)	63	84
All	All	1925/1947 (99%)	1889 (98%)	36 (2%)	50	77

5 of 36 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	C	95	ILE
2	S	546	CYS
1	C	135	GLU
2	S	454	ASN
1	B	4	LYS

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 15 such sidechains are listed below:

Mol	Chain	Res	Type
1	B	14	ASN
2	S	104	ASN
1	B	234	GLN
2	S	454	ASN
1	C	14	ASN

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates ⓘ

There are no oligosaccharides in this entry.

5.6 Ligand geometry ⓘ

Of 25 ligands modelled in this entry, 6 are monoatomic - leaving 19 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
3	SF4	A	265	1	0,12,12	-	-	-		
3	SF4	B	265	1	0,12,12	-	-	-		
5	GOL	Q	562	-	5,5,5	0.42	0	5,5,5	0.95	0
3	SF4	C	265	1	0,12,12	-	-	-		
3	SF4	A	267	1	0,12,12	-	-	-		
6	FCO	Q	550	2	3,6,6	2.60	1 (33%)	-		
5	GOL	Q	561	-	5,5,5	0.35	0	5,5,5	1.19	0
5	GOL	R	562	-	5,5,5	0.52	0	5,5,5	0.56	0
5	GOL	S	561	-	5,5,5	0.55	0	5,5,5	0.45	0
4	F3S	C	266	1	0,9,9	-	-	-		
5	GOL	A	271	-	5,5,5	0.30	0	5,5,5	0.60	0
6	FCO	R	550	2	3,6,6	1.31	0	-		
5	GOL	R	561	-	5,5,5	0.52	0	5,5,5	0.26	0
3	SF4	C	267	1	0,12,12	-	-	-		
3	SF4	B	267	1	0,12,12	-	-	-		
6	FCO	S	550	2	3,6,6	2.28	1 (33%)	-		
4	F3S	B	266	1	0,9,9	-	-	-		
5	GOL	R	563	-	5,5,5	0.39	0	5,5,5	0.39	0
4	F3S	A	266	1	0,9,9	-	-	-		

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the

Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns.
'-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	SF4	A	265	1	-	-	0/6/5/5
5	GOL	Q	561	-	-	0/4/4/4	-
5	GOL	Q	562	-	-	1/4/4/4	-
3	SF4	B	265	1	-	-	0/6/5/5
4	F3S	B	266	1	-	-	0/3/3/3
5	GOL	R	562	-	-	4/4/4/4	-
5	GOL	A	271	-	-	0/4/4/4	-
4	F3S	C	266	1	-	-	0/3/3/3
3	SF4	C	265	1	-	-	0/6/5/5
3	SF4	C	267	1	-	-	0/6/5/5
3	SF4	B	267	1	-	-	0/6/5/5
5	GOL	S	561	-	-	2/4/4/4	-
3	SF4	A	267	1	-	-	0/6/5/5
5	GOL	R	561	-	-	4/4/4/4	-
5	GOL	R	563	-	-	0/4/4/4	-
4	F3S	A	266	1	-	-	0/3/3/3

All (2) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
6	Q	550	FCO	O3-C3	-3.72	1.11	1.16
6	S	550	FCO	C2-N2	3.47	1.20	1.15

There are no bond angle outliers.

There are no chirality outliers.

5 of 11 torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
5	R	561	GOL	O1-C1-C2-C3
5	R	561	GOL	C1-C2-C3-O3
5	Q	562	GOL	O1-C1-C2-C3
5	R	562	GOL	O1-C1-C2-C3
5	R	562	GOL	C1-C2-C3-O3

There are no ring outliers.

4 monomers are involved in 7 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
5	Q	562	GOL	2	0
6	Q	550	FCO	1	0
5	S	561	GOL	2	0
5	R	561	GOL	2	0

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	262/264 (99%)	-1.60	0 100 100	29, 36, 38, 46	8 (3%)
1	B	262/264 (99%)	-1.45	0 100 100	29, 36, 37, 41	10 (3%)
1	C	260/264 (98%)	-1.62	0 100 100	27, 36, 38, 42	6 (2%)
2	Q	544/549 (99%)	-1.62	0 100 100	28, 36, 39, 42	15 (2%)
2	R	545/549 (99%)	-1.54	0 100 100	26, 36, 38, 41	15 (2%)
2	S	544/549 (99%)	-1.50	0 100 100	26, 36, 38, 41	14 (2%)
All	All	2417/2439 (99%)	-1.55	0 100 100	26, 36, 38, 46	68 (2%)

There are no RSRZ outliers to report.

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled ‘Q< 0.9’ lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
5	GOL	A	271	6/6	0.98	0.07	34,34,35,36	0

Continued on next page...

Continued from previous page...

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors($\text{\AA}^2$)	Q<0.9
5	GOL	R	561	6/6	0.98	0.07	46,49,49,50	0
5	GOL	R	563	6/6	0.98	0.06	45,47,49,50	0
5	GOL	S	561	6/6	0.98	0.05	45,48,49,50	0
5	GOL	Q	561	6/6	0.99	0.03	32,34,35,38	0
5	GOL	Q	562	6/6	0.99	0.03	35,36,37,39	0
3	SF4	B	267	8/8	0.99	0.03	35,35,36,37	0
5	GOL	R	562	6/6	0.99	0.04	44,45,46,46	0
4	F3S	B	266	7/7	0.99	0.05	36,38,39,42	0
3	SF4	B	265	8/8	0.99	0.06	30,32,33,34	0
7	NI	Q	551	1/1	0.99	0.01	39,39,39,39	0
7	NI	S	551	1/1	0.99	0.02	39,39,39,39	0
4	F3S	A	266	7/7	1.00	0.01	32,34,34,35	0
3	SF4	A	265	8/8	1.00	0.02	29,32,34,35	0
4	F3S	C	266	7/7	1.00	0.03	35,36,37,40	0
3	SF4	A	267	8/8	1.00	0.02	36,37,39,39	0
6	FCO	Q	550	7/7	1.00	0.03	34,35,37,37	0
6	FCO	R	550	7/7	1.00	0.03	36,36,40,41	0
6	FCO	S	550	7/7	1.00	0.05	35,37,40,40	0
3	SF4	C	265	8/8	1.00	0.01	33,34,35,36	0
7	NI	R	551	1/1	1.00	0.01	41,41,41,41	0
3	SF4	C	267	8/8	1.00	0.02	36,37,37,38	0
8	MG	Q	553	1/1	1.00	0.02	35,35,35,35	0
8	MG	R	553	1/1	1.00	0.04	34,34,34,34	0
8	MG	S	553	1/1	1.00	0.02	33,33,33,33	0

6.5 Other polymers [i](#)

There are no such residues in this entry.