



wwPDB X-ray Structure Validation Summary Report ⓘ

Aug 5, 2026 – 03:08 AM JST

PDB ID : 5HPW / pdb_00005hpw
Title : Mode of binding of antithyroid drug, propylthiouracil to lactoperoxidase: Binding studies and structure determination
Authors : Singh, R.P.; Singh, A.; Sharma, P.; Kaur, P.; Sharma, S.; Singh, T.P.
Deposited on : 2016-01-21
Resolution : 2.50 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4-5-2 with Phenix2.0
Mogul : 1.8.5 (274361), CSD as541be (2020)
Xtriage (Phenix) : 2.0
EDS : 3.0
Buster-report : wwPDB partial adaption of 1.1.7 (2018)
Percentile statistics : 20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4 : 9.0.015 (Gargrove)
Density-Fitness : 1.0.12
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.50


1 Overall quality at a glance ⓘ

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.50 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.




Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	5829 (2.50-2.50)
Clashscore	190562	6492 (2.50-2.50)
Ramachandran outliers	187476	6378 (2.50-2.50)
Sidechain outliers	187428	6380 (2.50-2.50)
RSRZ outliers	180081	5833 (2.50-2.50)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	595	4% 44% 47% 8%
1	B	595	3% 48% 44% 7%
1	C	595	5% 47% 43% 9%
1	D	595	5% 39% 49% 11%
2	E	2	50% 50%
2	G	2	50% 50%

Continued on next page...

Continued from previous page...

Mol	Chain	Length	Quality of chain
2	H	2	 100%
3	F	2	 100%

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
6	NO3	B	603	-	-	X	-
6	NO3	C	606	-	-	X	-
6	NO3	D	604	-	-	X	-
6	NO3	D	605	-	-	X	-
6	NO3	D	606	-	-	X	-
7	3CJ	A	607	-	X	X	-
7	3CJ	B	607	-	-	X	-
7	3CJ	C	607	-	-	X	-
7	3CJ	D	607	-	X	X	-

2 Entry composition [i](#)

There are 9 unique types of molecules in this entry. The entry contains 20141 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Lactoperoxidase.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	595	Total	C	N	O	S	0	0	0
			4753	3021	844	862	26			
1	B	595	Total	C	N	O	S	0	0	0
			4753	3021	844	862	26			
1	C	595	Total	C	N	O	S	0	0	0
			4753	3021	844	862	26			
1	D	595	Total	C	N	O	S	0	0	0
			4753	3021	844	862	26			

- Molecule 2 is an oligosaccharide called 2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose.




Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	Trace
2	E	2	Total	C	N	O	0	0	0
			28	16	2	10			
2	G	2	Total	C	N	O	0	0	0
			28	16	2	10			
2	H	2	Total	C	N	O	0	0	0
			28	16	2	10			

- Molecule 3 is an oligosaccharide called 2-acetamido-2-deoxy-beta-D-glucopyranose-(1-3)-2-acetamido-2-deoxy-beta-D-glucopyranose.



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	Trace
3	F	2	Total	C	N	O	0	0	0
			28	16	2	10			

- Molecule 4 is 2-acetamido-2-deoxy-beta-D-glucopyranose (CCD ID: NAG) (formula: $C_8H_{15}NO_6$).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
4	A	1	Total	C	N	O	0	0
			14	8	1	5		
4	A	1	Total	C	N	O	0	0
			14	8	1	5		
4	B	1	Total	C	N	O	0	0
			14	8	1	5		
4	C	1	Total	C	N	O	0	0
			14	8	1	5		
4	C	1	Total	C	N	O	0	0
			14	8	1	5		
4	D	1	Total	C	N	O	0	0
			14	8	1	5		
4	D	1	Total	C	N	O	0	0
			14	8	1	5		

- Molecule 5 is CALCIUM ION (CCD ID: CA) (formula: Ca).


Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
5	A	1	Total	Ca	0	0
			1	1		

Continued on next page...

Continued from previous page...

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
5	B	1	Total	Ca	0	0
			1	1		
5	C	1	Total	Ca	0	0
			1	1		
5	D	1	Total	Ca	0	0
			1	1		

- Molecule 6 is NITRATE ION (CCD ID: NO3) (formula: NO₃).




Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
6	A	1	Total	N	O	0	0
			4	1	3		
6	A	1	Total	N	O	0	0
			4	1	3		
6	A	1	Total	N	O	0	0
			4	1	3		
6	B	1	Total	N	O	0	0
			4	1	3		
6	B	1	Total	N	O	0	0
			4	1	3		
6	B	1	Total	N	O	0	0
			4	1	3		
6	C	1	Total	N	O	0	0
			4	1	3		
6	C	1	Total	N	O	0	0
			4	1	3		

Continued on next page...

Continued from previous page...


Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
6	C	1	Total	N	O	0	0
			4	1	3		
6	D	1	Total	N	O	0	0
			4	1	3		
6	D	1	Total	N	O	0	0
			4	1	3		
6	D	1	Total	N	O	0	0
			4	1	3		

- Molecule 7 is 6-propyl-2-thioxo-2,3-dihydropyrimidin-4(1H)-one (CCD ID: 3CJ) (formula: C₇H₁₀N₂OS).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
7	A	1	Total	C	N	O	S	0	0
			11	7	2	1	1		
7	B	1	Total	C	N	O	S	0	0
			11	7	2	1	1		
7	C	1	Total	C	N	O	S	0	0
			11	7	2	1	1		
7	D	1	Total	C	N	O	S	0	0
			11	7	2	1	1		

- Molecule 8 is PROTOPORPHYRIN IX CONTAINING FE (CCD ID: HEM) (formula: C₃₄H₃₂FeN₄O₄).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
8	A	1	Total 43	C 34	Fe 1	N 4	O 4	0	0
8	B	1	Total 43	C 34	Fe 1	N 4	O 4	0	0
8	C	1	Total 43	C 34	Fe 1	N 4	O 4	0	0
8	D	1	Total 43	C 34	Fe 1	N 4	O 4	0	0


- Molecule 9 is water.

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
9	A	158	Total O 158 158	0	0
9	B	157	Total O 157 157	0	0
9	C	168	Total O 168 168	0	0
9	D	168	Total O 168 168	0	0


3 Residue-property plots


These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


• Molecule 1: Lactoperoxidase




• Molecule 1: Lactoperoxidase







• Molecule 1: Lactoperoxidase



• Molecule 2: 2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose



- Molecule 2: 2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose

Chain G:  50% 50%

MAG1
MAG2

- Molecule 2: 2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose

Chain H:  100%

MAG1
MAG2

- Molecule 3: 2-acetamido-2-deoxy-beta-D-glucopyranose-(1-3)-2-acetamido-2-deoxy-beta-D-glucopyranose

Chain F:  100%

MAG1
MAG2

4 Data and refinement statistics

Property	Value	Source
Space group	P 1	Depositor
Cell constants a, b, c, α , β , γ	80.22Å 82.59Å 95.08Å 80.91° 73.71° 89.96°	Depositor
Resolution (Å)	42.50 – 2.50 42.50 – 2.50	Depositor EDS
% Data completeness (in resolution range)	91.7 (42.50-2.50) 91.7 (42.50-2.50)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	2.19 (at 2.51Å)	Xtriage
Refinement program	REFMAC 5.7.0032	Depositor
R, R_{free}	0.260 , 0.311 0.254 , 0.307	Depositor DCC
R_{free} test set	3683 reflections (4.63%)	wwPDB-VP
Wilson B-factor (Å ²)	41.2	Xtriage
Anisotropy	0.744	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.24 , 43.0	EDS
L-test for twinning ²	$\langle L \rangle = 0.47$, $\langle L^2 \rangle = 0.30$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.91	EDS
Total number of atoms	20141	wwPDB-VP
Average B, all atoms (Å ²)	45.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The analyses of the Patterson function reveals a significant off-origin peak that is 40.53 % of the origin peak, indicating pseudo-translational symmetry. The chance of finding a peak of this or larger height randomly in a structure without pseudo-translational symmetry is equal to 2.7003e-04. The detected translational NCS is most likely also responsible for the elevated intensity ratio.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: NAG, HEM, NO3, 3CJ, CA

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.68	5/4882 (0.1%)	1.22	59/6632 (0.9%)
1	B	0.62	2/4882 (0.0%)	1.19	48/6632 (0.7%)
1	C	0.68	6/4882 (0.1%)	1.19	56/6632 (0.8%)
1	D	0.63	1/4882 (0.0%)	1.23	66/6632 (1.0%)
All	All	0.66	14/19528 (0.1%)	1.21	229/26528 (0.9%)

The worst 5 of 14 bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	10	VAL	C-N	14.32	1.52	1.34
1	C	31	ARG	CA-C	-11.34	1.38	1.52
1	C	31	ARG	C-N	-8.94	1.22	1.33
1	C	240	ILE	C-N	7.17	1.43	1.33
1	B	107	VAL	C-O	-6.59	1.16	1.24

The worst 5 of 229 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	349	PHE	N-CA-C	-13.77	96.49	113.28
1	D	4	VAL	N-CA-C	-10.88	102.45	111.81
1	D	111	LEU	N-CA-C	10.67	122.49	111.07
1	A	124	HIS	N-CA-C	-10.53	100.54	113.50
1	A	492	ILE	N-CA-C	10.44	121.32	110.36

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts ⓘ

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	4753	0	4646	309	0
1	B	4753	0	4646	246	0
1	C	4753	0	4649	327	0
1	D	4753	0	4647	312	0
2	E	28	0	25	1	0
2	G	28	0	25	1	0
2	H	28	0	25	0	0
3	F	28	0	25	2	0
4	A	28	0	26	0	0
4	B	14	0	13	1	0
4	C	28	0	26	0	0
4	D	28	0	26	1	0
5	A	1	0	0	0	0
5	B	1	0	0	0	0
5	C	1	0	0	0	0
5	D	1	0	0	0	0
6	A	12	0	0	1	0
6	B	12	0	0	2	0
6	C	12	0	0	4	0
6	D	12	0	0	8	0
7	A	11	0	10	11	0
7	B	11	0	10	12	0
7	C	11	0	10	9	0
7	D	11	0	10	11	0
8	A	43	0	30	12	0
8	B	43	0	30	14	0
8	C	43	0	30	17	0
8	D	43	0	30	15	0
9	A	158	0	0	15	0
9	B	157	0	0	10	0
9	C	168	0	0	20	0
9	D	168	0	0	11	0
All	All	20141	0	18939	1217	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 32.

The worst 5 of 1217 close contacts within the same asymmetric unit are listed below, sorted by

their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:D:258:GLU:OE2	8:D:608:HEM:CMB	1.65	1.42
1:A:167:CYS:HB2	1:A:168:PRO:CD	1.55	1.32
1:D:258:GLU:OE2	8:D:608:HEM:HMB1	1.18	1.30
1:B:10:VAL:HG11	1:B:41:ARG:NH1	1.48	1.28
1:C:96:ARG:HD2	1:C:100:PHE:CD2	1.73	1.21

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	593/595 (100%)	520 (88%)	68 (12%)	5 (1%)	16	31
1	B	593/595 (100%)	524 (88%)	63 (11%)	6 (1%)	12	24
1	C	593/595 (100%)	523 (88%)	61 (10%)	9 (2%)	8	16
1	D	593/595 (100%)	515 (87%)	67 (11%)	11 (2%)	6	11
All	All	2372/2380 (100%)	2082 (88%)	259 (11%)	31 (1%)	9	18

5 of 31 Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	8	ALA
1	A	167	CYS
1	B	167	CYS
1	C	8	ALA
1	C	12	LEU

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	517/517 (100%)	469 (91%)	48 (9%)	8	18
1	B	517/517 (100%)	468 (90%)	49 (10%)	8	17
1	C	517/517 (100%)	467 (90%)	50 (10%)	8	17
1	D	517/517 (100%)	456 (88%)	61 (12%)	5	11
All	All	2068/2068 (100%)	1860 (90%)	208 (10%)	7	15

5 of 208 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	C	146	LYS
1	C	578	ASP
1	D	538	GLU
1	C	177	ARG
1	C	333	ASN

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 57 such sidechains are listed below:

Mol	Chain	Res	Type
1	C	94	GLN
1	D	403	ASN
1	C	266	HIS
1	D	333	ASN
1	D	128	GLN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates ⓘ

8 monosaccharides are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# $ Z > 2$	Counts	RMSZ	# $ Z > 2$
2	NAG	E	1	1,2	14,14,15	1.33	3 (21%)	17,19,21	1.45	1 (5%)
2	NAG	E	2	2	14,14,15	0.92	1 (7%)	17,19,21	2.78	6 (35%)
3	NAG	F	1	1,3	14,14,15	1.44	2 (14%)	17,19,21	2.31	6 (35%)
3	NAG	F	2	3	14,14,15	0.99	0	17,19,21	1.99	4 (23%)
2	NAG	G	1	1,2	14,14,15	0.27	0	17,19,21	0.60	0
2	NAG	G	2	2	14,14,15	0.29	0	17,19,21	0.61	0
2	NAG	H	1	1,2	14,14,15	0.84	0	17,19,21	1.49	3 (17%)
2	NAG	H	2	2	14,14,15	0.72	0	17,19,21	1.99	5 (29%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	NAG	E	1	1,2	-	5/6/23/26	0/1/1/1
2	NAG	E	2	2	-	4/6/23/26	0/1/1/1
3	NAG	F	1	1,3	-	2/6/23/26	0/1/1/1
3	NAG	F	2	3	-	6/6/23/26	0/1/1/1
2	NAG	G	1	1,2	-	0/6/23/26	0/1/1/1
2	NAG	G	2	2	-	0/6/23/26	0/1/1/1
2	NAG	H	1	1,2	-	6/6/23/26	0/1/1/1
2	NAG	H	2	2	-	4/6/23/26	0/1/1/1

The worst 5 of 6 bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
3	F	1	NAG	C1-C2	2.83	1.56	1.52

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	E	1	NAG	C2-N2	-2.81	1.41	1.46
2	E	1	NAG	O5-C1	-2.70	1.39	1.43
2	E	2	NAG	C2-N2	-2.50	1.42	1.46
3	F	1	NAG	O5-C1	-2.07	1.40	1.43

The worst 5 of 25 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	E	2	NAG	C1-O5-C5	8.57	123.80	112.19
3	F	1	NAG	O5-C5-C6	4.93	114.93	107.20
3	F	2	NAG	C4-C3-C2	-4.75	104.06	111.02
3	F	1	NAG	O3-C3-C2	4.74	119.27	109.47
2	E	1	NAG	C2-N2-C7	-4.35	116.70	122.90

There are no chirality outliers.

5 of 27 torsion outliers are listed below:


Mol	Chain	Res	Type	Atoms
2	E	1	NAG	C3-C2-N2-C7
2	E	1	NAG	O7-C7-N2-C2
2	H	1	NAG	C3-C2-N2-C7
3	F	1	NAG	C3-C2-N2-C7
3	F	2	NAG	C8-C7-N2-C2


There are no ring outliers.


4 monomers are involved in 4 short contacts:


Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	E	2	NAG	1	0
3	F	2	NAG	2	0
2	G	1	NAG	1	0
3	F	1	NAG	2	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for oligosaccharide.









5.6 Ligand geometry [i](#)

Of 31 ligands modelled in this entry, 4 are monoatomic - leaving 27 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
6	NO3	D	606	-	1,3,3	0.35	0	0,3,3	-	-
4	NAG	A	602	1	14,14,15	0.29	0	17,19,21	0.61	0
6	NO3	C	605	-	1,3,3	3.46	1 (100%)	0,3,3	-	-
4	NAG	C	601	1	14,14,15	0.30	0	17,19,21	0.62	0
6	NO3	C	604	-	1,3,3	0.34	0	0,3,3	-	-
4	NAG	B	601	1	14,14,15	1.18	1 (7%)	17,19,21	1.81	7 (41%)
8	HEM	B	606	1	50,50,50	1.92	15 (30%)	66,82,82	1.83	17 (25%)
8	HEM	D	608	1	50,50,50	1.82	17 (34%)	66,82,82	1.80	14 (21%)
4	NAG	A	601	1	14,14,15	1.01	0	17,19,21	2.38	8 (47%)
6	NO3	A	604	-	1,3,3	1.27	0	0,3,3	-	-
8	HEM	A	608	1,7	50,50,50	1.86	15 (30%)	66,82,82	1.88	12 (18%)
8	HEM	C	608	1	50,50,50	1.68	12 (24%)	66,82,82	1.91	19 (28%)
6	NO3	D	605	-	1,3,3	3.56	1 (100%)	0,3,3	-	-
6	NO3	B	604	-	1,3,3	3.57	1 (100%)	0,3,3	-	-
6	NO3	D	604	-	1,3,3	0.34	0	0,3,3	-	-
6	NO3	B	603	-	1,3,3	0.35	0	0,3,3	-	-
6	NO3	B	605	-	1,3,3	0.46	0	0,3,3	-	-
4	NAG	D	602	1	14,14,15	0.28	0	17,19,21	0.63	0
7	3CJ	C	607	-	11,11,11	2.32	2 (18%)	14,14,14	3.26	8 (57%)
7	3CJ	B	607	-	11,11,11	2.25	2 (18%)	14,14,14	3.15	7 (50%)
6	NO3	A	605	-	1,3,3	3.55	1 (100%)	0,3,3	-	-
6	NO3	C	606	-	1,3,3	0.32	0	0,3,3	-	-
4	NAG	C	602	1	14,14,15	0.29	0	17,19,21	0.62	0
7	3CJ	A	607	8	11,11,11	2.06	2 (18%)	14,14,14	3.22	9 (64%)
4	NAG	D	601	1	14,14,15	0.81	0	17,19,21	1.94	5 (29%)
6	NO3	A	606	-	1,3,3	0.34	0	0,3,3	-	-
7	3CJ	D	607	-	11,11,11	2.70	6 (54%)	14,14,14	3.59	10 (71%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
4	NAG	A	601	1	-	5/6/23/26	0/1/1/1
4	NAG	C	602	1	-	2/6/23/26	0/1/1/1
4	NAG	A	602	1	-	0/6/23/26	0/1/1/1
8	HEM	A	608	1,7	-	4/14/54/54	-

Continued on next page...

Continued from previous page...

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
7	3CJ	D	607	-	-	3/3/3/3	0/1/1/1
7	3CJ	A	607	8	-	1/3/3/3	0/1/1/1
8	HEM	C	608	1	-	4/14/54/54	-
4	NAG	C	601	1	-	4/6/23/26	0/1/1/1
4	NAG	D	601	1	-	4/6/23/26	0/1/1/1
8	HEM	B	606	1	-	4/14/54/54	-
4	NAG	D	602	1	-	0/6/23/26	0/1/1/1
4	NAG	B	601	1	-	2/6/23/26	0/1/1/1
7	3CJ	C	607	-	-	1/3/3/3	0/1/1/1
7	3CJ	B	607	-	-	1/3/3/3	0/1/1/1
8	HEM	D	608	1	-	2/14/54/54	-

The worst 5 of 76 bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
7	C	607	3CJ	O1-C4	6.80	1.37	1.24
7	D	607	3CJ	O1-C4	6.78	1.37	1.24
7	B	607	3CJ	O1-C4	6.53	1.37	1.24
7	A	607	3CJ	O1-C4	5.89	1.35	1.24
8	A	608	HEM	C4D-ND	-4.90	1.31	1.40

The worst 5 of 116 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
7	D	607	3CJ	C5-C2-N1	7.16	125.94	115.53
7	A	607	3CJ	C1-N2-C4	-6.31	118.94	125.29
7	B	607	3CJ	C1-N2-C4	-5.71	119.54	125.29
8	C	608	HEM	C1B-NB-C4B	5.70	110.96	105.07
7	C	607	3CJ	C1-N2-C4	-5.64	119.62	125.29

There are no chirality outliers.

5 of 37 torsion outliers are listed below:


Mol	Chain	Res	Type	Atoms
4	A	601	NAG	C3-C2-N2-C7
4	C	601	NAG	C8-C7-N2-C2
4	C	601	NAG	O7-C7-N2-C2
7	A	607	3CJ	C2-C5-C6-C7
7	B	607	3CJ	C2-C5-C6-C7


There are no ring outliers.


18 monomers are involved in 92 short contacts:


Mol	Chain	Res	Type	Clashes	Symm-Clashes
6	D	606	NO3	4	0
6	C	605	NO3	1	0
6	C	604	NO3	1	0
4	B	601	NAG	1	0
8	B	606	HEM	14	0
8	D	608	HEM	15	0
8	A	608	HEM	12	0
8	C	608	HEM	17	0
6	D	605	NO3	2	0
6	D	604	NO3	2	0
6	B	603	NO3	2	0
4	D	602	NAG	1	0
7	C	607	3CJ	9	0
7	B	607	3CJ	12	0
6	A	605	NO3	1	0
6	C	606	NO3	2	0
7	A	607	3CJ	11	0
7	D	607	3CJ	11	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.









5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2			OWAB(Å ²)	Q<0.9
1	A	595/595 (100%)	0.40	23 (3%)	43	38	16, 42, 78, 117	0
1	B	595/595 (100%)	0.39	18 (3%)	52	48	20, 43, 76, 100	0
1	C	595/595 (100%)	0.47	32 (5%)	31	27	20, 43, 83, 100	0
1	D	595/595 (100%)	0.45	31 (5%)	33	29	14, 41, 79, 100	0
All	All	2380/2380 (100%)	0.43	104 (4%)	39	34	14, 42, 79, 117	0

The worst 5 of 104 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	D	4	VAL	7.4
1	D	2	TRP	6.1
1	A	171	PRO	6.0
1	B	189	ALA	5.2
1	C	96	ARG	5.1

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled ‘Q< 0.9’ lists the number of atoms with occupancy less than 0.9.


Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
2	NAG	E	2	14/15	0.73	0.14	43,50,51,52	14
2	NAG	G	2	14/15	0.74	0.12	49,56,63,63	14

Continued on next page...

Continued from previous page...




Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors($\text{\AA}^2$)	Q<0.9
2	NAG	H	2	14/15	0.75	0.13	61,67,70,70	14
3	NAG	F	2	14/15	0.75	0.14	45,53,55,58	14
2	NAG	G	1	14/15	0.82	0.14	20,20,20,20	0
3	NAG	F	1	14/15	0.87	0.12	36,50,59,66	0
2	NAG	H	1	14/15	0.88	0.11	70,79,82,85	0
2	NAG	E	1	14/15	0.89	0.10	33,41,52,54	0

The following is a graphical depiction of the model fit to experimental electron density for oligosaccharide. Each fit is shown from different orientation to approximate a three-dimensional view.






Electron density around Chain G:


$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)



Electron density around Chain H:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)





6.4 Ligands ⓘ

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
4	NAG	A	601	14/15	0.55	0.14	64,79,81,81	0
4	NAG	C	601	14/15	0.63	0.17	78,86,89,90	0
6	NO3	C	605	4/4	0.67	0.31	24,24,27,31	0
4	NAG	D	602	14/15	0.75	0.13	46,54,59,60	0
7	3CJ	C	607	11/11	0.75	0.16	69,79,84,85	0
6	NO3	B	605	4/4	0.76	0.37	24,24,24,28	0
6	NO3	B	603	4/4	0.77	0.13	22,25,27,28	0
4	NAG	C	602	14/15	0.77	0.13	48,55,60,61	0
6	NO3	C	606	4/4	0.78	0.22	24,28,29,30	0
4	NAG	D	601	14/15	0.79	0.10	52,60,65,66	0
7	3CJ	B	607	11/11	0.79	0.12	48,52,58,61	0
6	NO3	B	604	4/4	0.79	0.25	26,26,27,29	0
7	3CJ	D	607	11/11	0.79	0.12	39,48,58,58	0

Continued on next page...




Continued from previous page...

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors($\text{\AA}^2$)	Q<0.9
6	NO3	D	605	4/4	0.82	0.27	25,26,27,28	0
6	NO3	C	604	4/4	0.82	0.35	24,27,29,31	0
6	NO3	D	606	4/4	0.85	0.24	23,23,26,30	0
6	NO3	A	604	4/4	0.86	0.21	23,23,24,26	0
4	NAG	B	601	14/15	0.87	0.13	55,63,73,75	0
7	3CJ	A	607	11/11	0.88	0.09	47,50,55,56	0
6	NO3	A	606	4/4	0.88	0.21	23,24,26,27	0
6	NO3	A	605	4/4	0.89	0.15	24,26,28,32	0
4	NAG	A	602	14/15	0.89	0.08	38,46,51,54	0
6	NO3	D	604	4/4	0.89	0.13	21,24,24,26	0
5	CA	C	603	1/1	0.91	0.07	46,46,46,46	0
8	HEM	B	606	43/43	0.93	0.11	33,42,54,62	0
5	CA	B	602	1/1	0.94	0.04	44,44,44,44	0
8	HEM	C	608	43/43	0.94	0.09	35,45,51,53	0
8	HEM	A	608	43/43	0.95	0.08	31,39,44,49	0
5	CA	A	603	1/1	0.95	0.05	43,43,43,43	0
5	CA	D	603	1/1	0.95	0.09	36,36,36,36	0
8	HEM	D	608	43/43	0.95	0.09	19,25,39,45	0

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.




Electron density around HEM B 606:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)






Electron density around HEM C 608:


$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)



Electron density around HEM A 608:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)





6.5 Other polymers ⓘ

There are no such residues in this entry.