



wwPDB X-ray Structure Validation Summary Report ⓘ

Mar 8, 2026 – 03:39 PM UTC

PDB ID : 2I6E / pdb_00002i6e
Title : Crystal structure of protein DR0370 from *Deinococcus radiodurans*, Pfam DUF178
Authors : Tyagi, R.; Kumaran, D.; Burley, S.K.; Swaminathan, S.; New York SGX Research Center for Structural Genomics (NYSGXRC)
Deposited on : 2006-08-28
Resolution : 2.50 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4-5-2 with Phenix2.0
Mogul : 2022.3.0, CSD as543be (2022)
Xtriage (Phenix) : 2.0
EDS : 3.0
Percentile statistics : 20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4 : 9.0.010 (Gargrove)
Density-Fitness : 1.0.12
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.49


1 Overall quality at a glance i

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.50 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	5829 (2.50-2.50)
Clashscore	190562	6492 (2.50-2.50)
Ramachandran outliers	187476	6378 (2.50-2.50)
Sidechain outliers	187428	6380 (2.50-2.50)
RSRZ outliers	180081	5833 (2.50-2.50)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	301	
1	B	301	
1	C	301	
1	D	301	
1	E	301	

Continued on next page...

Continued from previous page...

Mol	Chain	Length	Quality of chain
1	F	301	<p>1% 64% 24% 10%</p>
1	G	301	<p>2% 68% 19% 10%</p>
1	H	301	<p>4% 62% 26% 10%</p>

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
2	SO4	B	717	-	-	X	-
2	SO4	F	718	-	-	X	-

2 Entry composition [i](#)

There are 3 unique types of molecules in this entry. The entry contains 17623 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Hypothetical protein.

Mol	Chain	Residues	Atoms						ZeroOcc	AltConf	Trace
			Total	C	N	O	S	Se			
1	A	272	2110	1351	373	381	1	4	0	0	0
1	B	272	2110	1351	373	381	1	4	0	0	0
1	C	272	2110	1351	373	381	1	4	0	0	0
1	D	272	2110	1351	373	381	1	4	0	0	0
1	E	272	2110	1351	373	381	1	4	0	0	0
1	F	272	2110	1351	373	381	1	4	0	0	0
1	G	272	2110	1351	373	381	1	4	0	0	0
1	H	272	2110	1351	373	381	1	4	0	0	0

There are 120 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	1	MSE	-	expression tag	UNP Q9RXE3
A	2	SER	-	expression tag	UNP Q9RXE3
A	3	LEU	-	expression tag	UNP Q9RXE3
A	50	MSE	MET	modified residue	UNP Q9RXE3
A	115	MSE	MET	modified residue	UNP Q9RXE3
A	175	MSE	MET	modified residue	UNP Q9RXE3
A	223	MSE	MET	modified residue	UNP Q9RXE3
A	294	GLU	-	expression tag	UNP Q9RXE3
A	295	GLY	-	expression tag	UNP Q9RXE3
A	296	HIS	-	expression tag	UNP Q9RXE3
A	297	HIS	-	expression tag	UNP Q9RXE3
A	298	HIS	-	expression tag	UNP Q9RXE3
A	299	HIS	-	expression tag	UNP Q9RXE3

Continued on next page...

Continued from previous page...

Chain	Residue	Modelled	Actual	Comment	Reference
A	300	HIS	-	expression tag	UNP Q9RXE3
A	301	HIS	-	expression tag	UNP Q9RXE3
B	1	MSE	-	expression tag	UNP Q9RXE3
B	2	SER	-	expression tag	UNP Q9RXE3
B	3	LEU	-	expression tag	UNP Q9RXE3
B	50	MSE	MET	modified residue	UNP Q9RXE3
B	115	MSE	MET	modified residue	UNP Q9RXE3
B	175	MSE	MET	modified residue	UNP Q9RXE3
B	223	MSE	MET	modified residue	UNP Q9RXE3
B	294	GLU	-	expression tag	UNP Q9RXE3
B	295	GLY	-	expression tag	UNP Q9RXE3
B	296	HIS	-	expression tag	UNP Q9RXE3
B	297	HIS	-	expression tag	UNP Q9RXE3
B	298	HIS	-	expression tag	UNP Q9RXE3
B	299	HIS	-	expression tag	UNP Q9RXE3
B	300	HIS	-	expression tag	UNP Q9RXE3
B	301	HIS	-	expression tag	UNP Q9RXE3
C	1	MSE	-	expression tag	UNP Q9RXE3
C	2	SER	-	expression tag	UNP Q9RXE3
C	3	LEU	-	expression tag	UNP Q9RXE3
C	50	MSE	MET	modified residue	UNP Q9RXE3
C	115	MSE	MET	modified residue	UNP Q9RXE3
C	175	MSE	MET	modified residue	UNP Q9RXE3
C	223	MSE	MET	modified residue	UNP Q9RXE3
C	294	GLU	-	expression tag	UNP Q9RXE3
C	295	GLY	-	expression tag	UNP Q9RXE3
C	296	HIS	-	expression tag	UNP Q9RXE3
C	297	HIS	-	expression tag	UNP Q9RXE3
C	298	HIS	-	expression tag	UNP Q9RXE3
C	299	HIS	-	expression tag	UNP Q9RXE3
C	300	HIS	-	expression tag	UNP Q9RXE3
C	301	HIS	-	expression tag	UNP Q9RXE3
D	1	MSE	-	expression tag	UNP Q9RXE3
D	2	SER	-	expression tag	UNP Q9RXE3
D	3	LEU	-	expression tag	UNP Q9RXE3
D	50	MSE	MET	modified residue	UNP Q9RXE3
D	115	MSE	MET	modified residue	UNP Q9RXE3
D	175	MSE	MET	modified residue	UNP Q9RXE3
D	223	MSE	MET	modified residue	UNP Q9RXE3
D	294	GLU	-	expression tag	UNP Q9RXE3
D	295	GLY	-	expression tag	UNP Q9RXE3
D	296	HIS	-	expression tag	UNP Q9RXE3

Continued on next page...

Continued from previous page...


Chain	Residue	Modelled	Actual	Comment	Reference
D	297	HIS	-	expression tag	UNP Q9RXE3
D	298	HIS	-	expression tag	UNP Q9RXE3
D	299	HIS	-	expression tag	UNP Q9RXE3
D	300	HIS	-	expression tag	UNP Q9RXE3
D	301	HIS	-	expression tag	UNP Q9RXE3
E	1	MSE	-	expression tag	UNP Q9RXE3
E	2	SER	-	expression tag	UNP Q9RXE3
E	3	LEU	-	expression tag	UNP Q9RXE3
E	50	MSE	MET	modified residue	UNP Q9RXE3
E	115	MSE	MET	modified residue	UNP Q9RXE3
E	175	MSE	MET	modified residue	UNP Q9RXE3
E	223	MSE	MET	modified residue	UNP Q9RXE3
E	294	GLU	-	expression tag	UNP Q9RXE3
E	295	GLY	-	expression tag	UNP Q9RXE3
E	296	HIS	-	expression tag	UNP Q9RXE3
E	297	HIS	-	expression tag	UNP Q9RXE3
E	298	HIS	-	expression tag	UNP Q9RXE3
E	299	HIS	-	expression tag	UNP Q9RXE3
E	300	HIS	-	expression tag	UNP Q9RXE3
E	301	HIS	-	expression tag	UNP Q9RXE3
F	1	MSE	-	expression tag	UNP Q9RXE3
F	2	SER	-	expression tag	UNP Q9RXE3
F	3	LEU	-	expression tag	UNP Q9RXE3
F	50	MSE	MET	modified residue	UNP Q9RXE3
F	115	MSE	MET	modified residue	UNP Q9RXE3
F	175	MSE	MET	modified residue	UNP Q9RXE3
F	223	MSE	MET	modified residue	UNP Q9RXE3
F	294	GLU	-	expression tag	UNP Q9RXE3
F	295	GLY	-	expression tag	UNP Q9RXE3
F	296	HIS	-	expression tag	UNP Q9RXE3
F	297	HIS	-	expression tag	UNP Q9RXE3
F	298	HIS	-	expression tag	UNP Q9RXE3
F	299	HIS	-	expression tag	UNP Q9RXE3
F	300	HIS	-	expression tag	UNP Q9RXE3
F	301	HIS	-	expression tag	UNP Q9RXE3
G	1	MSE	-	expression tag	UNP Q9RXE3
G	2	SER	-	expression tag	UNP Q9RXE3
G	3	LEU	-	expression tag	UNP Q9RXE3
G	50	MSE	MET	modified residue	UNP Q9RXE3
G	115	MSE	MET	modified residue	UNP Q9RXE3
G	175	MSE	MET	modified residue	UNP Q9RXE3
G	223	MSE	MET	modified residue	UNP Q9RXE3

Continued on next page...

Continued from previous page...

Chain	Residue	Modelled	Actual	Comment	Reference
G	294	GLU	-	expression tag	UNP Q9RXE3
G	295	GLY	-	expression tag	UNP Q9RXE3
G	296	HIS	-	expression tag	UNP Q9RXE3
G	297	HIS	-	expression tag	UNP Q9RXE3
G	298	HIS	-	expression tag	UNP Q9RXE3
G	299	HIS	-	expression tag	UNP Q9RXE3
G	300	HIS	-	expression tag	UNP Q9RXE3
G	301	HIS	-	expression tag	UNP Q9RXE3
H	1	MSE	-	expression tag	UNP Q9RXE3
H	2	SER	-	expression tag	UNP Q9RXE3
H	3	LEU	-	expression tag	UNP Q9RXE3
H	50	MSE	MET	modified residue	UNP Q9RXE3
H	115	MSE	MET	modified residue	UNP Q9RXE3
H	175	MSE	MET	modified residue	UNP Q9RXE3
H	223	MSE	MET	modified residue	UNP Q9RXE3
H	294	GLU	-	expression tag	UNP Q9RXE3
H	295	GLY	-	expression tag	UNP Q9RXE3
H	296	HIS	-	expression tag	UNP Q9RXE3
H	297	HIS	-	expression tag	UNP Q9RXE3
H	298	HIS	-	expression tag	UNP Q9RXE3
H	299	HIS	-	expression tag	UNP Q9RXE3
H	300	HIS	-	expression tag	UNP Q9RXE3
H	301	HIS	-	expression tag	UNP Q9RXE3

- Molecule 2 is SULFATE ION (CCD ID: SO4) (formula: O₄S).



Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
2	A	1	Total O S 5 4 1	0	0
2	A	1	Total O S 5 4 1	0	0
2	B	1	Total O S 5 4 1	0	0
2	B	1	Total O S 5 4 1	0	0
2	B	1	Total O S 5 4 1	0	0
2	C	1	Total O S 5 4 1	0	0
2	C	1	Total O S 5 4 1	0	0
2	D	1	Total O S 5 4 1	0	0
2	D	1	Total O S 5 4 1	0	0
2	E	1	Total O S 5 4 1	0	0
2	E	1	Total O S 5 4 1	0	0
2	F	1	Total O S 5 4 1	0	0
2	F	1	Total O S 5 4 1	0	0
2	F	1	Total O S 5 4 1	0	0
2	G	1	Total O S 5 4 1	0	0
2	G	1	Total O S 5 4 1	0	0
2	G	1	Total O S 5 4 1	0	0
2	H	1	Total O S 5 4 1	0	0

- Molecule 3 is water.

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
3	A	56	Total O 56 56	0	0
3	B	77	Total O 77 77	0	0

Continued on next page...


Continued from previous page...

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	C	77	Total 77	O 77	0	0
3	D	67	Total 67	O 67	0	0
3	E	65	Total 65	O 65	0	0
3	F	121	Total 121	O 121	0	0
3	G	105	Total 105	O 105	0	0
3	H	85	Total 85	O 85	0	0


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: Hypothetical protein




- Molecule 1: Hypothetical protein




- Molecule 1: Hypothetical protein







• Molecule 1: Hypothetical protein



• Molecule 1: Hypothetical protein




• Molecule 1: Hypothetical protein





• Molecule 1: Hypothetical protein



• Molecule 1: Hypothetical protein



4 Data and refinement statistics

Property	Value	Source
Space group	P 1 21 1	Depositor
Cell constants a, b, c, α , β , γ	75.31Å 139.37Å 153.62Å 90.00° 92.79° 90.00°	Depositor
Resolution (Å)	48.03 – 2.50 48.03 – 2.50	Depositor EDS
% Data completeness (in resolution range)	95.1 (48.03-2.50) 95.2 (48.03-2.50)	Depositor EDS
R_{merge}	0.10	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	4.22 (at 2.51Å)	Xtrriage
Refinement program	CNS 1.1	Depositor
R, R_{free}	0.202 , 0.235 0.202 , 0.233	Depositor DCC
R_{free} test set	3150 reflections (2.93%)	wwPDB-VP
Wilson B-factor (Å ²)	27.2	Xtrriage
Anisotropy	0.709	Xtrriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.34 , 40.3	EDS
L-test for twinning ²	$\langle L \rangle = 0.48$, $\langle L^2 \rangle = 0.31$	Xtrriage
Estimated twinning fraction	0.034 for h,-k,-l	Xtrriage
F_o, F_c correlation	0.93	EDS
Total number of atoms	17623	wwPDB-VP
Average B, all atoms (Å ²)	32.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 8.06% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: SO4

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.41	0/2162	0.90	7/2949 (0.2%)
1	B	0.41	0/2162	0.90	2/2949 (0.1%)
1	C	0.43	0/2162	0.90	2/2949 (0.1%)
1	D	0.41	0/2162	0.93	8/2949 (0.3%)
1	E	0.41	0/2162	0.93	6/2949 (0.2%)
1	F	0.43	0/2162	0.93	7/2949 (0.2%)
1	G	0.43	0/2162	0.89	4/2949 (0.1%)
1	H	0.42	0/2162	0.93	3/2949 (0.1%)
All	All	0.42	0/17296	0.91	39/23592 (0.2%)

There are no bond length outliers.

The worst 5 of 39 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	H	110	THR	N-CA-C	7.50	119.46	111.28
1	D	110	THR	N-CA-C	7.04	121.45	113.01
1	E	23	HIS	N-CA-C	6.99	120.50	109.39
1	E	90	VAL	N-CA-C	-6.74	106.67	113.47
1	G	90	VAL	N-CA-C	-6.62	107.05	113.53

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2110	0	2102	65	0
1	B	2110	0	2102	67	0
1	C	2110	0	2102	59	0
1	D	2110	0	2102	68	0
1	E	2110	0	2102	82	0
1	F	2110	0	2102	59	0
1	G	2110	0	2102	59	0
1	H	2110	0	2102	73	0
2	A	10	0	0	0	0
2	B	15	0	0	3	0
2	C	10	0	0	0	0
2	D	10	0	0	0	0
2	E	10	0	0	1	0
2	F	15	0	0	2	0
2	G	15	0	0	0	0
2	H	5	0	0	0	0
3	A	56	0	0	8	0
3	B	77	0	0	9	0
3	C	77	0	0	9	0
3	D	67	0	0	5	0
3	E	65	0	0	10	0
3	F	121	0	0	16	0
3	G	105	0	0	7	0
3	H	85	0	0	14	0
All	All	17623	0	16816	518	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 15.

The worst 5 of 518 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:125:ARG:HB3	3:B:794:HOH:O	1.56	1.05
1:D:113:SER:HB3	3:D:778:HOH:O	1.63	0.97
1:H:229:ARG:HD2	3:H:800:HOH:O	1.71	0.90
1:H:113:SER:HB2	3:H:760:HOH:O	1.71	0.90
1:C:169:LEU:HD13	1:C:178:LEU:HD21	1.51	0.90

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	270/301 (90%)	243 (90%)	23 (8%)	4 (2%)	8	16
1	B	270/301 (90%)	256 (95%)	13 (5%)	1 (0%)	30	49
1	C	270/301 (90%)	257 (95%)	12 (4%)	1 (0%)	30	49
1	D	270/301 (90%)	256 (95%)	14 (5%)	0	100	100
1	E	270/301 (90%)	253 (94%)	17 (6%)	0	100	100
1	F	270/301 (90%)	253 (94%)	17 (6%)	0	100	100
1	G	270/301 (90%)	259 (96%)	11 (4%)	0	100	100
1	H	270/301 (90%)	259 (96%)	11 (4%)	0	100	100
All	All	2160/2408 (90%)	2036 (94%)	118 (6%)	6 (0%)	36	55

5 of 6 Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	174	THR
1	A	173	ARG
1	B	23	HIS
1	C	243	GLU
1	A	23	HIS

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	220/239 (92%)	213 (97%)	7 (3%)	34	62

Continued on next page...

Continued from previous page...

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles
1	B	220/239 (92%)	214 (97%)	6 (3%)	39 67
1	C	220/239 (92%)	211 (96%)	9 (4%)	27 53
1	D	220/239 (92%)	213 (97%)	7 (3%)	34 62
1	E	220/239 (92%)	211 (96%)	9 (4%)	27 53
1	F	220/239 (92%)	210 (96%)	10 (4%)	24 49
1	G	220/239 (92%)	207 (94%)	13 (6%)	18 37
1	H	220/239 (92%)	213 (97%)	7 (3%)	34 62
All	All	1760/1912 (92%)	1692 (96%)	68 (4%)	28 55

5 of 68 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	G	178	LEU
1	G	240	ARG
1	H	174	THR
1	D	111	SER
1	D	50	MSE

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 52 such sidechains are listed below:

Mol	Chain	Res	Type
1	E	97	HIS
1	F	180	HIS
1	H	239	GLN
1	E	239	GLN
1	F	72	HIS

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

18 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
2	SO4	A	701	-	4,4,4	0.38	0	6,6,6	0.07	0
2	SO4	C	711	-	4,4,4	0.30	0	6,6,6	0.11	0
2	SO4	D	712	-	4,4,4	0.37	0	6,6,6	0.17	0
2	SO4	H	716	-	4,4,4	0.37	0	6,6,6	0.09	0
2	SO4	C	703	-	4,4,4	0.32	0	6,6,6	0.10	0
2	SO4	B	702	-	4,4,4	0.32	0	6,6,6	0.07	0
2	SO4	F	706	-	4,4,4	0.33	0	6,6,6	0.06	0
2	SO4	B	710	-	4,4,4	0.35	0	6,6,6	0.11	0
2	SO4	G	708	-	4,4,4	0.36	0	6,6,6	0.09	0
2	SO4	E	713	-	4,4,4	0.38	0	6,6,6	0.14	0
2	SO4	G	707	-	4,4,4	0.35	0	6,6,6	0.14	0
2	SO4	F	718	-	4,4,4	0.15	0	6,6,6	0.40	0
2	SO4	G	715	-	4,4,4	0.36	0	6,6,6	0.11	0
2	SO4	E	705	-	4,4,4	0.35	0	6,6,6	0.13	0
2	SO4	A	709	-	4,4,4	0.38	0	6,6,6	0.08	0
2	SO4	F	714	-	4,4,4	0.25	0	6,6,6	0.16	0
2	SO4	D	704	-	4,4,4	0.35	0	6,6,6	0.16	0
2	SO4	B	717	-	4,4,4	0.37	0	6,6,6	0.17	0

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

4 monomers are involved in 6 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	B	710	SO4	1	0
2	E	713	SO4	1	0
2	F	718	SO4	2	0
2	B	717	SO4	2	0

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	268/301 (89%)	0.51	26 (9%) 13 11	20, 38, 63, 76	0
1	B	268/301 (89%)	0.01	4 (1%) 72 68	15, 30, 50, 71	0
1	C	268/301 (89%)	-0.06	3 (1%) 78 75	14, 30, 47, 64	0
1	D	268/301 (89%)	0.10	9 (3%) 48 43	14, 31, 54, 71	0
1	E	268/301 (89%)	0.23	16 (5%) 27 24	15, 33, 58, 74	0
1	F	268/301 (89%)	-0.23	4 (1%) 72 68	13, 25, 43, 70	0
1	G	268/301 (89%)	-0.24	7 (2%) 57 52	14, 25, 44, 68	0
1	H	268/301 (89%)	-0.11	12 (4%) 38 33	12, 26, 49, 69	0
All	All	2144/2408 (89%)	0.02	81 (3%) 44 39	12, 30, 54, 76	0

The worst 5 of 81 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	16	PRO	5.4
1	D	16	PRO	4.1
1	E	24	PHE	4.0
1	G	173	ARG	3.6
1	B	16	PRO	3.5

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
2	SO4	B	717	5/5	0.90	0.13	71,72,73,73	0
2	SO4	B	710	5/5	0.91	0.20	63,65,66,67	0
2	SO4	D	712	5/5	0.91	0.14	71,72,72,73	0
2	SO4	F	718	5/5	0.91	0.15	70,70,70,71	0
2	SO4	H	716	5/5	0.91	0.16	66,67,68,68	0
2	SO4	A	709	5/5	0.92	0.13	65,66,67,67	0
2	SO4	E	713	5/5	0.92	0.13	62,62,62,63	0
2	SO4	C	711	5/5	0.93	0.13	45,45,47,48	0
2	SO4	A	701	5/5	0.93	0.10	49,49,51,52	0
2	SO4	E	705	5/5	0.93	0.11	50,50,52,53	0
2	SO4	G	715	5/5	0.94	0.17	53,54,57,57	0
2	SO4	G	708	5/5	0.96	0.09	46,46,48,49	0
2	SO4	D	704	5/5	0.97	0.08	37,38,39,39	0
2	SO4	B	702	5/5	0.97	0.07	42,44,44,44	0
2	SO4	F	706	5/5	0.97	0.08	39,42,43,43	0
2	SO4	F	714	5/5	0.97	0.10	35,36,39,40	0
2	SO4	G	707	5/5	0.98	0.06	35,35,36,37	0
2	SO4	C	703	5/5	0.98	0.07	35,36,39,40	0

6.5 Other polymers [i](#)

There are no such residues in this entry.