



wwPDB X-ray Structure Validation Summary Report ⓘ

Mar 9, 2026 – 04:46 PM UTC

PDB ID : 4IGN / pdb_00004ign
Title : 2.32 Angstrom X-ray Crystal structure of R47A mutant of human ACMSD
Authors : Liu, F.; Liu, A.
Deposited on : 2012-12-17
Resolution : 2.33 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4-5-2 with Phenix2.0
Xtrriage (Phenix) : 2.0
EDS : 3.0
Percentile statistics : 20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4 : 9.0.010 (Gargrove)
Density-Fitness : 1.0.12
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.33 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.




Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	7754 (2.34-2.30)
Clashscore	190562	8383 (2.34-2.30)
Ramachandran outliers	187476	8303 (2.34-2.30)
Sidechain outliers	187428	8303 (2.34-2.30)
RSRZ outliers	180081	7760 (2.34-2.30)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	332	
1	B	332	
1	C	332	
1	D	332	
1	E	332	

Continued on next page...

Continued from previous page...

Mol	Chain	Length	Quality of chain
1	F	332	 <p>A horizontal bar chart representing the quality of chain. The bar is divided into four segments: a small red segment at the far left labeled '4%', a large green segment labeled '74%', a yellow segment labeled '21%', and a small orange segment at the far right labeled '5%'.</p>

2 Entry composition [i](#)

There are 3 unique types of molecules in this entry. The entry contains 16601 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called 2-amino-3-carboxymuconate-6-semialdehyde decarboxylase.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	332	2629	1700	440	468	21	0	0	0
1	B	332	2629	1700	440	468	21	0	0	0
1	C	332	2629	1700	440	468	21	0	0	0
1	D	332	2629	1700	440	468	21	0	0	0
1	E	332	2629	1700	440	468	21	0	0	0
1	F	332	2629	1700	440	468	21	0	0	0

There are 6 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	47	ALA	ARG	engineered mutation	UNP Q8TDX5
B	47	ALA	ARG	engineered mutation	UNP Q8TDX5
C	47	ALA	ARG	engineered mutation	UNP Q8TDX5
D	47	ALA	ARG	engineered mutation	UNP Q8TDX5
E	47	ALA	ARG	engineered mutation	UNP Q8TDX5
F	47	ALA	ARG	engineered mutation	UNP Q8TDX5

- Molecule 2 is ZINC ION (CCD ID: ZN) (formula: Zn).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
2	A	1	Total	Zn	0	0
			1	1		
2	B	1	Total	Zn	0	0
			1	1		
2	C	1	Total	Zn	0	0
			1	1		

Continued on next page...

Continued from previous page...

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
2	D	1	Total 1	Zn 1	0	0
2	E	1	Total 1	Zn 1	0	0
2	F	1	Total 1	Zn 1	0	0


- Molecule 3 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	A	142	Total 142	O 142	0	0
3	B	175	Total 175	O 175	0	0
3	C	148	Total 148	O 148	0	0
3	D	145	Total 145	O 145	0	0
3	E	129	Total 129	O 129	0	0
3	F	82	Total 82	O 82	0	0


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: 2-amino-3-carboxymuconate-6-semialdehyde decarboxylase



- Molecule 1: 2-amino-3-carboxymuconate-6-semialdehyde decarboxylase





- Molecule 1: 2-amino-3-carboxymuconate-6-semialdehyde decarboxylase









- Molecule 1: 2-amino-3-carboxymuconate-6-semialdehyde decarboxylase



- Molecule 1: 2-amino-3-carboxymuconate-6-semialdehyde decarboxylase



- Molecule 1: 2-amino-3-carboxymuconate-6-semialdehyde decarboxylase



4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, α , β , γ	89.14Å 101.69Å 232.61Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	50.00 – 2.33 50.00 – 2.33	Depositor EDS
% Data completeness (in resolution range)	98.3 (50.00-2.33) 98.3 (50.00-2.33)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	2.78 (at 2.34Å)	Xtrriage
Refinement program	REFMAC 5.6.0117	Depositor
R, R_{free}	0.208 , 0.276 (Not available) , 0.235	Depositor DCC
R_{free} test set	4492 reflections (4.93%)	wwPDB-VP
Wilson B-factor (Å ²)	29.7	Xtrriage
Anisotropy	0.153	Xtrriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.30 , 24.4	EDS
L-test for twinning ²	$\langle L \rangle = 0.51$, $\langle L^2 \rangle = 0.34$	Xtrriage
Estimated twinning fraction	No twinning to report.	Xtrriage
F_o, F_c correlation	0.94	EDS
Total number of atoms	16601	wwPDB-VP
Average B, all atoms (Å ²)	30.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 7.74% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality i

5.1 Standard geometry i

Bond lengths and bond angles in the following residue types are not validated in this section:
ZN

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	1.10	3/2700 (0.1%)	1.13	6/3655 (0.2%)
1	B	1.13	3/2700 (0.1%)	1.22	20/3655 (0.5%)
1	C	1.11	3/2700 (0.1%)	1.17	9/3655 (0.2%)
1	D	1.10	2/2700 (0.1%)	1.10	4/3655 (0.1%)
1	E	1.02	1/2700 (0.0%)	1.12	6/3655 (0.2%)
1	F	0.98	0/2700	1.13	11/3655 (0.3%)
All	All	1.07	12/16200 (0.1%)	1.14	56/21930 (0.3%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	A	0	1
1	C	0	1
1	D	0	1
All	All	0	3

The worst 5 of 12 bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	C	146	HIS	CG-CD2	6.54	1.43	1.35
1	A	54	TRP	CD2-CE2	6.39	1.52	1.41
1	A	148	ASN	CA-C	6.08	1.58	1.52
1	C	6	HIS	CA-C	-5.81	1.46	1.53
1	C	146	HIS	ND1-CE1	5.80	1.38	1.32

The worst 5 of 56 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	F	121	ALA	CA-C-N	10.65	133.15	119.84
1	F	121	ALA	C-N-CA	10.65	133.15	119.84
1	F	44	LYS	N-CA-C	8.25	120.50	108.86
1	B	189	LEU	CA-C-N	-7.89	111.59	119.56
1	B	189	LEU	C-N-CA	-7.89	111.59	119.56

There are no chirality outliers.

All (3) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	A	41	LYS	Peptide
1	C	1	MET	Peptide
1	D	41	LYS	Peptide

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2629	0	2635	71	0
1	B	2629	0	2635	65	0
1	C	2629	0	2635	77	0
1	D	2629	0	2635	67	0
1	E	2629	0	2635	54	0
1	F	2629	0	2635	53	0
2	A	1	0	0	0	0
2	B	1	0	0	0	0
2	C	1	0	0	0	0
2	D	1	0	0	0	0
2	E	1	0	0	0	0
2	F	1	0	0	0	0
3	A	142	0	0	11	0
3	B	175	0	0	8	0
3	C	148	0	0	15	0
3	D	145	0	0	8	0
3	E	129	0	0	5	0
3	F	82	0	0	1	0
All	All	16601	0	15810	370	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 12.

The worst 5 of 370 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:D:42:ASP:HA	3:D:512:HOH:O	1.42	1.15
1:B:32:HIS:HB3	3:B:631:HOH:O	1.44	1.13
1:B:332:ARG:CD	1:B:332:ARG:H	1.62	1.12
1:C:175:PRO:HG2	1:C:195:MET:HE3	1.28	1.11
1:B:13:GLU:HB2	3:B:519:HOH:O	1.50	1.11

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	330/332 (99%)	315 (96%)	14 (4%)	1 (0%)	36	45
1	B	330/332 (99%)	315 (96%)	13 (4%)	2 (1%)	21	25
1	C	330/332 (99%)	311 (94%)	18 (6%)	1 (0%)	36	45
1	D	330/332 (99%)	316 (96%)	13 (4%)	1 (0%)	36	45
1	E	330/332 (99%)	308 (93%)	21 (6%)	1 (0%)	36	45
1	F	330/332 (99%)	303 (92%)	20 (6%)	7 (2%)	5	4
All	All	1980/1992 (99%)	1868 (94%)	99 (5%)	13 (1%)	18	22

5 of 13 Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	E	42	ASP
1	B	42	ASP
1	B	43	GLY

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	C	42	ASP
1	F	34	LYS

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles
1	A	284/284 (100%)	257 (90%)	27 (10%)	8 10
1	B	284/284 (100%)	261 (92%)	23 (8%)	11 14
1	C	284/284 (100%)	260 (92%)	24 (8%)	10 13
1	D	284/284 (100%)	260 (92%)	24 (8%)	10 13
1	E	284/284 (100%)	264 (93%)	20 (7%)	14 18
1	F	284/284 (100%)	259 (91%)	25 (9%)	9 12
All	All	1704/1704 (100%)	1561 (92%)	143 (8%)	10 13

5 of 143 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	E	290	THR
1	F	20	ARG
1	F	100	LEU
1	B	319	LEU
1	B	318	LYS

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 21 such sidechains are listed below:

Mol	Chain	Res	Type
1	D	251	ASN
1	E	238	HIS
1	F	317	ASN
1	E	249	GLN
1	E	155	GLN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

Of 6 ligands modelled in this entry, 6 are monoatomic - leaving 0 for Mogul analysis.

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	332/332 (100%)	-0.21	4 (1%) 76 78	14, 26, 45, 67	0
1	B	332/332 (100%)	-0.40	3 (0%) 81 82	13, 22, 41, 72	0
1	C	332/332 (100%)	-0.21	6 (1%) 67 70	14, 26, 49, 86	0
1	D	332/332 (100%)	-0.35	4 (1%) 76 78	15, 25, 42, 70	0
1	E	332/332 (100%)	-0.07	9 (2%) 56 59	18, 30, 52, 90	0
1	F	332/332 (100%)	0.17	13 (3%) 43 46	19, 34, 69, 103	0
All	All	1992/1992 (100%)	-0.18	39 (1%) 65 67	13, 26, 52, 103	0

The worst 5 of 39 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	F	1	MET	6.1
1	D	1	MET	5.9
1	A	1	MET	5.4
1	C	1	MET	5.1
1	E	1	MET	5.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
2	ZN	A	401	1/1	1.00	0.02	19,19,19,19	0
2	ZN	B	401	1/1	1.00	0.03	19,19,19,19	0
2	ZN	C	401	1/1	1.00	0.01	22,22,22,22	0
2	ZN	D	401	1/1	1.00	0.01	25,25,25,25	0
2	ZN	E	401	1/1	1.00	0.01	26,26,26,26	0
2	ZN	F	401	1/1	1.00	0.01	29,29,29,29	0

6.5 Other polymers [i](#)

There are no such residues in this entry.