Full wwPDB X-ray Structure Validation Report ⓘ

Mar 6, 2026 – 07:25 PM UTC

PDB ID	:	8IU9 / pdb_00008iu9
Title	:	Crystal structure of GH66 endodextranase from Flavobacterium johnsoniae in complex with glucose
Authors	:	Nakamura, S.; Miyazaki, T.
Deposited on	:	2023-03-24
Resolution	:	1.80 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at *validation@mail.wwpdb.org*
A user guide is available at
https://www.wwpdb.org/validation/2017/XrayValidationReportHelp
with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at
http://www.wwpdb.org/validation/2017/FAQs#types.

The following versions of software and data (see references ⓘ) were used in the production of this report:


MolProbity	:	4-5-2 with Phenix2.0
Mogul	:	2022.3.0, CSD as543be (2022)
Xtriage (Phenix)	:	2.0
EDS	:	3.0
Buster-report	:	wwPDB partial adaption of 1.1.7 (2018)
Percentile statistics	:	20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4	:	9.0.010 (Gargrove)
Density-Fitness	:	1.0.12
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.49

1 Overall quality at a glance (i)

The following experimental techniques were used to determine the structure:
X-RAY DIFFRACTION

The reported resolution of this entry is 1.80 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	7662 (1.80-1.80)
Clashscore	190562	8479 (1.80-1.80)
Ramachandran outliers	187476	8391 (1.80-1.80)
Sidechain outliers	187428	8390 (1.80-1.80)
RSRZ outliers	180081	7663 (1.80-1.80)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for >=3, 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions <=5% The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	576	% 78% 16% • 5%

2 Entry composition (i)

There are 5 unique types of molecules in this entry. The entry contains 8900 atoms, of which 4233 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.


- Molecule 1 is a protein called Candidate dextranase Glycoside hydrolase family 66.

Mol	Chain	Residues	Atoms						ZeroOcc	AltConf	Trace
			Total	C	H	N	O	S			
1	A	548	8564	2784	4197	726	842	15	124	2	0

There are 23 discrepancies between the modelled and reference sequences:


Chain	Residue	Modelled	Actual	Comment	Reference
A	11	MET	-	initiating methionine	UNP A5FBI2
A	12	GLY	-	expression tag	UNP A5FBI2
A	13	SER	-	expression tag	UNP A5FBI2
A	14	SER	-	expression tag	UNP A5FBI2
A	15	HIS	-	expression tag	UNP A5FBI2
A	16	HIS	-	expression tag	UNP A5FBI2
A	17	HIS	-	expression tag	UNP A5FBI2
A	18	HIS	-	expression tag	UNP A5FBI2
A	19	HIS	-	expression tag	UNP A5FBI2
A	20	HIS	-	expression tag	UNP A5FBI2
A	21	SER	-	expression tag	UNP A5FBI2
A	22	SER	-	expression tag	UNP A5FBI2
A	23	GLY	-	expression tag	UNP A5FBI2
A	24	LEU	-	expression tag	UNP A5FBI2
A	25	VAL	-	expression tag	UNP A5FBI2
A	26	PRO	-	expression tag	UNP A5FBI2
A	27	ARG	-	expression tag	UNP A5FBI2
A	28	GLY	-	expression tag	UNP A5FBI2
A	29	SER	-	expression tag	UNP A5FBI2
A	30	HIS	-	expression tag	UNP A5FBI2
A	31	MET	-	expression tag	UNP A5FBI2
A	32	ALA	-	expression tag	UNP A5FBI2
A	33	SER	-	expression tag	UNP A5FBI2

- Molecule 2 is alpha-D-glucopyranose (CCD ID: GLC) (formula: $C_6H_{12}O_6$) (labeled as "Ligand of Interest" by depositor).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
2	A	1	Total	C	H	O	4	0
			24	6	12	6		
2	A	1	Total	C	H	O	4	0
			24	6	12	6		

- Molecule 3 is beta-D-glucopyranose (CCD ID: BGC) (formula: $C_6H_{12}O_6$) (labeled as "Ligand of Interest" by depositor).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
3	A	1	Total	C	H	O	4	0
			24	6	12	6		

- Molecule 4 is SODIUM ION (CCD ID: NA) (formula: Na).

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
4	A	1	Total Na 1 1	0	0


- Molecule 5 is water.

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
5	A	263	Total O 263 263	0	0

3 Residue-property plots (i)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density (RSRZ > 2). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

• Molecule 1: Candidate dextranase Glycoside hydrolase family 66



4 Data and refinement statistics (i)

Property	Value	Source
Space group	C 1 2 1	Depositor
Cell constants a, b, c, α, β, γ	155.18Å 48.38Å 76.49Å 90.00° 104.41° 90.00°	Depositor
Resolution (Å)	47.26 – 1.80 47.26 – 1.80	Depositor EDS
% Data completeness (in resolution range)	98.3 (47.26-1.80) 98.3 (47.26-1.80)	Depositor EDS
R_{merge}	0.13	Depositor
R_{sym}	(Not available)	Depositor
$< I/\sigma(I) >$ [1]	2.98 (at 1.79Å)	Xtriage
Refinement program	REFMAC 5.8.0405	Depositor
R, R_{free}	0.206 , 0.259 0.211 , 0.263	Depositor DCC
R_{free} test set	2464 reflections (4.80%)	wwPDB-VP
Wilson B-factor (Å^2)	19.7	Xtriage
Anisotropy	0.560	Xtriage
Bulk solvent k_{sol}(e/Å^3), B_{sol}(Å^2)	0.43 , 36.0	EDS
L-test for twinning[2]	$< \|L\| > = 0.49$, $< L^2 > = 0.32$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o,F_c correlation	0.95	EDS
Total number of atoms	8900	wwPDB-VP
Average B, all atoms (Å^2)	29.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 10.87% of the height of the origin peak. No significant pseudotranslation is detected.*

[1] Intensities estimated from amplitudes.

[2] Theoretical values of $< |L| >$, $< L^2 >$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: BGC, NA, GLC

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	#$\|Z\|$ >5	RMSZ	#$\|Z\|$ >5
1	A	1.03	2/4478 (0.0%)	1.45	22/6082 (0.4%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand.A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	A	0	2

All (2) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	159	HIS	ND1-CE1	5.45	1.38	1.32
1	A	505	HIS	CG-CD2	5.36	1.41	1.35

All (22) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(^{o})	Ideal(^{o})
1	A	83	THR	CA-CB-OG1	-8.48	96.88	109.60
1	A	92	LEU	CA-C-O	8.16	124.59	119.29
1	A	216	PHE	CA-CB-CG	8.02	121.82	113.80
1	A	133	ARG	NE-CZ-NH2	7.97	126.37	119.20
1	A	452	ASP	CA-CB-CG	7.23	119.83	112.60
1	A	244	HIS	CA-CB-CG	6.77	120.57	113.80
1	A	484	TRP	N-CA-C	-6.22	101.50	110.40
1	A	290	PHE	CA-CB-CG	5.99	119.79	113.80
1	A	469	ASN	CA-CB-CG	-5.84	106.76	112.60
1	A	480	ASN	CB-CA-C	5.78	121.16	111.22
1	A	552	ASP	CA-CB-CG	5.72	118.33	112.60

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(^{o})	Ideal(^{o})
1	A	206	GLN	CB-CA-C	5.49	119.57	110.90
1	A	228	ASP	CA-CB-CG	5.47	118.07	112.60
1	A	554	ASN	CB-CA-C	-5.46	104.13	111.89
1	A	148	ARG	NE-CZ-NH1	-5.39	116.11	121.50
1	A	436	ASN	CA-C-O	-5.35	114.85	120.63
1	A	424	GLU	CA-C-N	5.31	130.56	122.11
1	A	424	GLU	C-N-CA	5.31	130.56	122.11
1	A	419	HIS	CB-CA-C	5.30	118.97	110.16
1	A	390	ASN	CA-CB-CG	5.17	117.77	112.60
1	A	462	ARG	CA-C-O	-5.11	116.92	121.67
1	A	261	TYR	CB-CA-C	5.08	118.80	109.46

There are no chirality outliers.

All (2) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	A	193	ARG	Sidechain
1	A	71	ARG	Sidechain

5.2 Too-close contacts (i)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	4367	4197	4182	50	0
2	A	24	24	24	1	0
3	A	12	12	12	0	0
4	A	1	0	0	0	0
5	A	263	0	0	3	0
All	All	4667	4233	4218	50	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 6.

All (50) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:139:GLU:HG3	1:A:143:ILE:HD11	1.47	0.96
1:A:42:ILE:HD12	1:A:117:LEU:HB3	1.62	0.81
1:A:583:ILE:HD12	1:A:583:ILE:N	2.11	0.66
1:A:42:ILE:HD11	1:A:65:PHE:CD1	2.32	0.65
1:A:582:MET:C	1:A:583:ILE:HD12	2.28	0.58
1:A:42:ILE:HD11	1:A:65:PHE:HD1	1.68	0.57
1:A:323:LYS:HE2	2:A:603:GLC:O3	2.06	0.56
1:A:583:ILE:N	1:A:583:ILE:CD1	2.71	0.54
1:A:116:ILE:HD13	1:A:554:ASN:ND2	2.22	0.54
1:A:279:ASP:OD2	1:A:283:LYS:NZ	2.39	0.53
1:A:409:ASP:OD2	1:A:421:GLU:OE2	2.27	0.52
1:A:136:PHE:CE1	1:A:423:GLY:N	2.78	0.51
1:A:71:ARG:HG2	1:A:81:GLU:HG2	1.91	0.50
1:A:191:ILE:HG23	1:A:256:PHE:CE1	2.47	0.50
1:A:532:GLN:NE2	1:A:575:PRO:HA	2.27	0.49
1:A:130:LYS:HD3	1:A:131:PHE:CZ	2.48	0.49
1:A:191:ILE:HD11	1:A:193:ARG:HD3	1.95	0.48
1:A:250:HIS:HB2	1:A:295:LEU:HD13	1.96	0.48
1:A:563:PHE:CD1	1:A:563:PHE:C	2.92	0.48
1:A:188:ASN:HA	1:A:193:ARG:O	2.14	0.47
1:A:432:PHE:N	1:A:433:PRO:CD	2.78	0.47
1:A:38:VAL:HG13	1:A:445[A]:SER:OG	2.15	0.47
1:A:132:PRO:HD2	1:A:462:ARG:NH1	2.30	0.47
1:A:189:ASP:OD1	1:A:193:ARG:HG2	2.15	0.47
1:A:68:SER:HA	1:A:106:TYR:O	2.15	0.47
1:A:73:THR:HA	1:A:77:LYS:O	2.16	0.46
1:A:480:ASN:ND2	1:A:490:LYS:HG2	2.30	0.45
1:A:301:VAL:HG23	1:A:309:ALA:HB3	1.99	0.45
1:A:315:PHE:N	1:A:316:PRO:CD	2.79	0.45
1:A:366:LEU:HG	1:A:517:TRP:CD1	2.52	0.45
1:A:479:LEU:HD23	1:A:530:ILE:HD13	1.99	0.44
1:A:189:ASP:HB2	5:A:822:HOH:O	2.17	0.44
1:A:352:LEU:HD11	1:A:377:SER:HB3	2.00	0.44
1:A:420:LEU:C	1:A:420:LEU:HD23	2.43	0.44
1:A:262:ILE:HD11	1:A:295:LEU:HD12	2.00	0.44
1:A:352:LEU:O	1:A:383:THR:HA	2.18	0.44
1:A:573:LYS:HE3	1:A:573:LYS:HB2	1.82	0.44
1:A:410:ALA:HB1	1:A:580:TRP:HB2	2.00	0.43
1:A:107:GLN:HB2	1:A:117:LEU:HD11	2.00	0.43
1:A:439[A]:MET:HG2	1:A:443:LEU:HD23	2.01	0.43
1:A:96:ASP:CG	5:A:890:HOH:O	2.62	0.43
1:A:187:TRP:CZ2	1:A:195:VAL:HG11	2.54	0.43

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:240:LYS:HE3	1:A:248:ASP:HA	2.00	0.42
1:A:484:TRP:CG	1:A:485:PRO:HA	2.54	0.42
1:A:63:LYS:HE3	1:A:63:LYS:HB3	1.72	0.42
1:A:389:MET:HE3	1:A:389:MET:HB3	1.84	0.42
1:A:107:GLN:O	1:A:114:GLN:HA	2.20	0.42
1:A:243:ASN:O	1:A:244:HIS:C	2.63	0.41
1:A:191:ILE:HG12	5:A:780:HOH:O	2.20	0.41
1:A:472:ILE:HD11	1:A:535:LEU:HD13	2.03	0.41

There are no symmetry-related clashes.

5.3 Torsion angles (i)

5.3.1 Protein backbone (i)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	546/576 (95%)	526 (96%)	20 (4%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains (i)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	469/489 (96%)	460 (98%)	9 (2%)	50	41

All (9) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	39	THR
1	A	221	TYR
1	A	486	PRO
1	A	497	GLN
1	A	542	LYS
1	A	548	TYR
1	A	564	SER
1	A	571	ASN
1	A	583	ILE

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (14) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	74	HIS
1	A	109	ASN
1	A	230	ASN
1	A	234	GLN
1	A	276	GLN
1	A	326	ASN
1	A	378	ASN
1	A	379	ASN
1	A	401	ASN
1	A	480	ASN
1	A	532	GLN
1	A	536	ASN
1	A	539	HIS
1	A	571	ASN

5.3.3 RNA (i)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains (i)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates (i)

There are no oligosaccharides in this entry.

5.6 Ligand geometry (i)

Of 4 ligands modelled in this entry, 1 is monoatomic - leaving 3 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\#\|Z\| > 2$	Counts	RMSZ	$\#\|Z\| > 2$
3	BGC	A	602	-	12,12,12	0.91	1 (8%)	17,17,17	1.16	1 (5%)
2	GLC	A	601	-	12,12,12	1.18	2 (16%)	17,17,17	1.29	2 (11%)
2	GLC	A	603	-	12,12,12	1.27	1 (8%)	17,17,17	1.39	2 (11%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	BGC	A	602	-	-	0/2/22/22	0/1/1/1
2	GLC	A	601	-	-	2/2/22/22	0/1/1/1
2	GLC	A	603	-	-	0/2/22/22	0/1/1/1

All (4) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	A	603	GLC	C3-C2	2.75	1.59	1.52
2	A	601	GLC	C4-C5	2.74	1.58	1.53
3	A	602	BGC	O5-C1	2.32	1.48	1.42
2	A	601	GLC	O1-C1	2.04	1.46	1.39

All (5) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(^{o})	Ideal(^{o})
2	A	603	GLC	O1-C1-C2	3.63	119.51	108.98
2	A	601	GLC	O1-C1-O5	-2.79	102.13	110.41
2	A	601	GLC	O1-C1-C2	2.53	116.31	108.98
2	A	603	GLC	O2-C2-C3	2.35	115.91	110.38
3	A	602	BGC	O5-C1-C2	2.21	114.19	110.30

There are no chirality outliers.

All (2) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
2	A	601	GLC	C4-C5-C6-O6
2	A	601	GLC	O5-C5-C6-O6


There are no ring outliers.

1 monomer is involved in 1 short contact:


Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	A	603	GLC	1	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less then 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.


Ligand BGC A 602




Bond lengths



Bond angles




Torsions




Rings


Ligand GLC A 601




Bond lengths



Bond angles




Torsions




Rings


Ligand GLC A 603




Bond lengths



Bond angles



Torsions



Rings

5.7 Other polymers

There are no such residues in this entry.

5.8 Polymer linkage issues

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

In the following table, the column labelled '#RSRZ> 2' contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95^{th} percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled 'Q< 0.9' lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2			OWAB(Å^2)	Q<0.9
1	A	548/576 (95%)	-0.10	7 (1%)	75	75	10, 25, 49, 80	2 (0%)

All (7) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	38	VAL	5.0
1	A	39	THR	3.2
1	A	111	ASP	2.9
1	A	112	GLY	2.6
1	A	113	THR	2.5
1	A	300	ASN	2.0
1	A	574	VAL	2.0

6.2 Non-standard residues in protein, DNA, RNA chains

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates

There are no oligosaccharides in this entry.

6.4 Ligands

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95^{th} percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å^2)	Q<0.9
2	GLC	A	603	12/12	0.88	0.10	31,34,38,48	4

Continued on next page...


Continued from previous page...

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å^2)	Q<0.9
3	BGC	A	602	12/12	0.93	0.07	24,27,38,38	4
2	GLC	A	601	12/12	0.94	0.07	19,26,38,38	4
4	NA	A	604	1/1	0.96	0.08	29,29,29,29	0

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.


Electron density around GLC A 603:

$2mF_o$-DF_c (at 0.7 rmsd) in gray
mF_o-DF_c (at 3 rmsd) in purple (negative) and green (positive)




Electron density around BGC A 602:

$2mF_o$-DF_c (at 0.7 rmsd) in gray
mF_o-DF_c (at 3 rmsd) in purple (negative) and green (positive)



Electron density around GLC A 601:

$2mF_o$-DF_c (at 0.7 rmsd) in gray
mF_o-DF_c (at 3 rmsd) in purple (negative) and green (positive)



6.5 Other polymers (i)

There are no such residues in this entry.