



wwPDB X-ray Structure Validation Summary Report ⓘ

Aug 4, 2026 – 12:02 AM EDT

PDB ID : 1KFY / pdb_00001kfy
Title : QUINOL-FUMARATE REDUCTASE WITH QUINOL INHIBITOR 2-[1-(4-CHLORO-PHENYL)-ETHYL]-4,6-DINITRO-PHENOL
Authors : Iverson, T.M.; Luna-Chavez, C.; Croal, L.R.; Cecchini, G.; Rees, D.C.
Deposited on : 2001-11-24
Resolution : 3.60 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4-5-2 with Phenix2.0
Mogul : 2022.3.0, CSD as543be (2022)
Xtriage (Phenix) : **NOT EXECUTED**
EDS : **NOT EXECUTED**
Buster-report : wwPDB partial adaption of 1.1.7 (2018)
Percentile statistics : 20250101.v01 (using entries in the PDB archive January 1st 2025)
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.50


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 3.60 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
Clashscore	190562	1827 (3.70-3.50)
Ramachandran outliers	187476	1773 (3.70-3.50)
Sidechain outliers	187428	1772 (3.70-3.50)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$.

Note EDS was not executed.

Mol	Chain	Length	Quality of chain
1	A	602	
1	M	602	
2	B	243	
2	N	243	
3	C	130	
3	O	130	
4	D	119	
4	P	119	

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
10	BRS	B	700	-	-	X	-
5	OAA	A	702	-	X	-	-
8	F3S	N	245	-	-	X	-
9	SF4	N	246	-	-	X	-

2 Entry composition

There are 11 unique types of molecules in this entry. The entry contains 16957 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called FUMARATE REDUCTASE FLAVOPROTEIN.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	577	Total	C	N	O	S	0	0	0
			4448	2775	802	840	31			
1	M	577	Total	C	N	O	S	0	0	0
			4448	2775	802	840	31			

- Molecule 2 is a protein called Fumarate reductase iron-sulfur protein.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
2	B	243	Total	C	N	O	S	0	0	0
			1888	1189	323	357	19			
2	N	243	Total	C	N	O	S	0	0	0
			1888	1189	323	357	19			


- Molecule 3 is a protein called FUMARATE REDUCTASE 15 KDA HYDROPHOBIC PROTEIN.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
3	C	130	Total	C	N	O	S	0	0	0
			1058	720	166	169	3			
3	O	130	Total	C	N	O	S	0	0	0
			1058	720	166	169	3			

- Molecule 4 is a protein called FUMARATE REDUCTASE 13 KDA HYDROPHOBIC PROTEIN.


Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
4	D	119	Total	C	N	O	S	0	0	0
			926	626	151	142	7			
4	P	119	Total	C	N	O	S	0	0	0
			926	626	151	142	7			

- Molecule 5 is OXALOACETATE ION (CCD ID: OAA) (formula: $C_4H_3O_5$).




Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
5	A	1	Total C O 9 4 5	0	0
5	M	1	Total C O 9 4 5	0	0

- Molecule 6 is FLAVIN-ADENINE DINUCLEOTIDE (CCD ID: FAD) (formula: $\text{C}_{27}\text{H}_{33}\text{N}_9\text{O}_{15}\text{P}_2$).




Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
6	A	1	Total 53	C 27	N 9	O 15	P 2	0	0
6	M	1	Total 53	C 27	N 9	O 15	P 2	0	0

- Molecule 7 is FE2/S2 (INORGANIC) CLUSTER (CCD ID: FES) (formula: Fe_2S_2).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
7	B	1	Total	Fe	S	0	0
			4	2	2		
7	N	1	Total	Fe	S	0	0
			4	2	2		

- Molecule 8 is FE3-S4 CLUSTER (CCD ID: F3S) (formula: Fe_3S_4).




Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
8	B	1	Total	Fe	S	0	0
			7	3	4		

Continued on next page...

Continued from previous page...


Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
8	N	1	Total	Fe	S	0	0
			7	3	4		

- Molecule 9 is IRON/SULFUR CLUSTER (CCD ID: SF4) (formula: Fe_4S_4).




Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
9	B	1	Total	Fe	S	0	0
			8	4	4		
9	N	1	Total	Fe	S	0	0
			8	4	4		

- Molecule 10 is 2-[1-(4-CHLORO-PHENYL)-ETHYL]-4,6-DINITRO-PHENOL (CCD ID: BRS) (formula: $\text{C}_{14}\text{H}_{11}\text{ClN}_2\text{O}_5$).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
10	B	1	Total 22	C 14	Cl 1	N 2	O 5	0	0
10	N	1	Total 22	C 14	Cl 1	N 2	O 5	0	0

- Molecule 11 is O-DODECANYL OCTAETHYLENE GLYCOL (CCD ID: CE1) (formula: $C_{28}H_{58}O_9$).




Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
11	D	1	Total	C	O	0	0
			37	28	9		

Continued on next page...


Continued from previous page...

Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
11	O	1	Total	C	O	0	0
			37	28	9		
11	P	1	Total	C	O	0	0
			37	28	9		




- Molecule 2: Fumarate reductase iron-sulfur protein

Chain B: 63% 34%




- Molecule 2: Fumarate reductase iron-sulfur protein

Chain N: 46% 49%




- Molecule 3: FUMARATE REDUCTASE 15 KDA HYDROPHOBIC PROTEIN

Chain C: 58% 33% 8%




- Molecule 3: FUMARATE REDUCTASE 15 KDA HYDROPHOBIC PROTEIN

Chain O:  52% 38% 8% .





• Molecule 4: FUMARATE REDUCTASE 13 KDA HYDROPHOBIC PROTEIN

Chain D:  47% 47% . .



• Molecule 4: FUMARATE REDUCTASE 13 KDA HYDROPHOBIC PROTEIN

Chain P:  45% 49% 6%



4 Data and refinement statistics

Xtriage (Phenix) and EDS were not executed - this section is therefore incomplete.

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, α , β , γ	96.15Å 137.81Å 270.99Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	50.00 – 3.60	Depositor
% Data completeness (in resolution range)	(Not available) (50.00-3.60)	Depositor
R_{merge}	(Not available)	Depositor
R_{sym}	0.10	Depositor
Refinement program	CNS	Depositor
R, R_{free}	0.264 , 0.301	Depositor
Estimated twinning fraction	No twinning to report.	Xtriage
Total number of atoms	16957	wwPDB-VP
Average B, all atoms (Å ²)	50.0	wwPDB-VP

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: CE1, OAA, FAD, BRS, SF4, FES, F3S

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.87	3/4540 (0.1%)	1.15	26/6139 (0.4%)
1	M	0.72	2/4540 (0.0%)	1.11	24/6139 (0.4%)
2	B	0.80	1/1931 (0.1%)	1.16	9/2617 (0.3%)
2	N	0.74	2/1931 (0.1%)	1.19	10/2617 (0.4%)
3	C	0.96	3/1094 (0.3%)	1.19	9/1496 (0.6%)
3	O	0.88	2/1094 (0.2%)	1.20	10/1496 (0.7%)
4	D	1.00	3/956 (0.3%)	1.29	11/1303 (0.8%)
4	P	0.91	3/956 (0.3%)	1.19	6/1303 (0.5%)
All	All	0.83	19/17042 (0.1%)	1.16	105/23110 (0.5%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
3	O	0	1

The worst 5 of 19 bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
4	D	117	THR	CA-C	-12.36	1.36	1.52
4	P	117	THR	CA-C	-10.53	1.38	1.52
4	D	3	PRO	C-O	-8.86	1.13	1.24
1	A	135	MET	SD-CE	8.80	2.01	1.79
3	C	65	ASN	C-O	-8.29	1.13	1.24

The worst 5 of 105 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	N	241	LYS	CA-C-N	-12.81	103.83	119.84
2	N	241	LYS	C-N-CA	-12.81	103.83	119.84
4	D	117	THR	N-CA-C	10.96	134.14	110.80
3	C	65	ASN	N-CA-C	10.04	132.01	109.81
2	N	65	CYS	N-CA-C	9.10	123.98	113.15

There are no chirality outliers.

All (1) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
3	O	129	TYR	Sidechain

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	4448	0	4335	162	0
1	M	4448	0	4335	284	0
2	B	1888	0	1837	86	0
2	N	1888	0	1837	130	0
3	C	1058	0	1108	65	0
3	O	1058	0	1108	69	0
4	D	926	0	971	83	0
4	P	926	0	971	78	0
5	A	9	0	2	1	0
5	M	9	0	2	2	0
6	A	53	0	31	7	0
6	M	53	0	31	11	0
7	B	4	0	0	0	0
7	N	4	0	0	0	0
8	B	7	0	0	0	0
8	N	7	0	0	2	0
9	B	8	0	0	0	0
9	N	8	0	0	7	0
10	B	22	0	11	13	0
10	N	22	0	11	5	0
11	D	37	0	58	5	0
11	O	37	0	58	1	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
11	P	37	0	58	2	0
All	All	16957	0	16764	872	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 26.

The worst 5 of 872 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:135:MET:SD	1:A:135:MET:CE	2.01	1.48
1:A:44:HIS:NE2	6:A:703:FAD:C8M	1.81	1.41
1:M:44:HIS:NE2	6:M:803:FAD:C8M	1.83	1.38
1:M:44:HIS:NE2	6:M:803:FAD:HM82	0.88	1.21
1:A:44:HIS:NE2	6:A:703:FAD:HM82	0.85	1.17

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	575/602 (96%)	525 (91%)	43 (8%)	7 (1%)	10	41
1	M	575/602 (96%)	506 (88%)	59 (10%)	10 (2%)	7	35
2	B	241/243 (99%)	218 (90%)	19 (8%)	4 (2%)	7	35
2	N	241/243 (99%)	212 (88%)	23 (10%)	6 (2%)	4	28
3	C	128/130 (98%)	116 (91%)	9 (7%)	3 (2%)	5	30
3	O	128/130 (98%)	114 (89%)	10 (8%)	4 (3%)	3	25
4	D	117/119 (98%)	103 (88%)	13 (11%)	1 (1%)	14	46
4	P	117/119 (98%)	106 (91%)	10 (8%)	1 (1%)	14	46
All	All	2122/2188 (97%)	1900 (90%)	186 (9%)	36 (2%)	7	35

5 of 36 Ramachandran outliers are listed below:

Mol	Chain	Res	Type
2	B	242	PRO
1	M	1	GLN
1	M	244	THR
3	O	18	LYS
1	A	79	GLN

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	460/475 (97%)	447 (97%)	13 (3%)	38	60
1	M	460/475 (97%)	434 (94%)	26 (6%)	18	46
2	B	205/205 (100%)	199 (97%)	6 (3%)	37	60
2	N	205/205 (100%)	199 (97%)	6 (3%)	37	60
3	C	111/111 (100%)	105 (95%)	6 (5%)	20	47
3	O	111/111 (100%)	104 (94%)	7 (6%)	16	44
4	D	97/97 (100%)	92 (95%)	5 (5%)	21	48
4	P	97/97 (100%)	94 (97%)	3 (3%)	35	59
All	All	1746/1776 (98%)	1674 (96%)	72 (4%)	27	53

5 of 72 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
2	N	148	CYS
4	P	118	ILE
2	N	189	LYS
3	O	37	TRP
3	C	103	MET

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 54 such sidechains are listed below:

Mol	Chain	Res	Type
1	M	182	GLN
1	M	409	GLN
3	O	65	ASN
1	M	186	ASN
1	M	264	GLN

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates ⓘ

There are no oligosaccharides in this entry.

5.6 Ligand geometry ⓘ

15 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
10	BRS	N	800	-	23,23,23	2.34	6 (26%)	26,33,33	1.57	6 (23%)
9	SF4	B	246	2	0,12,12	-	-	-	-	-
5	OAA	A	702	-	8,8,8	2.22	4 (50%)	8,10,10	2.21	3 (37%)
11	CE1	O	811	-	36,36,36	1.10	0	35,35,35	1.60	10 (28%)
10	BRS	B	700	-	23,23,23	2.54	10 (43%)	26,33,33	1.87	7 (26%)
11	CE1	D	710	-	36,36,36	1.17	2 (5%)	35,35,35	1.56	8 (22%)
7	FES	B	244	2	0,4,4	-	-	-	-	-
6	FAD	A	703	-	58,58,58	2.24	12 (20%)	85,89,89	1.64	21 (24%)

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
6	FAD	M	803	-	58,58,58	1.73	14 (24%)	85,89,89	1.54	14 (16%)
7	FES	N	244	2	0,4,4	-	-	-	-	-
11	CE1	P	810	-	36,36,36	1.10	3 (8%)	35,35,35	1.67	9 (25%)
8	F3S	N	245	2	0,9,9	-	-	-	-	-
9	SF4	N	246	2	0,12,12	-	-	-	-	-
5	OAA	M	802	-	8,8,8	1.28	0	8,10,10	1.09	1 (12%)
8	F3S	B	245	2	0,9,9	-	-	-	-	-

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
10	BRS	N	800	-	-	5/12/16/16	0/2/2/2
9	SF4	B	246	2	-	-	0/6/5/5
5	OAA	A	702	-	-	4/8/8/8	-
11	CE1	O	811	-	-	17/34/34/34	-
10	BRS	B	700	-	-	6/12/16/16	0/2/2/2
11	CE1	D	710	-	-	17/34/34/34	-
7	FES	B	244	2	-	-	0/1/1/1
6	FAD	A	703	-	-	7/34/50/50	0/6/6/6
6	FAD	M	803	-	-	5/34/50/50	0/6/6/6
11	CE1	P	810	-	-	14/34/34/34	-
7	FES	N	244	2	-	-	0/1/1/1
8	F3S	N	245	2	-	-	0/3/3/3
9	SF4	N	246	2	-	-	0/6/5/5
5	OAA	M	802	-	-	4/8/8/8	-
8	F3S	B	245	2	-	-	0/3/3/3

The worst 5 of 51 bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
6	A	703	FAD	PA-O3P	-9.33	1.49	1.59
6	A	703	FAD	P-O3P	-9.07	1.49	1.59
10	N	800	BRS	C12-N1	6.60	1.57	1.45
6	M	803	FAD	PA-O3P	-5.79	1.53	1.59
10	B	700	BRS	C17-C16	5.37	1.47	1.38

The worst 5 of 79 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
6	A	703	FAD	N3A-C2A-N1A	-5.36	120.47	128.58
6	M	803	FAD	N3A-C2A-N1A	-5.34	120.50	128.58
6	A	703	FAD	N3A-C4A-N9A	4.00	133.98	127.17
6	M	803	FAD	C4'-C3'-C2'	-3.81	107.22	113.57
6	M	803	FAD	N3A-C4A-N9A	3.71	133.48	127.17

There are no chirality outliers.

5 of 79 torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
5	A	702	OAA	O3-C3-C4-O4
5	A	702	OAA	O3-C3-C4-O5
5	A	702	OAA	C2-C3-C4-O5
5	M	802	OAA	O3-C3-C4-O4
5	M	802	OAA	O3-C3-C4-O5


There are no ring outliers.


11 monomers are involved in 56 short contacts:


Mol	Chain	Res	Type	Clashes	Symm-Clashes
10	N	800	BRS	5	0
5	A	702	OAA	1	0
11	O	811	CE1	1	0
10	B	700	BRS	13	0
11	D	710	CE1	5	0
6	A	703	FAD	7	0
6	M	803	FAD	11	0
11	P	810	CE1	2	0
8	N	245	F3S	2	0
9	N	246	SF4	7	0
5	M	802	OAA	2	0


The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient

equivalents in the CSD to analyse the geometry.









5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

EDS was not executed - this section is therefore empty.

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

EDS was not executed - this section is therefore empty.

6.3 Carbohydrates [i](#)

EDS was not executed - this section is therefore empty.

6.4 Ligands [i](#)

EDS was not executed - this section is therefore empty.

6.5 Other polymers [i](#)

EDS was not executed - this section is therefore empty.