Full wwPDB NMR Structure Validation Report

Mar 26, 2026 – 02:09 AM UTC

PDB ID	:	2KNG / pdb_00002kng
Title	:	Solution structure of C-domain of Lsr2
Authors	:	Li, Y.; Xia, B.
Deposited on	:	2009-08-22

This is a Full wwPDB NMR Structure Validation Report for a publicly released PDB entry.

We welcome your comments at *validation@mail.wwpdb.org*
A user guide is available at
https://www.wwpdb.org/validation/2017/NMRValidationReportHelp
with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at
http://www.wwpdb.org/validation/2017/FAQs#types.

The following versions of software and data (see references ⓘ) were used in the production of this report:


MolProbity	:	4-5-2 with Phenix2.0
Percentile statistics	:	20250101.v01 (using entries in the PDB archive January 1st 2025)
wwPDB-RCI	:	v_1n_11_5_13_A (Berjanski et al., 2005)
PANAV	:	Wang et al. (2010)
wwPDB-ShiftChecker	:	v1.2
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.49

1 Overall quality at a glance (i)

The following experimental techniques were used to determine the structure:
SOLUTION NMR

The overall completeness of chemical shifts assignment was not calculated.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	NMR archive (#Entries)
Clashscore	229148	14424
Ramachandran outliers	224038	12848
Sidechain outliers	223484	12823

The table below summarises the geometric issues observed across the polymeric chains and their fit to the experimental data. The red, orange, yellow and green segments indicate the fraction of residues that contain outliers for >=3, 2, 1 and 0 types of geometric quality criteria. A cyan segment indicates the fraction of residues that are not part of the well-defined cores, and a grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions <=5%

Mol	Chain	Length	Quality of chain
1	A	55	65% 35%

2 Ensemble composition and analysis (i)

This entry contains 20 models. Model 4 is the overall representative, medoid model (most similar to other models). The authors have identified model 1 as representative, based on the following criterion: *lowest energy*.

The following residues are included in the computation of the global validation metrics.

Well-defined (core) protein residues			
Well-defined core	**Residue range (total)**	**Backbone RMSD (Å)**	**Medoid model**
1	A:79-A:114 (36)	0.62	4

Ill-defined regions of proteins are excluded from the global statistics.

Ligands and non-protein polymers are included in the analysis.

The models can be grouped into 2 clusters and 2 single-model clusters were found.

Cluster number	Models
1	1, 2, 4, 6, 10, 11, 16, 17, 18
2	3, 5, 7, 8, 9, 12, 13, 14, 15
Single-model clusters	19; 20

3 Entry composition (i)

There is only 1 type of molecule in this entry. The entry contains 845 atoms, of which 411 are hydrogens and 0 are deuteriums.

- Molecule 1 is a protein called Protein lsr2.

Mol	Chain	Residues	Atoms					Trace
1	A	55	Total	C	H	N	O	0
			845	257	411	99	78	

There are 8 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	113	LEU	-	expression tag	UNP P65648
A	114	GLU	-	expression tag	UNP P65648
A	115	HIS	-	expression tag	UNP P65648
A	116	HIS	-	expression tag	UNP P65648
A	117	HIS	-	expression tag	UNP P65648
A	118	HIS	-	expression tag	UNP P65648
A	119	HIS	-	expression tag	UNP P65648
A	120	HIS	-	expression tag	UNP P65648


4 Residue-property plots

4.1 Average score per residue in the NMR ensemble

These plots are provided for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic is the same as shown in the summary in section 1 of this report. The second graphic shows the sequence where residues are colour-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. Stretches of 2 or more consecutive residues without any outliers are shown as green connectors. Residues which are classified as ill-defined in the NMR ensemble, are shown in cyan with an underline colour-coded according to the previous scheme. Residues which were present in the experimental sample, but not modelled in the final structure are shown in grey.

• Molecule 1: Protein lsr2

Chain A: 




4.2 Scores per residue for each member of the ensemble

Colouring as in section 4.1 above.

4.2.1 Score per residue for model 1

• Molecule 1: Protein lsr2


Chain A: 



4.2.2 Score per residue for model 2

• Molecule 1: Protein lsr2

Chain A: 



4.2.3 Score per residue for model 3

• Molecule 1: Protein lsr2

Chain A:



4.2.4 Score per residue for model 4 (medoid)

• Molecule 1: Protein lsr2

Chain A:



4.2.5 Score per residue for model 5

• Molecule 1: Protein lsr2

Chain A:



4.2.6 Score per residue for model 6

• Molecule 1: Protein lsr2

Chain A:



4.2.7 Score per residue for model 7

• Molecule 1: Protein lsr2

Chain A:



4.2.8 Score per residue for model 8

• Molecule 1: Protein lsr2

Chain A:



4.2.9 Score per residue for model 9

• Molecule 1: Protein lsr2

Chain A:



4.2.10 Score per residue for model 10

• Molecule 1: Protein lsr2

Chain A:



4.2.11 Score per residue for model 11

• Molecule 1: Protein lsr2

Chain A:



4.2.12 Score per residue for model 12

• Molecule 1: Protein lsr2

Chain A:



4.2.13 Score per residue for model 13

• Molecule 1: Protein lsr2

Chain A:



4.2.14 Score per residue for model 14

• Molecule 1: Protein lsr2

Chain A:



4.2.15 Score per residue for model 15

• Molecule 1: Protein lsr2

Chain A:



4.2.16 Score per residue for model 16

• Molecule 1: Protein lsr2

Chain A:



4.2.17 Score per residue for model 17

• Molecule 1: Protein lsr2

Chain A:



4.2.18 Score per residue for model 18

• Molecule 1: Protein lsr2

Chain A:



4.2.19 Score per residue for model 19

• Molecule 1: Protein lsr2

Chain A:



4.2.20 Score per residue for model 20

• Molecule 1: Protein lsr2

Chain A:



5 Refinement protocol and experimental data overview

The models were refined using the following method: *simulated annealing.*

Of the 100 calculated structures, 20 were deposited, based on the following criterion: *structures with the lowest energy.*

The following table shows the software used for structure solution, optimisation and refinement.

Software name	Classification	Version
Amber	refinement	9

No chemical shift data was provided.

6 Model quality

6.1 Standard geometry

There are no covalent bond-length or bond-angle outliers.

There are no bond-length outliers.

There are no bond-angle outliers.

There are no chirality outliers.

There are no planarity outliers.

6.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in each chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes averaged over the ensemble.

Mol	Chain	Non-H	H(model)	H(added)	Clashes
All	All	5640	5480	5440	-

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is -.

There are no clashes.

6.3 Torsion angles

6.3.1 Protein backbone

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all PDB entries followed by that with respect to all NMR entries. The Analysed column shows the number of residues for which the backbone conformation was analysed and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	36/55 (65%)	36±1 (99±2%)	0±1 (1±2%)	0±0 (0±0%)	100	100
All	All	720/1100 (65%)	710 (99%)	10 (1%)	0 (0%)	100	100

There are no Ramachandran outliers.

6.3.2 Protein sidechains

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all PDB entries followed by that with respect to all NMR entries. The Analysed column shows the number of residues for which the sidechain conformation was analysed and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	27/41 (66%)	27±1 (99±2%)	0±1 (1±2%)	59	94
All	All	540/820 (66%)	533 (99%)	7 (1%)	59	94

All 3 unique residues with a non-rotameric sidechain are listed below. They are sorted by the frequency of occurrence in the ensemble.

Mol	Chain	Res	Type	Models (Total)
1	A	104	VAL	4
1	A	114	GLU	2
1	A	99	ARG	1

6.3.3 RNA

There are no RNA molecules in this entry.

6.4 Non-standard residues in protein, DNA, RNA chains

There are no non-standard protein/DNA/RNA residues in this entry.

6.5 Carbohydrates

There are no oligosaccharides in this entry.

6.6 Ligand geometry

There are no ligands in this entry.

6.7 Other polymers

There are no such molecules in this entry.

6.8 Polymer linkage issues

There are no chain breaks in this entry.

7 Chemical shift validation

No chemical shift data were provided