wwPDB X-ray Structure Validation Summary Report (i)

Aug 5, 2026 – 04:10 PM JST

PDB ID	:	6KY7 / pdb_00006ky7
Title	:	Crystal structure of yak lactoperoxidase at 2.27 A resolution
Authors	:	Singh, P.K.; Viswanathan, V.; Sharma, P.; Rani, C.; Sharma, S.; Singh, T.P.
Deposited on	:	2019-09-16
Resolution	:	2.27 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at *validation@mail.wwpdb.org*
A user guide is available at
https://www.wwpdb.org/validation/2017/XrayValidationReportHelp
with specific help available everywhere you see the (i) symbol.

The types of validation reports are described at
http://www.wwpdb.org/validation/2017/FAQs#types.

The following versions of software and data (see references (i)) were used in the production of this report:

MolProbity	:	4-5-2 with Phenix2.0
Mogul	:	1.8.5 (274361), CSD as541be (2020)
Xtriage (Phenix)	:	2.0
EDS	:	3.0
Buster-report	:	wwPDB partial adaption of 1.1.7 (2018)
Percentile statistics	:	20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4	:	9.0.015 (Gargrove)
Density-Fitness	:	1.0.12
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.50

1 Overall quality at a glance (i)

The following experimental techniques were used to determine the structure:
X-RAY DIFFRACTION

The reported resolution of this entry is 2.27 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	9078 (2.30-2.26)
Clashscore	190562	9802 (2.30-2.26)
Ramachandran outliers	187476	9690 (2.30-2.26)
Sidechain outliers	187428	9691 (2.30-2.26)
RSRZ outliers	180081	9085 (2.30-2.26)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for >=3, 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions <=5% The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	595	4% 71% 23% 6%

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
6	OSM	A	620	-	-	X	-
6	OSM	A	622	-	-	X	-

Continued on next page...

Continued from previous page...

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
8	PEO	A	628	-	-	X	-

2 Entry composition

There are 9 unique types of molecules in this entry. The entry contains 5264 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Lactoperoxidase.

Mol	Chain	Residues	Atoms						ZeroOcc	AltConf	Trace
1	A	595	Total	C	N	O	P	S	0	0	0
			4763	3025	847	864	1	26			

There are 2 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	161	SER	PHE	conflict	UNP L8ICE9
A	254	ALA	PHE	conflict	UNP L8ICE9

- Molecule 2 is 2-acetamido-2-deoxy-beta-D-glucopyranose (CCD ID: NAG) (formula: $C_8H_{15}NO_6$) (labeled as "Ligand of Interest" by depositor).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
2	A	1	Total	C	N	O	0	0
			14	8	1	5		
2	A	1	Total	C	N	O	0	0
			14	8	1	5		

Continued on next page...

Continued from previous page...

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
2	A	1	Total C N O 14 8 1 5	0	0
2	A	1	Total C N O 14 8 1 5	0	0

- Molecule 3 is CALCIUM ION (CCD ID: CA) (formula: Ca).

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
3	A	1	Total Ca 1 1	0	0

- Molecule 4 is PROTOPORPHYRIN IX CONTAINING FE (CCD ID: HEM) (formula: $C_{34}H_{32}FeN_4O_4$) (labeled as "Ligand of Interest" by depositor).



Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
4	A	1	Total C Fe N O 43 34 1 4 4	0	0

- Molecule 5 is IODIDE ION (CCD ID: IOD) (formula: I).

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
5	A	13	Total I 13 13	0	0

- Molecule 6 is 1-(OXIDOSULFANYL)METHANAMINE (CCD ID: OSM) (formula: CH_5NOS) (labeled as "Ligand of Interest" by depositor).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
6	A	1	Total	C	N	O	S	0	0
			4	1	1	1	1		
6	A	1	Total	C	N	O	S	0	0
			4	1	1	1	1		
6	A	1	Total	C	N	O	S	0	0
			4	1	1	1	1		
6	A	1	Total	C	N	O	S	0	0
			4	1	1	1	1		

- Molecule 7 is THIOCYANATE ION (CCD ID: SCN) (formula: CNS).



Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
7	A	1	Total C N S 3 1 1 1	0	0
7	A	1	Total C N S 3 1 1 1	0	0
7	A	1	Total C N S 3 1 1 1	0	0
7	A	1	Total C N S 3 1 1 1	0	0

- Molecule 8 is HYDROGEN PEROXIDE (CCD ID: PEO) (formula: H_2O_2).



Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
8	A	1	Total O 2 2	0	0

- Molecule 9 is water.

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
9	A	358	Total O 358 358	0	0

3 Residue-property plots (i)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density (RSRZ > 2). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: Lactoperoxidase

Chain A:



4 Data and refinement statistics (i)

Property	Value	Source
Space group	P 1 21 1	Depositor
Cell constants a, b, c, α, β, γ	54.08Å 80.64Å 76.13Å 90.00° 102.80° 90.00°	Depositor
Resolution (Å)	52.79 – 2.27 52.79 – 2.27	Depositor EDS
% Data completeness (in resolution range)	100.0 (52.79-2.27) 100.0 (52.79-2.27)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	0.08	Depositor
$< I/\sigma(I) >$ [1]	1.52 (at 2.27Å)	Xtriage
Refinement program	REFMAC 5.8.0257	Depositor
R, R_{free}	0.196 , 0.266 0.209 , 0.269	Depositor DCC
R_{free} test set	1020 reflections (3.45%)	wwPDB-VP
Wilson B-factor (Å^2)	38.0	Xtriage
Anisotropy	0.685	Xtriage
Bulk solvent k_{sol}(e/Å^3), B_{sol}(Å^2)	0.34 , 54.2	EDS
L-test for twinning[2]	$< \|L\| > = 0.49$, $< L^2 > = 0.33$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o,F_c correlation	0.95	EDS
Total number of atoms	5264	wwPDB-VP
Average B, all atoms (Å^2)	58.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 5.59% of the height of the origin peak. No significant pseudotranslation is detected.*

[1] Intensities estimated from amplitudes.

[2] Theoretical values of $< |L| >$, $< L^2 >$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: CA, OSM, NAG, SEP, HEM, SCN, PEO, IOD

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	#$\|Z\|>5$	RMSZ	#$\|Z\|>5$
1	A	1.01	0/4878	1.41	6/6617 (0.1%)

There are no bond length outliers.

The worst 5 of 6 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(^{o})	Ideal(^{o})
1	A	170	PRO	N-CA-C	8.13	120.62	110.70
1	A	194	GLY	CA-C-O	-7.40	117.13	122.23
1	A	453	ARG	CA-C-N	5.14	125.80	119.99
1	A	453	ARG	C-N-CA	5.14	125.80	119.99
1	A	362	ASP	CA-CB-CG	5.12	117.72	112.60

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	4763	0	4679	136	0
2	A	56	0	52	0	0
3	A	1	0	0	0	0
4	A	43	0	30	4	0
5	A	13	0	0	3	0
6	A	16	0	20	5	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
7	A	12	0	0	2	0
8	A	2	0	0	2	0
9	A	358	0	0	22	0
All	All	5264	0	4781	141	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 15.

The worst 5 of 141 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
5:A:608:IOD:I	7:A:626:SCN:N	2.56	1.09
1:A:424:PRO:O	5:A:615:IOD:I	2.48	1.00
1:A:167:CYS:CB	1:A:168:PRO:HD2	1.96	0.96
1:A:360:ARG:NH2	1:A:389:ASP:OD2	1.99	0.95
1:A:167:CYS:HB2	1:A:168:PRO:HD2	1.48	0.94

There are no symmetry-related clashes.

5.3 Torsion angles (i)

5.3.1 Protein backbone (i)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
1	A	592/595 (100%)	552 (93%)	31 (5%)	9 (2%)	8 7

5 of 9 Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	64	ARG
1	A	167	CYS
1	A	170	PRO
1	A	581	THR

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	A	583	ASP

5.3.2 Protein sidechains (i)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles
1	A	516/516 (100%)	468 (91%)	48 (9%)	8 10

5 of 48 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	317	LEU
1	A	407	MET
1	A	322	GLN
1	A	363	GLU
1	A	464	LEU

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 16 such sidechains are listed below:

Mol	Chain	Res	Type
1	A	545	GLN
1	A	520	GLN
1	A	333	ASN
1	A	497	ASN
1	A	322	GLN

5.3.3 RNA (i)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains (i)

1 non-standard protein/DNA/RNA residue is modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\#\|Z\| > 2$	Counts	RMSZ	$\#\|Z\| > 2$
1	SEP	A	198	1	8,9,10	1.13	0	8,12,14	1.24	1 (12%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
1	SEP	A	198	1	-	3/5/8/10	-

There are no bond length outliers.

All (1) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(^{o})	Ideal(^{o})
1	A	198	SEP	P-OG-CB	2.28	124.58	118.30

There are no chirality outliers.

All (3) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
1	A	198	SEP	CB-OG-P-O3P
1	A	198	SEP	CB-OG-P-O1P
1	A	198	SEP	CB-OG-P-O2P

There are no ring outliers.

1 monomer is involved in 2 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
1	A	198	SEP	2	0

5.5 Carbohydrates (i)

There are no oligosaccharides in this entry.

5.6 Ligand geometry (i)

Of 28 ligands modelled in this entry, 14 are monoatomic - leaving 14 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\#\|Z\| > 2$	Counts	RMSZ	$\#\|Z\| > 2$
2	NAG	A	601	1	14,14,15	0.71	0	17,19,21	1.17	2 (11%)
7	SCN	A	627	-	1,2,2	0.68	0	0,1,1	-	-
6	OSM	A	623	-	1,3,3	0.08	0	0,2,2	-	-
6	OSM	A	620	-	1,3,3	0.01	0	0,2,2	-	-
6	OSM	A	622	-	1,3,3	0.05	0	0,2,2	-	-
7	SCN	A	624	-	1,2,2	0.02	0	0,1,1	-	-
7	SCN	A	626	-	1,2,2	1.03	0	0,1,1	-	-
2	NAG	A	604	1	14,14,15	0.44	0	17,19,21	1.11	2 (11%)
4	HEM	A	606	8,1	50,50,50	1.64	11 (22%)	66,82,82	1.70	12 (18%)
2	NAG	A	603	1	14,14,15	1.04	1 (7%)	17,19,21	1.53	1 (5%)
7	SCN	A	625	-	1,2,2	0.73	0	0,1,1	-	-
2	NAG	A	602	1	14,14,15	0.29	0	17,19,21	1.11	1 (5%)
6	OSM	A	621	-	1,3,3	0.09	0	0,2,2	-	-
8	PEO	A	628	4	1,1,1	0.49	0	-		

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	NAG	A	601	1	-	0/6/23/26	0/1/1/1
6	OSM	A	623	-	-	0/0/1/1	-
6	OSM	A	620	-	-	0/0/1/1	-
6	OSM	A	622	-	-	0/0/1/1	-

Continued on next page...

Continued from previous page...

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	NAG	A	604	1	-	2/6/23/26	0/1/1/1
4	HEM	A	606	8,1	-	4/14/54/54	-
2	NAG	A	603	1	-	2/6/23/26	0/1/1/1
2	NAG	A	602	1	-	2/6/23/26	0/1/1/1
6	OSM	A	621	-	-	0/0/1/1	-

The worst 5 of 12 bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
4	A	606	HEM	FE-NB	5.15	2.10	1.94
4	A	606	HEM	FE-NC	3.82	2.08	1.95
4	A	606	HEM	C4D-ND	-3.75	1.33	1.40
2	A	603	NAG	O5-C1	-3.00	1.38	1.43
4	A	606	HEM	C1B-NB	-2.82	1.35	1.40

The worst 5 of 18 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(^{o})	Ideal(^{o})
2	A	603	NAG	C1-O5-C5	-5.14	105.23	112.19
4	A	606	HEM	CHA-C4D-ND	4.67	130.13	124.37
4	A	606	HEM	CHC-C4B-NB	4.33	129.13	124.42
4	A	606	HEM	CMD-C2D-C1D	4.14	131.35	125.04
4	A	606	HEM	C1B-NB-C4B	3.78	108.97	105.07

There are no chirality outliers.

5 of 10 torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
2	A	604	NAG	O5-C5-C6-O6
2	A	604	NAG	C4-C5-C6-O6
2	A	603	NAG	C4-C5-C6-O6
2	A	603	NAG	O5-C5-C6-O6
2	A	602	NAG	C4-C5-C6-O6

There are no ring outliers.

7 monomers are involved in 11 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
7	A	627	SCN	1	0
6	A	620	OSM	2	0

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Clashes	Symm-Clashes
6	A	622	OSM	2	0
7	A	626	SCN	1	0
4	A	606	HEM	4	0
6	A	621	OSM	1	0
8	A	628	PEO	2	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less then 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.

Ligand NAG A 601



Ligand OSM A 623



Bond lengths



Bond angles



Torsions



Rings

Ligand OSM A 620



Bond lengths



Bond angles



Torsions



Rings

Ligand OSM A 622



Bond lengths



Bond angles



Torsions



Rings

Ligand NAG A 604



Bond lengths



Bond angles



Torsions



Rings

Ligand HEM A 606



Bond lengths



Bond angles



Torsions



Rings

Ligand NAG A 603



Bond lengths



Bond angles



Torsions



Rings

Ligand NAG A 602



Bond lengths



Bond angles



Torsions



Rings

Ligand OSM A 621



5.7 Other polymers (i)

There are no such residues in this entry.

5.8 Polymer linkage issues (i)

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

In the following table, the column labelled '#RSRZ> 2' contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95^{th} percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled 'Q< 0.9' lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å^2)	Q<0.9
1	A	594/595 (99%)	0.29	22 (3%) 45 47	31, 52, 112, 177	0

The worst 5 of 22 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	174	SER	4.2
1	A	175	LEU	3.8
1	A	2	TRP	3.6
1	A	171	PRO	3.6
1	A	169	THR	3.4

6.2 Non-standard residues in protein, DNA, RNA chains

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95^{th} percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å^2)	Q<0.9
1	SEP	A	198	10/11	0.92	0.12	37,52,62,66	0

6.3 Carbohydrates

There are no oligosaccharides in this entry.

6.4 Ligands

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum,

median, 95^{th} percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å^2)	Q<0.9
2	NAG	A	601	14/15	0.69	0.14	75,86,98,101	0
2	NAG	A	603	14/15	0.81	0.12	52,62,67,80	0
2	NAG	A	604	14/15	0.82	0.10	77,89,96,96	0
2	NAG	A	602	14/15	0.85	0.13	72,78,94,99	0
6	OSM	A	621	4/4	0.87	0.12	54,56,60,60	0
8	PEO	A	628	2/2	0.95	0.10	42,42,42,60	0
5	IOD	A	618	1/1	0.96	0.09	76,76,76,76	1
6	OSM	A	620	4/4	0.96	0.11	30,31,33,36	0
5	IOD	A	614	1/1	0.96	0.07	81,81,81,81	0
6	OSM	A	622	4/4	0.96	0.06	49,50,50,51	0
7	SCN	A	625	3/3	0.96	0.08	56,56,58,62	0
5	IOD	A	616	1/1	0.96	0.09	74,74,74,74	1
5	IOD	A	617	1/1	0.97	0.06	74,74,74,74	1
4	HEM	A	606	43/43	0.97	0.07	28,37,42,48	0
6	OSM	A	623	4/4	0.97	0.12	26,28,34,34	0
7	SCN	A	624	3/3	0.97	0.07	44,44,52,58	0
5	IOD	A	619	1/1	0.97	0.12	81,81,81,81	1
7	SCN	A	627	3/3	0.97	0.08	44,44,50,60	0
5	IOD	A	611	1/1	0.97	0.05	65,65,65,65	1
5	IOD	A	608	1/1	0.98	0.12	67,67,67,67	1
5	IOD	A	609	1/1	0.98	0.04	60,60,60,60	0
5	IOD	A	610	1/1	0.98	0.05	59,59,59,59	1
3	CA	A	605	1/1	0.98	0.03	46,46,46,46	0
7	SCN	A	626	3/3	0.98	0.07	30,30,34,35	3
5	IOD	A	607	1/1	0.98	0.04	51,51,51,51	1
5	IOD	A	615	1/1	0.98	0.05	56,56,56,56	0
5	IOD	A	613	1/1	0.99	0.04	44,44,44,44	0
5	IOD	A	612	1/1	0.99	0.03	69,69,69,69	0

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.

Electron density around NAG A 601:

$2mF_o$-DF_c (at 0.7 rmsd) in gray
mF_o-DF_c (at 3 rmsd) in purple (negative) and green (positive)







Electron density around NAG A 603:

$2mF_o$-DF_c (at 0.7 rmsd) in gray
mF_o-DF_c (at 3 rmsd) in purple (negative) and green (positive)







Electron density around NAG A 604:

$2mF_o$-DF_c (at 0.7 rmsd) in gray
mF_o-DF_c (at 3 rmsd) in purple (negative) and green (positive)







Electron density around NAG A 602:

$2mF_o$-DF_c (at 0.7 rmsd) in gray
mF_o-DF_c (at 3 rmsd) in purple (negative) and green (positive)







Electron density around OSM A 621:

$2mF_o$-DF_c (at 0.7 rmsd) in gray
mF_o-DF_c (at 3 rmsd) in purple (negative) and green (positive)







Electron density around OSM A 620:

$2mF_o$-DF_c (at 0.7 rmsd) in gray
mF_o-DF_c (at 3 rmsd) in purple (negative) and green (positive)







Electron density around OSM A 622:

$2mF_o$-DF_c (at 0.7 rmsd) in gray
mF_o-DF_c (at 3 rmsd) in purple (negative) and green (positive)







Electron density around HEM A 606:

$2mF_o$-DF_c (at 0.7 rmsd) in gray
mF_o-DF_c (at 3 rmsd) in purple (negative) and green (positive)







Electron density around OSM A 623:

$2mF_o$-DF_c (at 0.7 rmsd) in gray
mF_o-DF_c (at 3 rmsd) in purple (negative) and green (positive)



6.5 Other polymers (i)

There are no such residues in this entry.