



# Full wwPDB X-ray Structure Validation Report ⓘ

Mar 6, 2026 – 05:31 PM UTC

PDB ID : 1LOO / pdb\_00001loo  
Title : Crystal Structure of the Mouse-Muscle Adenylosuccinate Synthetase Ligated with GTP  
Authors : Iancu, C.V.; Borza, T.; Fromm, H.J.; Honzatko, R.B.  
Deposited on : 2002-05-06  
Resolution : 2.20 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4-5-2 with Phenix2.0  
Mogul : 2022.3.0, CSD as543be (2022)  
Xtrriage (Phenix) : 2.0  
EDS : 3.0  
Buster-report : wwPDB partial adaption of 1.1.7 (2018)  
Percentile statistics : 20250101.v01 (using entries in the PDB archive January 1st 2025)  
CCP4 : 9.0.010 (Gargrove)  
Density-Fitness : 1.0.12  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : 2.49


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

*X-RAY DIFFRACTION*

The reported resolution of this entry is 2.20 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
Clashscore	190562	6851 (2.20-2.20)
Ramachandran outliers	187476	6768 (2.20-2.20)
Sidechain outliers	187428	6769 (2.20-2.20)
RSRZ outliers	180081	6166 (2.20-2.20)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	457	

## 2 Entry composition [i](#)


There are 3 unique types of molecules in this entry. The entry contains 3485 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called adenylosuccinate synthetase.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	431	3353	2133	589	619	12	0	0	0

- Molecule 2 is GUANOSINE-5'-TRIPHOSPHATE (CCD ID: GTP) (formula: C<sub>10</sub>H<sub>16</sub>N<sub>5</sub>O<sub>14</sub>P<sub>3</sub>).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
			Total	C	N	O	P		
2	A	1	32	10	5	14	3	0	0


- Molecule 3 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
			Total	O		
3	A	100	100	100	0	0

### 3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: adenylosuccinate synthetase



## 4 Data and refinement statistics

Property	Value	Source
Space group	P 43 21 2	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	69.94Å 69.94Å 198.09Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	10.00 – 2.20 10.00 – 2.20	Depositor EDS
% Data completeness (in resolution range)	(Not available) (10.00-2.20) 98.1 (10.00-2.20)	Depositor EDS
$R_{merge}$	0.04	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	2.17 (at 2.20Å)	Xtrriage
Refinement program	CNS	Depositor
R, $R_{free}$	0.244 , 0.276 0.249 , (Not available)	Depositor DCC
$R_{free}$ test set	No test flags present.	wwPDB-VP
Wilson B-factor (Å <sup>2</sup> )	48.1	Xtrriage
Anisotropy	0.016	Xtrriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.42 , 49.2	EDS
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.50$ , $\langle L^2 \rangle = 0.33$	Xtrriage
Estimated twinning fraction	No twinning to report.	Xtrriage
$F_o, F_c$ correlation	0.94	EDS
Total number of atoms	3485	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	48.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 6.66% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

## 5 Model quality [i](#)

### 5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: GTP

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z  >5	RMSZ	# Z  >5
1	A	0.40	0/3423	0.88	7/4628 (0.2%)

There are no bond length outliers.

All (7) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	104	ILE	N-CA-C	7.04	117.97	108.11
1	A	277	VAL	N-CA-C	5.99	116.72	110.62
1	A	245	GLY	CA-C-N	-5.95	115.37	119.66
1	A	245	GLY	C-N-CA	-5.95	115.37	119.66
1	A	264	PHE	N-CA-C	5.23	119.81	113.38
1	A	61	SER	N-CA-C	5.20	116.89	109.14
1	A	234	ASP	N-CA-C	-5.02	101.92	109.15

There are no chirality outliers.

There are no planarity outliers.

### 5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	3353	0	3359	100	0
2	A	32	0	12	0	0
3	A	100	0	0	1	0
All	All	3485	0	3371	100	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 15.

All (100) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:178:THR:HG23	1:A:199:LEU:HD13	1.59	0.84
1:A:163:THR:HG22	1:A:164:LYS:H	1.47	0.79
1:A:185:LEU:HD22	1:A:225:ALA:HB2	1.71	0.72
1:A:233:ARG:HG3	1:A:233:ARG:HH11	1.54	0.72
1:A:86:PRO:HD3	1:A:98:ILE:HD11	1.72	0.72
1:A:105:HIS:CE1	1:A:107:PRO:HG2	2.27	0.70
1:A:95:VAL:HG11	1:A:129:ILE:HD12	1.74	0.69
1:A:262:ILE:HD13	1:A:262:ILE:O	1.95	0.66
1:A:229:ARG:HH21	1:A:229:ARG:HB2	1.60	0.66
1:A:62:ARG:HG2	1:A:62:ARG:HH11	1.61	0.65
1:A:163:THR:HG22	1:A:164:LYS:N	2.11	0.65
1:A:150:VAL:HA	1:A:154:ALA:HB2	1.78	0.65
1:A:229:ARG:HB2	1:A:229:ARG:NH2	2.11	0.64
1:A:193:SER:HB3	1:A:218:LEU:HD21	1.80	0.63
1:A:155:GLN:OE1	1:A:156:GLU:HG2	2.00	0.61
1:A:177:ARG:HH11	1:A:177:ARG:HG2	1.66	0.60
1:A:95:VAL:CG1	1:A:129:ILE:HD12	2.30	0.60
1:A:75:VAL:HG11	1:A:90:ILE:HG21	1.84	0.60
1:A:153:GLN:HA	1:A:157:GLY:HA3	1.84	0.60
1:A:75:VAL:HG23	1:A:78:LYS:NZ	2.18	0.59
1:A:117:GLU:HA	1:A:121:LEU:O	2.03	0.59
1:A:31:SER:O	1:A:243:LEU:HD22	2.03	0.58
1:A:129:ILE:HD13	1:A:238:PHE:CE2	2.37	0.58
1:A:78:LYS:NZ	1:A:78:LYS:HB2	2.19	0.57
1:A:331:VAL:HG13	1:A:332:THR:HG23	1.86	0.57
1:A:351:HIS:HD2	1:A:356:PHE:H	1.50	0.57
1:A:145:ASP:HB2	1:A:169:PRO:HG2	1.87	0.56
1:A:115:LYS:HD2	1:A:115:LYS:C	2.31	0.56
1:A:122:LYS:HD3	1:A:123:ASP:H	1.68	0.56
1:A:233:ARG:HG3	1:A:233:ARG:NH1	2.20	0.56
1:A:185:LEU:CD2	1:A:225:ALA:HB2	2.36	0.55
1:A:349:TYR:HA	1:A:352:MET:CE	2.37	0.55
1:A:122:LYS:HD3	1:A:123:ASP:N	2.22	0.55
1:A:317:GLU:H	1:A:317:GLU:CD	2.14	0.54
1:A:123:ASP:HA	1:A:125:GLU:OE2	2.06	0.54
1:A:189:PHE:CZ	1:A:222:LYS:HB2	2.41	0.54
1:A:442:LYS:HD3	1:A:443:TRP:CH2	2.43	0.54

*Continued on next page...*

*Continued from previous page...*

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:178:THR:HG23	1:A:199:LEU:CD1	2.34	0.54
1:A:198:ASN:O	1:A:202:GLN:HG3	2.07	0.54
1:A:92:THR:HG22	1:A:127:ARG:NH2	2.23	0.53
1:A:144:VAL:O	1:A:148:GLN:HG3	2.09	0.53
1:A:143:ALA:O	1:A:147:LEU:HD13	2.10	0.52
1:A:380:LYS:HE2	1:A:383:GLY:O	2.09	0.52
1:A:204:GLN:HG3	1:A:212:ILE:HD12	1.92	0.52
1:A:229:ARG:HH21	1:A:229:ARG:CB	2.22	0.51
1:A:323:GLN:NE2	1:A:336:LYS:HB3	2.25	0.51
1:A:210:LEU:HD12	1:A:212:ILE:HG13	1.92	0.51
1:A:349:TYR:HA	1:A:352:MET:HE2	1.91	0.51
1:A:66:GLY:HA2	1:A:273:SER:OG	2.11	0.51
1:A:237:TYR:CE1	1:A:241:GLU:HG3	2.45	0.51
1:A:75:VAL:HG23	1:A:78:LYS:HZ2	1.77	0.50
1:A:442:LYS:HD3	1:A:443:TRP:CZ3	2.47	0.50
1:A:262:ILE:HD13	1:A:262:ILE:C	2.37	0.50
1:A:323:GLN:OE1	1:A:336:LYS:HD3	2.12	0.49
1:A:153:GLN:HB3	1:A:158:LYS:O	2.12	0.49
1:A:80:TYR:OH	1:A:120:GLY:HA3	2.13	0.48
1:A:321:LEU:HD23	1:A:370:LEU:HD21	1.95	0.48
1:A:144:VAL:HG21	1:A:203:HIS:HD2	1.77	0.48
1:A:200:ALA:HB1	1:A:212:ILE:HD13	1.93	0.48
1:A:434:GLU:HG2	1:A:441:VAL:HG23	1.96	0.47
1:A:404:GLU:HG2	1:A:432:PHE:CZ	2.50	0.47
1:A:351:HIS:CD2	1:A:356:PHE:H	2.31	0.47
1:A:106:LEU:CD2	1:A:136:LEU:HD22	2.45	0.47
1:A:60:VAL:HG11	1:A:88:GLY:HA3	1.96	0.46
1:A:276:THR:HB	1:A:354:ASN:OD1	2.15	0.46
1:A:160:ILE:HG22	1:A:161:GLY:N	2.29	0.46
1:A:150:VAL:HG12	1:A:150:VAL:O	2.16	0.45
1:A:196:PHE:HZ	1:A:217:GLN:HE21	1.64	0.45
1:A:75:VAL:HG11	1:A:90:ILE:CG2	2.46	0.45
1:A:193:SER:CB	1:A:218:LEU:HD21	2.45	0.45
1:A:43:ASP:O	1:A:299:LYS:HE2	2.17	0.45
1:A:259:LEU:HD11	1:A:346:ILE:HG12	1.99	0.45
1:A:341:TRP:CE2	1:A:374:LYS:HB2	2.52	0.45
1:A:113:ALA:HB1	1:A:124:TRP:CZ2	2.52	0.44
1:A:146:GLY:O	1:A:150:VAL:HG23	2.16	0.44
1:A:92:THR:HA	1:A:127:ARG:HE	1.83	0.44
1:A:38:GLY:N	1:A:46:LYS:HD3	2.33	0.44
1:A:67:ASN:HB3	1:A:256:ASN:HD21	1.84	0.43

*Continued on next page...*

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:67:ASN:CB	1:A:256:ASN:HD21	2.31	0.43
1:A:78:LYS:N	1:A:78:LYS:HD3	2.34	0.43
1:A:319:GLY:O	1:A:323:GLN:HB2	2.19	0.43
1:A:163:THR:CG2	1:A:164:LYS:H	2.23	0.43
1:A:160:ILE:HG23	1:A:268:PRO:O	2.18	0.43
1:A:132:ASP:HB3	1:A:232:VAL:CG1	2.49	0.42
1:A:62:ARG:HG2	1:A:62:ARG:NH1	2.29	0.42
1:A:210:LEU:HD13	1:A:211:GLU:N	2.35	0.42
1:A:41:TRP:NE1	1:A:304:ARG:NH2	2.68	0.42
1:A:98:ILE:HG21	1:A:104:ILE:HD11	2.02	0.42
1:A:443:TRP:HA	1:A:454:ILE:O	2.20	0.42
1:A:93:LYS:NZ	1:A:93:LYS:HB3	2.34	0.41
1:A:125:GLU:H	1:A:125:GLU:CD	2.28	0.41
1:A:351:HIS:CD2	1:A:356:PHE:N	2.88	0.41
1:A:392:ASN:OD1	1:A:394:GLU:HB2	2.20	0.41
1:A:166:GLY:C	1:A:169:PRO:HD2	2.45	0.41
1:A:182:ILE:O	1:A:186:LEU:HD13	2.20	0.41
1:A:259:LEU:HA	1:A:265:GLY:HA2	2.02	0.41
1:A:135:HIS:HD2	3:A:546:HOH:O	2.03	0.41
1:A:163:THR:C	1:A:165:LYS:H	2.29	0.41
1:A:105:HIS:NE2	1:A:107:PRO:HG2	2.36	0.40
1:A:177:ARG:HG2	1:A:177:ARG:NH1	2.34	0.40

There are no symmetry-related clashes.

## 5.3 Torsion angles [i](#)

### 5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
1	A	429/457 (94%)	406 (95%)	19 (4%)	4 (1%)	<b>14</b> <b>14</b>

All (4) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	160	ILE
1	A	151	GLN
1	A	214	VAL
1	A	150	VAL

### 5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles
1	A	354/373 (95%)	344 (97%)	10 (3%)	38 52

All (10) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	78	LYS
1	A	106	LEU
1	A	152	ARG
1	A	153	GLN
1	A	155	GLN
1	A	178	THR
1	A	185	LEU
1	A	210	LEU
1	A	262	ILE
1	A	347	LEU

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (14) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	67	ASN
1	A	135	HIS
1	A	148	GLN
1	A	153	GLN
1	A	159	ASN
1	A	198	ASN
1	A	202	GLN
1	A	203	HIS

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type
1	A	204	GLN
1	A	217	GLN
1	A	244	HIS
1	A	324	ASN
1	A	351	HIS
1	A	382	ASN

### 5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

### 5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

### 5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

### 5.6 Ligand geometry [i](#)

1 ligand is modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z  > 2	Counts	RMSZ	# Z  > 2
2	GTP	A	458	-	33,34,34	1.24	3 (9%)	50,54,54	0.65	1 (2%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	GTP	A	458	-	-	9/22/38/38	0/3/3/3

All (3) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	A	458	GTP	PA-O3A	4.10	1.63	1.59
2	A	458	GTP	PB-O3B	3.70	1.63	1.59
2	A	458	GTP	PB-O3A	3.57	1.63	1.59

All (1) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	A	458	GTP	O3B-PB-O1B	-2.09	104.41	110.70

There are no chirality outliers.

All (9) torsion outliers are listed below:


Mol	Chain	Res	Type	Atoms
2	A	458	GTP	PB-O3B-PG-O2G
2	A	458	GTP	C5'-O5'-PA-O3A
2	A	458	GTP	C5'-O5'-PA-O1A
2	A	458	GTP	O4'-C4'-C5'-O5'
2	A	458	GTP	C3'-C4'-C5'-O5'
2	A	458	GTP	PA-O3A-PB-O2B
2	A	458	GTP	PB-O3A-PA-O2A
2	A	458	GTP	PB-O3B-PG-O1G
2	A	458	GTP	PB-O3B-PG-O3G

There are no ring outliers.

No monomer is involved in short contacts.

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient

equivalents in the CSD to analyse the geometry.



## 5.7 Other polymers [i](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

## 6 Fit of model and data [i](#)

### 6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	431/457 (94%)	0.05	13 (3%) 52 49	26, 43, 79, 101	0

All (13) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	162	THR	5.3
1	A	155	GLN	4.2
1	A	153	GLN	4.0
1	A	150	VAL	3.9
1	A	160	ILE	3.9
1	A	163	THR	3.8
1	A	146	GLY	3.6
1	A	66	GLY	2.9
1	A	154	ALA	2.8
1	A	28	ALA	2.5
1	A	161	GLY	2.5
1	A	159	ASN	2.2
1	A	147	LEU	2.1

### 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

### 6.3 Carbohydrates [i](#)


There are no oligosaccharides in this entry.

## 6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å <sup>2</sup> )	Q<0.9
2	GTP	A	458	32/32	0.90	0.09	35,43,52,57	5

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.



## 6.5 Other polymers [i](#)

There are no such residues in this entry.