



# Full wwPDB X-ray Structure Validation Report ⓘ

Mar 6, 2026 – 09:45 PM UTC

PDB ID : 5MAP / pdb\_00005map  
Title : X-ray generated oxyferrous complex of DtpA from Streptomyces lividans  
Authors : Moreno Chicano, T.; Chaplin, A.K.; Worrall, J.A.R.; Strange, R.W.; Hough, M.A.  
Deposited on : 2016-11-04  
Resolution : 1.49 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4-5-2 with Phenix2.0  
Mogul : 2022.3.0, CSD as543be (2022)  
Xtrriage (Phenix) : 2.0  
EDS : 3.0  
Buster-report : wwPDB partial adaption of 1.1.7 (2018)  
Percentile statistics : 20250101.v01 (using entries in the PDB archive January 1st 2025)  
CCP4 : 9.0.010 (Gargrove)  
Density-Fitness : 1.0.12  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : 2.49


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

*X-RAY DIFFRACTION*



The reported resolution of this entry is 1.49 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	180053	4037 (1.50-1.50)
Clashscore	190562	4235 (1.50-1.50)
Ramachandran outliers	187476	4153 (1.50-1.50)
Sidechain outliers	187428	4150 (1.50-1.50)
RSRZ outliers	180081	4039 (1.50-1.50)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	376	
1	B	376	

## 2 Entry composition i

There are 4 unique types of molecules in this entry. The entry contains 6332 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called DtpA.


Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	367	Total 2808	C 1740	N 529	O 527	S 12	11	2	0
1	B	369	Total 2846	C 1762	N 538	O 535	S 11	11	6	0

- Molecule 2 is PROTOPORPHYRIN IX CONTAINING FE (CCD ID: HEM) (formula:  $C_{34}H_{32}FeN_4O_4$ ).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
			Total	C	Fe	N	O		
2	A	1	Total 43	C 34	Fe 1	N 4	O 4	0	0
2	B	1	Total 43	C 34	Fe 1	N 4	O 4	0	0

- Molecule 3 is OXYGEN MOLECULE (CCD ID: OXY) (formula:  $O_2$ ).



Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
3	A	1	Total O 2 2	0	0
3	B	1	Total O 2 2	0	0

- Molecule 4 is water.

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
4	A	281	Total O 282 282	0	1
4	B	305	Total O 306 306	0	1



## 4 Data and refinement statistics

Property	Value	Source
Space group	P 1 21 1	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	59.79Å 70.63Å 77.66Å 90.00° 93.00° 90.00°	Depositor
Resolution (Å)	40.00 – 1.49 40.00 – 1.49	Depositor EDS
% Data completeness (in resolution range)	98.1 (40.00-1.49) 98.2 (40.00-1.49)	Depositor EDS
$R_{merge}$	0.11	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	1.62 (at 1.49Å)	Xtrriage
Refinement program	REFMAC 5.8.0155	Depositor
R, $R_{free}$	0.218 , 0.252 0.225 , 0.259	Depositor DCC
$R_{free}$ test set	4986 reflections (4.74%)	wwPDB-VP
Wilson B-factor (Å <sup>2</sup> )	10.8	Xtrriage
Anisotropy	0.191	Xtrriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.39 , 43.4	EDS
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.48$ , $\langle L^2 \rangle = 0.31$	Xtrriage
Estimated twinning fraction	No twinning to report.	Xtrriage
$F_o, F_c$ correlation	0.92	EDS
Total number of atoms	6332	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	13.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 6.17% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

## 5 Model quality [i](#)

### 5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: OXY, HEM

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# $ Z  > 5$	RMSZ	# $ Z  > 5$
1	A	1.20	5/2870 (0.2%)	1.12	3/3882 (0.1%)
1	B	1.19	2/2917 (0.1%)	1.13	5/3943 (0.1%)
All	All	1.20	7/5787 (0.1%)	1.12	8/7825 (0.1%)

All (7) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	205	VAL	N-CA	7.93	1.55	1.46
1	A	407	LYS	C-O	5.62	1.30	1.24
1	A	374	TYR	C-O	-5.59	1.17	1.23
1	A	350	ILE	C-O	-5.29	1.17	1.24
1	B	160	PHE	C-O	-5.21	1.18	1.24
1	A	98	GLY	C-O	-5.18	1.17	1.23
1	B	198	ILE	CA-C	-5.12	1.46	1.52

All (8) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	261	ALA	N-CA-C	8.62	120.68	111.28
1	B	269	VAL	CA-C-N	-6.54	113.90	120.31
1	B	269	VAL	C-N-CA	-6.54	113.90	120.31
1	A	242	THR	CA-C-N	6.16	129.87	120.82
1	A	242	THR	C-N-CA	6.16	129.87	120.82
1	B	377	GLY	N-CA-C	5.40	118.59	111.52
1	B	383	VAL	CA-C-N	-5.20	114.37	119.99
1	B	383	VAL	C-N-CA	-5.20	114.37	119.99

There are no chirality outliers.

There are no planarity outliers.

## 5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2808	0	2721	9	0
1	B	2846	0	2759	13	0
2	A	43	0	30	2	0
2	B	43	0	30	0	0
3	A	2	0	0	0	0
3	B	2	0	0	0	0
4	A	282	0	0	0	1
4	B	306	0	0	6	1
All	All	6332	0	5540	23	1

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 2.

All (23) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:331:GLU:O	4:B:1001:HOH:O	2.12	0.67
1:B:376:ASP:OD1	4:B:1002:HOH:O	2.13	0.66
1:B:256:PRO:O	4:B:1003:HOH:O	2.15	0.64
1:A:75:LEU:O	1:A:148:SER:HA	2.14	0.48
1:A:184:HIS:ND1	1:A:419:ARG:HD2	2.29	0.47
1:A:195:TRP:CZ2	1:A:197:GLN:HB2	2.49	0.47
2:A:901:HEM:HBB2	2:A:901:HEM:HMB2	1.97	0.47
1:A:262:ASP:OD1	1:A:262:ASP:N	2.42	0.46
1:B:335:MET:SD	1:B:337:LEU:HD21	2.56	0.45
1:B:81:MET:HG2	4:B:1264:HOH:O	2.17	0.44
1:B:134:ALA:C	4:B:1015:HOH:O	2.61	0.43
1:B:128:LEU:HD21	1:B:134:ALA:HB2	2.01	0.43
1:A:195:TRP:CH2	1:A:197:GLN:HB2	2.54	0.43
1:B:167[B]:GLU:HA	1:B:167[B]:GLU:OE1	2.18	0.42
1:B:135:GLY:N	4:B:1015:HOH:O	2.51	0.42
1:B:419[B]:ARG:HH21	1:B:419[B]:ARG:HD2	1.71	0.42
1:A:94[B]:MET:SD	1:A:243:ALA:HB2	2.60	0.41
1:B:94:MET:SD	1:B:243:ALA:HB2	2.61	0.41
1:B:295:ARG:NH2	1:B:419[B]:ARG:HD3	2.35	0.41

*Continued on next page...*

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:76:GLY:O	1:A:146:GLY:HA3	2.20	0.41
1:A:198:ILE:CD1	1:A:210:LEU:HA	2.51	0.41
2:A:901:HEM:HBB2	2:A:901:HEM:CMB	2.51	0.41
1:A:72:LEU:HA	1:B:72:LEU:HA	2.03	0.40

All (1) symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
4:A:1225:HOH:O	4:B:1114:HOH:O[1_655]	2.11	0.09

### 5.3 Torsion angles [i](#)

#### 5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	365/376 (97%)	355 (97%)	10 (3%)	0	100	100
1	B	371/376 (99%)	361 (97%)	10 (3%)	0	100	100
All	All	736/752 (98%)	716 (97%)	20 (3%)	0	100	100

There are no Ramachandran outliers to report.

#### 5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	283/285 (99%)	281 (99%)	2 (1%)	76	57
1	B	287/285 (101%)	285 (99%)	2 (1%)	76	57
All	All	570/570 (100%)	566 (99%)	4 (1%)	76	57

All (4) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	163	ARG
1	A	394	GLN
1	B	394	GLN
1	B	416	GLN

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (1) such sidechains are listed below:

Mol	Chain	Res	Type
1	B	158	HIS

### 5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

## 5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

## 5.6 Ligand geometry [i](#)

4 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the

expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z  > 2	Counts	RMSZ	# Z  > 2
3	OXY	B	902	2	1,1,1	0.27	0	-		
2	HEM	A	901	1,3	50,50,50	1.45	9 (18%)	67,82,82	1.52	11 (16%)
3	OXY	A	902	2	1,1,1	0.23	0	-		
2	HEM	B	901	1,3	50,50,50	1.38	7 (14%)	67,82,82	1.67	17 (25%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	HEM	A	901	1,3	-	4/14/54/54	-
2	HEM	B	901	1,3	-	4/14/54/54	-

All (16) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	B	901	HEM	FE-NB	4.21	2.07	1.94
2	A	901	HEM	FE-NB	4.14	2.07	1.94
2	A	901	HEM	C1B-NB	-3.81	1.33	1.40
2	B	901	HEM	CHB-C1B	3.71	1.45	1.38
2	A	901	HEM	FE-NC	2.81	2.04	1.95
2	B	901	HEM	O2A-CGA	-2.80	1.21	1.30
2	B	901	HEM	C4B-NB	-2.70	1.33	1.38
2	A	901	HEM	C4D-ND	-2.55	1.35	1.40
2	B	901	HEM	FE-NC	2.52	2.03	1.95
2	A	901	HEM	C3B-C4B	2.44	1.49	1.44
2	A	901	HEM	O1D-CGD	2.41	1.30	1.22
2	A	901	HEM	C3C-C2C	2.31	1.41	1.37
2	A	901	HEM	CHB-C1B	2.24	1.43	1.38
2	A	901	HEM	C1C-C2C	-2.17	1.41	1.45
2	B	901	HEM	O1D-CGD	2.09	1.28	1.22
2	B	901	HEM	CBD-CGD	2.01	1.55	1.50

All (28) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	A	901	HEM	C3B-C2B-C1B	4.06	109.46	106.41

*Continued on next page...*

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	A	901	HEM	CHC-C4B-NB	3.83	128.54	124.42
2	B	901	HEM	C1B-NB-C4B	3.68	109.56	105.21
2	A	901	HEM	CHA-C4D-ND	3.64	128.87	124.37
2	B	901	HEM	C3D-C4D-ND	3.51	114.02	110.17
2	B	901	HEM	CHD-C1D-C2D	-3.45	119.58	125.03
2	B	901	HEM	C4D-ND-C1D	-3.27	101.33	105.21
2	B	901	HEM	CHB-C1B-NB	3.27	128.41	124.37
2	B	901	HEM	C2D-C1D-ND	3.26	113.67	109.90
2	A	901	HEM	CHD-C4C-NC	3.25	128.00	124.45
2	A	901	HEM	C4B-C3B-C2B	-3.20	104.34	107.28
2	A	901	HEM	CHA-C4D-C3D	-3.08	119.55	125.23
2	B	901	HEM	CAD-C3D-C4D	3.00	129.93	124.70
2	A	901	HEM	C1B-NB-C4B	2.81	108.53	105.21
2	B	901	HEM	CBA-CAA-C2A	2.74	120.11	112.53
2	A	901	HEM	CHD-C1D-ND	2.66	127.28	124.42
2	A	901	HEM	CAD-C3D-C4D	2.62	129.26	124.70
2	B	901	HEM	CHA-C4D-C3D	-2.59	120.44	125.23
2	B	901	HEM	CHC-C4B-NB	2.59	127.21	124.42
2	B	901	HEM	C3B-C4B-NB	-2.57	107.62	109.47
2	B	901	HEM	C1A-CHA-C4D	-2.51	120.34	126.25
2	A	901	HEM	CHD-C1D-C2D	-2.44	121.18	125.03
2	B	901	HEM	CHA-C1A-NA	2.36	128.15	123.86
2	B	901	HEM	CMD-C2D-C1D	2.34	128.69	125.03
2	B	901	HEM	CHB-C4A-NA	2.23	127.91	123.86
2	B	901	HEM	CHD-C1D-ND	2.15	126.73	124.42
2	A	901	HEM	C1A-CHA-C4D	-2.12	121.27	126.25
2	B	901	HEM	C2A-C1A-NA	-2.10	107.82	110.15

There are no chirality outliers.

All (8) torsion outliers are listed below:


Mol	Chain	Res	Type	Atoms
2	A	901	HEM	CAA-CBA-CGA-O1A
2	A	901	HEM	CAA-CBA-CGA-O2A
2	B	901	HEM	CAA-CBA-CGA-O2A
2	B	901	HEM	CAA-CBA-CGA-O1A
2	A	901	HEM	CAD-CBD-CGD-O2D
2	B	901	HEM	CAD-CBD-CGD-O1D
2	B	901	HEM	CAD-CBD-CGD-O2D
2	A	901	HEM	CAD-CBD-CGD-O1D


There are no ring outliers.

1 monomer is involved in 2 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	A	901	HEM	2	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.





## 5.7 Other polymers [i](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

## 6 Fit of model and data

### 6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	367/376 (97%)	0.76	34 (9%) 14 15	5, 12, 23, 35	18 (4%)
1	B	369/376 (98%)	0.62	27 (7%) 21 23	4, 11, 24, 40	12 (3%)
All	All	736/752 (97%)	0.69	61 (8%) 17 19	4, 12, 24, 40	30 (4%)

All (61) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	B	337	LEU	5.8
1	B	136	SER	5.7
1	A	261	ALA	5.3
1	B	344	GLY	4.5
1	A	259	GLY	4.5
1	B	342	GLY	3.9
1	B	135	GLY	3.8
1	A	343	SER	3.8
1	B	340	THR	3.7
1	B	219	ALA	3.3
1	A	257	LYS	3.2
1	A	239	HIS	3.2
1	A	163	ARG	3.0
1	B	134	ALA	3.0
1	A	435	GLY	3.0
1	B	333	THR	2.9
1	A	441	ARG	2.8
1	B	218	GLY	2.8
1	B	69	ALA	2.8
1	A	342	GLY	2.8
1	B	271	GLU	2.8
1	B	327	SER	2.7
1	B	239	HIS	2.7
1	B	272	GLU	2.6

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type	RSRZ
1	B	332	SER	2.6
1	A	327	SER	2.5
1	B	343	SER	2.5
1	A	271	GLU	2.5
1	B	341	ASP	2.5
1	A	432	ALA	2.4
1	A	436	GLU	2.4
1	A	437	TYR	2.4
1	B	108	GLY	2.4
1	A	344	GLY	2.4
1	A	340	THR	2.3
1	A	172	VAL	2.3
1	A	433	ALA	2.3
1	B	326	GLY	2.3
1	A	258	PRO	2.2
1	A	110	GLY	2.2
1	A	337	LEU	2.2
1	B	238	ALA	2.2
1	A	184	HIS	2.2
1	A	265	ARG	2.2
1	A	282	TRP	2.2
1	B	236	ALA	2.2
1	A	262	ASP	2.2
1	A	166	LEU	2.1
1	A	173	ALA	2.1
1	A	175	ASP	2.1
1	B	115	ALA	2.1
1	B	346	LEU	2.1
1	A	399	ARG	2.1
1	A	280	PRO	2.1
1	B	380	ALA	2.0
1	A	444	GLU	2.0
1	B	338	GLU	2.0
1	A	218	GLY	2.0
1	B	280	PRO	2.0
1	A	161	PHE	2.0
1	A	398	LEU	2.0

## 6.2 Non-standard residues in protein, DNA, RNA chains

There are no non-standard protein/DNA/RNA residues in this entry.

### 6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

### 6.4 Ligands [i](#)




In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.


Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å <sup>2</sup> )	Q<0.9
3	OXY	B	902	2/2	0.77	0.15	13,13,13,20	0
3	OXY	A	902	2/2	0.85	0.13	16,16,16,17	0
2	HEM	B	901	43/43	0.97	0.07	6,6,9,10	0
2	HEM	A	901	43/43	0.98	0.07	7,7,9,10	0

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.

**Electron density around HEM B 901:**

$2mF_o-DF_c$  (at 0.7 rmsd) in gray  
 $mF_o-DF_c$  (at 3 rmsd) in purple (negative)  
and green (positive)





## 6.5 Other polymers [i](#)

There are no such residues in this entry.