



wwPDB X-ray Structure Validation Summary Report ⓘ

Aug 5, 2026 – 11:40 PM EDT

PDB ID : 7MH4 / pdb_00007mh4
Title : Crystal structure of R. sphaeroides Photosynthetic Reaction Center variant;
Y(M210)3-bromotyrosine
Authors : Mathews, I.; Weaver, J.; Boxer, S.G.
Deposited on : 2021-04-14
Resolution : 2.48 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4-5-2 with Phenix2.0
Mogul : 2022.3.0, CSD as543be (2022)
Xtriage (Phenix) : 2.0
EDS : 3.0
Buster-report : wwPDB partial adaption of 1.1.7 (2018)
Percentile statistics : 20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4 : 9.0.015 (Gargrove)
Density-Fitness : 1.0.12
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.50


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.48 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	7589 (2.50-2.46)
Clashscore	190562	8295 (2.50-2.46)
Ramachandran outliers	187476	8164 (2.50-2.46)
Sidechain outliers	187428	8166 (2.50-2.46)
RSRZ outliers	180081	7593 (2.50-2.46)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	H	266	2% 79% 10% • 11%
2	L	282	3% 93% 7%
3	M	308	2% 88% 9% •

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
6	BPH	L	305	X	-	-	-

2 Entry composition

There are 12 unique types of molecules in this entry. The entry contains 7335 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Reaction center protein H chain.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	H	238	Total	C	N	O	S	0	1	0
			1826	1169	312	336	9			

There are 7 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
H	260	VAL	-	expression tag	UNP P0C0Y7
H	261	HIS	-	expression tag	UNP P0C0Y7
H	262	HIS	-	expression tag	UNP P0C0Y7
H	263	HIS	-	expression tag	UNP P0C0Y7
H	264	HIS	-	expression tag	UNP P0C0Y7
H	265	HIS	-	expression tag	UNP P0C0Y7
H	266	HIS	-	expression tag	UNP P0C0Y7


- Molecule 2 is a protein called Reaction center protein L chain.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
2	L	281	Total	C	N	O	S	0	1	0
			2240	1513	356	363	8			

- Molecule 3 is a protein called Reaction center protein M chain.


Mol	Chain	Residues	Atoms						ZeroOcc	AltConf	Trace
3	M	300	Total	Br	C	N	O	S	0	0	0
			2396	2	1599	392	393	10			

- Molecule 4 is LAURYL DIMETHYLAMINE-N-OXIDE (CCD ID: LDA) (formula: C₁₄H₃₁NO).




Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
4	H	1	Total	C	N	O	0	0
			16	14	1	1		
4	M	1	Total	C	N	O	0	0
			16	14	1	1		
4	M	1	Total	C	N	O	0	0
			16	14	1	1		
4	M	1	Total	C	N	O	0	0
			16	14	1	1		
4	M	1	Total	C	N	O	0	0
			16	14	1	1		

- Molecule 5 is BACTERIOCHLOROPHYLL A (CCD ID: BCL) (formula: $C_{55}H_{74}MgN_4O_6$) (labeled as "Ligand of Interest" by depositor).




Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
5	L	1	Total 66	C 55	Mg 1	N 4	O 6	0	0
5	L	1	Total 66	C 55	Mg 1	N 4	O 6	0	0
5	M	1	Total 51	C 40	Mg 1	N 4	O 6	0	0
5	M	1	Total 66	C 55	Mg 1	N 4	O 6	0	0

- Molecule 6 is BACTERIOPHEOPHYTIN A (CCD ID: BPH) (formula: C₅₅H₇₆N₄O₆) (labeled as "Ligand of Interest" by depositor).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
6	L	1	Total	C	N	O	0	0
			65	55	4	6		
6	L	1	Total	C	N	O	0	0
			55	45	4	6		

- Molecule 7 is UBIQUINONE-10 (CCD ID: U10) (formula: C₅₉H₉₀O₄) (labeled as "Ligand of Interest" by depositor).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
7	L	1	Total	C	O	0	0
			18	14	4		
7	M	1	Total	C	O	0	0
			48	44	4		


- Molecule 8 is CHLORIDE ION (CCD ID: CL) (formula: Cl).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
8	L	1	Total	Cl	0	0
			1	1		

- Molecule 9 is FE (III) ION (CCD ID: FE) (formula: Fe).


Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
9	M	1	Total	Fe	0	0
			1	1		

- Molecule 10 is SPHEROIDENE (CCD ID: SPO) (formula: C₄₁H₆₀O).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
10	M	1	Total	C	O	0	0
			42	41	1		

- Molecule 11 is CARDIOLIPIN (CCD ID: CDL) (formula: $C_{81}H_{156}O_{17}P_2$).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
11	M	1	Total	C	O	P	0	0
			69	50	17	2		

- Molecule 12 is water.


Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
12	H	115	Total 115	O 115	0	0
12	L	60	Total 60	O 60	0	0
12	M	69	Total 70	O 70	0	1

3 Residue-property plots [i](#)


These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: Reaction center protein H chain

Chain H: 




- Molecule 2: Reaction center protein L chain

Chain L: 



- Molecule 3: Reaction center protein M chain

Chain M: 



4 Data and refinement statistics

Property	Value	Source
Space group	P 31 2 1	Depositor
Cell constants a, b, c, α , β , γ	141.34Å 141.34Å 186.87Å 90.00° 90.00° 120.00°	Depositor
Resolution (Å)	39.00 – 2.48 39.00 – 2.48	Depositor EDS
% Data completeness (in resolution range)	99.7 (39.00-2.48) 99.7 (39.00-2.48)	Depositor EDS
R_{merge}	0.11	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.82 (at 2.48Å)	Xtriage
Refinement program	REFMAC 5.8.0267	Depositor
R, R_{free}	0.168 , 0.193 0.175 , 0.197	Depositor DCC
R_{free} test set	3768 reflections (4.90%)	wwPDB-VP
Wilson B-factor (Å ²)	46.2	Xtriage
Anisotropy	0.140	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.29 , 46.5	EDS
L-test for twinning ²	$\langle L \rangle = 0.50$, $\langle L^2 \rangle = 0.33$	Xtriage
Estimated twinning fraction	0.010 for -h,-k,l	Xtriage
F_o, F_c correlation	0.95	EDS
Total number of atoms	7335	wwPDB-VP
Average B, all atoms (Å ²)	51.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 2.66% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: U10, LDA, CDL, BCL, SPO, CL, FE, DBY, BPH

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	H	1.00	1/1875 (0.1%)	1.32	2/2552 (0.1%)
2	L	0.99	0/2328	1.39	3/3186 (0.1%)
3	M	0.98	0/2472	1.42	4/3373 (0.1%)
All	All	0.99	1/6675 (0.0%)	1.38	9/9111 (0.1%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	H	0	1

All (1) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	H	49	PRO	C-O	-5.51	1.17	1.23

The worst 5 of 9 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	M	230	GLY	CA-C-N	6.54	127.35	120.03
3	M	230	GLY	C-N-CA	6.54	127.35	120.03
2	L	210	ASP	CA-CB-CG	5.59	118.19	112.60
1	H	11	ASP	CA-C-N	5.49	127.64	120.28
1	H	11	ASP	C-N-CA	5.49	127.64	120.28

There are no chirality outliers.

All (1) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	H	138	ALA	Peptide

5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	H	1826	0	1826	14	0
2	L	2240	0	2197	12	0
3	M	2396	0	2304	21	0
4	H	16	0	31	0	0
4	M	64	0	124	0	0
5	L	132	0	148	7	0
5	M	117	0	115	10	0
6	L	120	0	129	9	0
7	L	18	0	15	1	0
7	M	48	0	63	0	0
8	L	1	0	0	0	0
9	M	1	0	0	0	0
10	M	42	0	60	0	0
11	M	69	0	82	0	0
12	H	115	0	0	1	0
12	L	60	0	0	0	0
12	M	70	0	0	0	0
All	All	7335	0	7094	53	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 4.

The worst 5 of 53 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
3:M:16:ALA:HB1	3:M:32:VAL:HG11	1.56	0.87
5:L:301:BCL:ND	3:M:210:DBY:BR2	2.69	0.80
6:L:302:BPH:HBB3	6:L:302:BPH:HHC	1.64	0.77
5:L:301:BCL:HBB2	5:L:301:BCL:HHC	1.72	0.71
3:M:197:PHE:CZ	5:M:402:BCL:HBB2	2.27	0.69

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	H	237/266 (89%)	233 (98%)	4 (2%)	0	100	100
2	L	280/282 (99%)	271 (97%)	9 (3%)	0	100	100
3	M	297/308 (96%)	287 (97%)	9 (3%)	1 (0%)	36	53
All	All	814/856 (95%)	791 (97%)	22 (3%)	1 (0%)	48	66

All (1) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
3	M	301	HIS

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	H	194/215 (90%)	187 (96%)	7 (4%)	31	55
2	L	221/221 (100%)	214 (97%)	7 (3%)	34	58
3	M	234/240 (98%)	229 (98%)	5 (2%)	47	71
All	All	649/676 (96%)	630 (97%)	19 (3%)	37	62

5 of 19 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
3	M	39	LEU
3	M	216	PHE
3	M	278	LEU
3	M	72	ILE
2	L	150	ILE

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (3) such sidechains are listed below:

Mol	Chain	Res	Type
2	L	159	ASN
2	L	183	ASN
2	L	264	GLN

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

1 non-standard protein/DNA/RNA residue is modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
3	DBY	M	210	3	13,14,15	0.79	0	14,19,21	2.07	4 (28%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	DBY	M	210	3	-	2/5/6/8	0/1/1/1

There are no bond length outliers.

All (4) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	M	210	DBY	BR1-CE1-CZ	3.83	123.09	118.80
3	M	210	DBY	CB-CG-CD1	3.77	126.90	120.43
3	M	210	DBY	CB-CG-CD2	-3.37	114.64	120.43
3	M	210	DBY	BR2-CE2-CZ	3.15	122.33	118.80

There are no chirality outliers.

All (2) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
3	M	210	DBY	CA-CB-CG-CD1
3	M	210	DBY	CA-CB-CG-CD2

There are no ring outliers.

1 monomer is involved in 4 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
3	M	210	DBY	4	0

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

Of 17 ligands modelled in this entry, 2 are monoatomic - leaving 15 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
11	CDL	M	410	-	68,68,99	0.37	0	74,80,111	0.51	1 (1%)
4	LDA	M	404	-	13,15,15	0.13	0	14,17,17	0.28	0

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
5	BCL	M	401	3	54,59,74	1.95	15 (27%)	61,97,115	2.32	21 (34%)
7	U10	M	408	-	48,48,63	0.64	1 (2%)	60,61,79	0.59	1 (1%)
5	BCL	M	402	3	69,74,74	1.68	12 (17%)	79,115,115	2.06	23 (29%)
10	SPO	M	409	-	41,41,41	1.24	7 (17%)	47,50,50	1.11	3 (6%)
5	BCL	L	304	2	69,74,74	1.67	14 (20%)	79,115,115	1.95	20 (25%)
4	LDA	M	403	-	13,15,15	0.26	0	14,17,17	0.31	0
7	U10	L	303	-	18,18,63	1.00	1 (5%)	24,25,79	1.04	2 (8%)
4	LDA	H	301	-	13,15,15	0.17	0	14,17,17	0.26	0
5	BCL	L	301	2	69,74,74	1.73	12 (17%)	79,115,115	2.13	23 (29%)
6	BPH	L	305	-	49,60,70	0.91	2 (4%)	47,89,101	1.14	4 (8%)
4	LDA	M	405	-	13,15,15	0.15	0	14,17,17	0.20	0
6	BPH	L	302	-	59,70,70	0.75	2 (3%)	59,101,101	1.06	4 (6%)
4	LDA	M	406	-	13,15,15	0.19	0	14,17,17	0.18	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
11	CDL	M	410	-	-	23/79/79/110	-
4	LDA	M	404	-	-	4/13/13/13	-
5	BCL	M	401	3	-	1/23/119/137	-
7	U10	M	408	-	-	4/45/69/87	0/1/1/1
5	BCL	M	402	3	-	2/41/137/137	-
10	SPO	M	409	-	-	6/47/47/47	-
5	BCL	L	304	2	-	2/41/137/137	-
4	LDA	M	403	-	-	3/13/13/13	-
7	U10	L	303	-	-	5/9/33/87	0/1/1/1
4	LDA	H	301	-	-	6/13/13/13	-
5	BCL	L	301	2	-	8/41/137/137	-
6	BPH	L	305	-	1/1/16/22	7/25/93/105	0/5/6/6
4	LDA	M	405	-	-	4/13/13/13	-
6	BPH	L	302	-	-	7/37/105/105	0/5/6/6
4	LDA	M	406	-	-	3/13/13/13	-

The worst 5 of 66 bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
5	L	301	BCL	O2D-CGD	5.34	1.46	1.33
5	L	304	BCL	O2D-CGD	4.92	1.45	1.33
5	M	402	BCL	O2D-CGD	4.88	1.45	1.33
5	M	401	BCL	C3D-C4D	-4.57	1.33	1.44
5	L	304	BCL	C3D-C4D	-4.57	1.33	1.44

The worst 5 of 102 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
5	M	401	BCL	CMD-C2D-C1D	6.63	136.40	124.73
5	M	402	BCL	CMD-C2D-C1D	6.55	136.26	124.73
5	L	304	BCL	CMD-C2D-C1D	6.50	136.18	124.73
5	L	301	BCL	CMD-C2D-C1D	6.43	136.05	124.73
5	M	401	BCL	C2B-C1B-NB	5.60	116.13	110.33

All (1) chirality outliers are listed below:

Mol	Chain	Res	Type	Atom
6	L	305	BPH	C8

5 of 85 torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
4	H	301	LDA	C2-C1-N1-CM1
4	H	301	LDA	C2-C1-N1-CM2
4	M	404	LDA	N1-C1-C2-C3
7	L	303	U10	C1-C6-C7-C8
7	L	303	U10	C5-C6-C7-C8


There are no ring outliers.


7 monomers are involved in 27 short contacts:












Mol	Chain	Res	Type	Clashes	Symm-Clashes
5	M	401	BCL	4	0
5	M	402	BCL	7	0
5	L	304	BCL	1	0
7	L	303	U10	1	0
5	L	301	BCL	6	0
6	L	305	BPH	6	0
6	L	302	BPH	3	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths,


bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.






Ligand BCL M 402	
 Bond lengths	 Bond angles
 Torsions	 Rings
Ligand SPO M 409	
 Bond lengths	 Bond angles
 Torsions	 Rings
Ligand BCL L 304	
 Bond lengths	 Bond angles
 Torsions	 Rings


Ligand U10 L 303




Ligand BCL L 301



Ligand BPH L 305



Ligand BPH L 302



5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	H	238/266 (89%)	-0.09	4 (1%) 69 66	29, 47, 67, 97	1 (0%)
2	L	281/282 (99%)	0.14	9 (3%) 50 47	22, 46, 79, 96	1 (0%)
3	M	299/308 (97%)	0.03	6 (2%) 65 62	31, 49, 80, 100	0
All	All	818/856 (95%)	0.03	19 (2%) 61 58	22, 47, 78, 100	2 (0%)

The worst 5 of 19 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
3	M	301	HIS	4.1
2	L	281	GLY	4.1
3	M	3	TYR	3.4
1	H	18[A]	TYR	3.0
2	L	155	ASP	2.9

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled ‘Q < 0.9’ lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
3	DBY	M	210	14/15	0.98	0.06	32,35,37,41	2

6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.4 Ligands ⓘ




In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q<0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
4	LDA	M	406	16/16	0.81	0.36	85,90,120,128	0
4	LDA	M	405	16/16	0.86	0.32	91,102,121,126	0
7	U10	L	303	18/63	0.88	0.27	35,40,43,43	18
11	CDL	M	410	69/100	0.89	0.19	58,77,98,112	0
4	LDA	M	404	16/16	0.90	0.24	73,82,105,107	0
7	U10	M	408	48/63	0.91	0.19	33,48,101,103	0
4	LDA	M	403	16/16	0.92	0.20	65,74,84,85	0
10	SPO	M	409	42/42	0.93	0.15	49,62,92,98	0
8	CL	L	306	1/1	0.93	0.11	82,82,82,82	0
6	BPH	L	305	55/65	0.94	0.12	39,49,90,94	0
4	LDA	H	301	16/16	0.94	0.15	58,68,85,87	0
5	BCL	L	301	66/66	0.95	0.10	34,38,70,74	0
5	BCL	M	401	51/66	0.95	0.10	35,38,63,67	0
6	BPH	L	302	65/65	0.95	0.09	30,37,51,54	0
5	BCL	M	402	66/66	0.96	0.09	35,38,72,87	0
5	BCL	L	304	66/66	0.96	0.09	34,38,55,66	0
9	FE	M	407	1/1	0.99	0.03	33,33,33,33	0




The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.

Electron density around U10 L 303:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)



**Electron density around CDL M 410:**

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)






Electron density around U10 M 408:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)




**Electron density around SPO M 409:**

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)






Electron density around BPH L 305:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)






Electron density around BCL L 301:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)





Electron density around BCL M 401:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)





Electron density around BPH L 302:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)






Electron density around BCL M 402:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)

**Electron density around BCL L 304:**

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)



6.5 Other polymers [i](#)

There are no such residues in this entry.