wwPDB X-ray Structure Validation Summary Report (i)

Sep 9, 2026 – 09:09 am BST

PDB ID	:	5N78 / pdb_00005n78
Title	:	Crystal structure of the cytosolic domain of the CorA Mg2+ channel from Escherichia coli in complex with magnesium and cobalt hexammine
Authors	:	Lerche, M.; Sandhu, H.; Flockner, L.; Hogbom, M.; Rapp, M.
Deposited on	:	2017-02-20
Resolution	:	2.85 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at *validation@mail.wwpdb.org*
A user guide is available at
https://www.wwpdb.org/validation/2017/XrayValidationReportHelp
with specific help available everywhere you see the (i) symbol.

The types of validation reports are described at
http://www.wwpdb.org/validation/2017/FAQs#types.

The following versions of software and data (see references (i)) were used in the production of this report:


MolProbity	:	4-5-2 with Phenix2.0
Mogul	:	1.8.4, CSD as541be (2020)
Xtriage (Phenix)	:	2.0
EDS	:	3.0
Percentile statistics	:	20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4	:	9.0.015 (Gargrove)
Density-Fitness	:	1.0.12
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.50

1 Overall quality at a glance (i)

The following experimental techniques were used to determine the structure:
X-RAY DIFFRACTION

The reported resolution of this entry is 2.85 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	1407 (2.88-2.84)
Clashscore	190562	1446 (2.88-2.84)
Ramachandran outliers	187476	1406 (2.88-2.84)
Sidechain outliers	187428	1407 (2.88-2.84)
RSRZ outliers	180081	1408 (2.88-2.84)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for >=3, 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions <=5% The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	257	93% 7%
1	B	257	88% 10%
1	C	257	86% 12%
1	D	257	93% 6%
1	E	257	91% 9%

2 Entry composition (i)

There are 5 unique types of molecules in this entry. The entry contains 10611 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.


- Molecule 1 is a protein called Magnesium transport protein CorA.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	257	Total	C	N	O	S	0	0	0
			2081	1308	360	405	8			
1	B	257	Total	C	N	O	S	0	0	0
			2088	1310	362	408	8			
1	C	254	Total	C	N	O	S	0	0	0
			2059	1296	355	400	8			
1	D	257	Total	C	N	O	S	0	0	0
			2076	1304	359	405	8			
1	E	257	Total	C	N	O	S	0	0	0
			2082	1307	359	408	8			

- Molecule 2 is MAGNESIUM ION (CCD ID: MG) (formula: Mg).


Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
2	A	2	Total	Mg	0	0
			2	2		
2	B	1	Total	Mg	0	0
			1	1		
2	C	1	Total	Mg	0	0
			1	1		
2	D	1	Total	Mg	0	0
			1	1		
2	E	1	Total	Mg	0	0
			1	1		

- Molecule 3 is COBALT HEXAMMINE(III) (CCD ID: NCO) (formula: $CoH_{18}N_6$).



Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
3	A	1	Total Co N 7 1 6	0	0
3	A	1	Total Co N 7 1 6	0	0
3	A	1	Total Co N 7 1 6	0	0
3	B	1	Total Co N 7 1 6	0	0
3	B	1	Total Co N 7 1 6	0	0
3	B	1	Total Co N 7 1 6	0	0
3	C	1	Total Co N 7 1 6	0	0
3	C	1	Total Co N 7 1 6	0	0
3	D	1	Total Co N 7 1 6	0	0
3	D	1	Total Co N 7 1 6	0	0
3	D	1	Total Co N 7 1 6	0	0
3	E	1	Total Co N 7 1 6	0	0
3	E	1	Total Co N 7 1 6	0	0
3	E	1	Total Co N 7 1 6	0	0

- Molecule 4 is 2-(2-METHOXYETHOXY)ETHANOL (CCD ID: PG0) (formula: $C_5H_{12}O_3$).



Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
4	B	1	Total C O 8 5 3	0	0
4	C	1	Total C O 8 5 3	0	0
4	E	1	Total C O 8 5 3	0	0

- Molecule 5 is water.


Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
5	A	15	Total O 15 15	0	0
5	B	23	Total O 23 23	0	0
5	C	23	Total O 23 23	0	0
5	D	18	Total O 18 18	0	0
5	E	18	Total O 18 18	0	0

3 Residue-property plots (i)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density (RSRZ > 2). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


• Molecule 1: Magnesium transport protein CorA

Chain A:




• Molecule 1: Magnesium transport protein CorA

Chain B:




• Molecule 1: Magnesium transport protein CorA

Chain C:




• Molecule 1: Magnesium transport protein CorA

Chain D:



• Molecule 1: Magnesium transport protein CorA

Chain E:



4 Data and refinement statistics (i)

Property	Value	Source
Space group	P 1 21 1	Depositor
Cell constants a, b, c, α, β, γ	73.72Å 117.32Å 91.86Å 90.00° 103.14° 90.00°	Depositor
Resolution (Å)	29.44 – 2.85 29.44 – 2.85	Depositor EDS
% Data completeness (in resolution range)	98.1 (29.44-2.85) 98.2 (29.44-2.85)	Depositor EDS
R_{merge}	0.12	Depositor
R_{sym}	(Not available)	Depositor
$< I/\sigma(I) >$ [1]	1.94 (at 2.85Å)	Xtriage
Refinement program	REFMAC 5.8.0073	Depositor
R, R_{free}	0.178 , 0.218 0.182 , 0.215	Depositor DCC
R_{free} test set	1769 reflections (4.97%)	wwPDB-VP
Wilson B-factor (Å^2)	65.3	Xtriage
Anisotropy	0.342	Xtriage
Bulk solvent k_{sol}(e/Å^3), B_{sol}(Å^2)	0.30 , 43.5	EDS
L-test for twinning[2]	$< \|L\| > = 0.50$, $< L^2 > = 0.33$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o,F_c correlation	0.96	EDS
Total number of atoms	10611	wwPDB-VP
Average B, all atoms (Å^2)	78.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 4.67% of the height of the origin peak. No significant pseudotranslation is detected.*

[1] Intensities estimated from amplitudes.

[2] Theoretical values of $< |L| >$, $< L^2 >$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: PG0, MG, NCO

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	#$\|Z\|>5$	RMSZ	#$\|Z\|>5$
1	A	0.84	0/2114	0.88	1/2853 (0.0%)
1	B	0.90	0/2120	0.95	2/2860 (0.1%)
1	C	0.90	0/2090	0.93	4/2819 (0.1%)
1	D	0.79	0/2108	0.87	1/2845 (0.0%)
1	E	0.86	0/2114	0.90	0/2853
All	All	0.86	0/10546	0.91	8/14230 (0.1%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand.A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	A	0	1
1	B	0	3
1	C	0	2
1	D	0	1
1	E	0	1
All	All	0	8

There are no bond length outliers.

The worst 5 of 8 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(^{o})	Ideal(^{o})
1	B	167	GLN	N-CA-C	12.00	129.64	111.04
1	C	120	ARG	N-CA-C	-9.57	95.61	109.96
1	C	204	ARG	CB-CG-CD	6.56	126.38	111.30
1	C	121	SER	N-CA-C	6.28	120.34	111.92
1	D	48	GLN	N-CA-C	5.71	117.96	108.20

There are no chirality outliers.

5 of 8 planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	A	47	GLY	Peptide
1	B	164	GLU	Peptide
1	B	166	HIS	Peptide
1	B	47	GLY	Peptide
1	C	47	GLY	Peptide

5.2 Too-close contacts (i)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2081	0	2019	8	0
1	B	2088	0	2031	17	0
1	C	2059	0	2004	16	0
1	D	2076	0	2017	5	0
1	E	2082	0	2020	10	1
2	A	2	0	0	0	0
2	B	1	0	0	0	0
2	C	1	0	0	0	0
2	D	1	0	0	0	0
2	E	1	0	0	0	0
3	A	21	0	0	0	1
3	B	21	0	0	2	0
3	C	14	0	0	1	0
3	D	21	0	0	0	0
3	E	21	0	0	1	0
4	B	8	0	12	1	0
4	C	8	0	12	0	0
4	E	8	0	12	2	0
5	A	15	0	0	0	0
5	B	23	0	0	4	0
5	C	23	0	0	6	0
5	D	18	0	0	1	0
5	E	18	0	0	2	0
All	All	10611	0	10127	52	1

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including

hydrogen atoms). The all-atom clashscore for this structure is 3.

The worst 5 of 52 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:C:85:ASP:OD1	3:C:304:NCO:N1	2.00	0.93
1:B:149:GLU:OE1	3:B:305:NCO:N4	2.04	0.91
1:C:209:PRO:HD3	5:C:406:HOH:O	1.75	0.85
1:B:30:ASP:OD2	5:B:401:HOH:O	2.03	0.76
1:C:118:ARG:O	1:C:120:ARG:O	2.04	0.76

All (1) symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:E:19:GLU:OE1	3:A:304:NCO:N4[1_556]	1.96	0.24

5.3 Torsion angles (i)

5.3.1 Protein backbone (i)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	255/257 (99%)	245 (96%)	10 (4%)	0	100	100
1	B	255/257 (99%)	245 (96%)	10 (4%)	0	100	100
1	C	250/257 (97%)	242 (97%)	8 (3%)	0	100	100
1	D	255/257 (99%)	247 (97%)	8 (3%)	0	100	100
1	E	255/257 (99%)	247 (97%)	8 (3%)	0	100	100
All	All	1270/1285 (99%)	1226 (96%)	44 (4%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains (i)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	222/227 (98%)	213 (96%)	9 (4%)	27	52
1	B	224/227 (99%)	215 (96%)	9 (4%)	28	53
1	C	220/227 (97%)	211 (96%)	9 (4%)	27	52
1	D	222/227 (98%)	213 (96%)	9 (4%)	27	52
1	E	223/227 (98%)	212 (95%)	11 (5%)	22	45
All	All	1111/1135 (98%)	1064 (96%)	47 (4%)	26	51

5 of 47 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	D	16	GLU
1	D	218	GLU
1	D	23	LEU
1	D	153	SER
1	E	16	GLU

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 13 such sidechains are listed below:

Mol	Chain	Res	Type
1	C	86	HIS
1	C	242	GLN
1	E	242	GLN
1	D	242	GLN
1	E	86	HIS

5.3.3 RNA (i)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains (i)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates (i)

There are no oligosaccharides in this entry.

5.6 Ligand geometry (i)

Of 23 ligands modelled in this entry, 6 are monoatomic - leaving 17 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\#\|Z\| > 2$	Counts	RMSZ	$\#\|Z\| > 2$
4	PG0	C	302	-	7,7,7	0.63	0	6,6,6	0.41	0
3	NCO	E	305	-	6,6,6	0.67	0	-		
3	NCO	D	302	-	6,6,6	0.76	0	-		
3	NCO	B	305	-	6,6,6	0.66	0	-		
3	NCO	D	304	-	6,6,6	0.72	0	-		
3	NCO	C	303	-	6,6,6	0.70	0	-		
4	PG0	B	302	-	7,7,7	0.80	0	6,6,6	0.78	0
3	NCO	A	303	-	6,6,6	0.79	0	-		
3	NCO	B	304	-	6,6,6	0.70	0	-		
3	NCO	C	304	-	6,6,6	0.64	0	-		
3	NCO	D	303	-	6,6,6	0.70	0	-		
3	NCO	E	303	-	6,6,6	0.66	0	-		
4	PG0	E	302	-	7,7,7	0.64	0	6,6,6	0.47	0
3	NCO	E	304	-	6,6,6	0.67	0	-		
3	NCO	A	304	-	6,6,6	0.64	0	-		
3	NCO	B	303	-	6,6,6	0.73	0	-		
3	NCO	A	305	-	6,6,6	0.65	0	-		

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
4	PG0	C	302	-	-	0/5/5/5	-
4	PG0	B	302	-	-	2/5/5/5	-
4	PG0	E	302	-	-	3/5/5/5	-

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

All (5) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
4	B	302	PG0	OTT-C1-C2-O1
4	E	302	PG0	OTT-C1-C2-O1
4	B	302	PG0	C3-C4-O2-C5
4	E	302	PG0	C4-C3-O1-C2
4	E	302	PG0	O1-C3-C4-O2

There are no ring outliers.

6 monomers are involved in 8 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
3	B	305	NCO	2	0
4	B	302	PG0	1	0
3	C	304	NCO	1	0
4	E	302	PG0	2	0
3	E	304	NCO	1	0
3	A	304	NCO	0	1

5.7 Other polymers (i)

There are no such residues in this entry.

5.8 Polymer linkage issues (i)

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

In the following table, the column labelled '#RSRZ> 2' contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95^{th} percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled 'Q< 0.9' lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2			OWAB(Å^2)	Q<0.9
1	A	257/257 (100%)	-0.56	0	100	100	42, 74, 122, 187	0
1	B	257/257 (100%)	-0.59	0	100	100	37, 65, 117, 161	0
1	C	254/257 (98%)	-0.59	2 (0%)	82	78	40, 70, 111, 171	0
1	D	257/257 (100%)	-0.48	1 (0%)	88	87	49, 81, 140, 190	0
1	E	257/257 (100%)	-0.51	2 (0%)	82	78	44, 71, 120, 183	0
All	All	1282/1285 (99%)	-0.55	5 (0%)	88	87	37, 72, 126, 190	0

All (5) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	E	167	GLN	4.6
1	C	167	GLN	2.4
1	D	165	GLY	2.3
1	C	207	ARG	2.1
1	E	166	HIS	2.0

6.2 Non-standard residues in protein, DNA, RNA chains

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates

There are no oligosaccharides in this entry.

6.4 Ligands

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum,

median, 95^{th} percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å^2)	Q<0.9
3	NCO	B	304	7/7	0.65	0.09	145,153,161,164	0
3	NCO	E	305	7/7	0.68	0.07	169,173,177,180	0
3	NCO	C	304	7/7	0.70	0.09	163,167,172,175	0
3	NCO	E	303	7/7	0.71	0.13	172,181,191,196	0
3	NCO	C	303	7/7	0.74	0.09	149,154,165,165	0
3	NCO	B	305	7/7	0.75	0.11	149,158,164,166	0
3	NCO	D	304	7/7	0.76	0.09	142,152,155,166	0
3	NCO	D	303	7/7	0.78	0.10	152,156,159,163	0
3	NCO	D	302	7/7	0.81	0.09	120,131,140,144	0
3	NCO	E	304	7/7	0.82	0.08	138,142,151,158	0
4	PG0	E	302	8/8	0.82	0.16	90,104,108,109	0
3	NCO	A	305	7/7	0.83	0.09	134,144,148,155	0
3	NCO	A	304	7/7	0.84	0.10	127,147,149,155	0
4	PG0	B	302	8/8	0.85	0.19	82,104,113,116	0
4	PG0	C	302	8/8	0.87	0.14	88,98,107,108	0
3	NCO	A	303	7/7	0.90	0.08	106,112,120,122	0
2	MG	D	301	1/1	0.92	0.12	78,78,78,78	0
2	MG	B	301	1/1	0.95	0.10	58,58,58,58	0
2	MG	A	301	1/1	0.95	0.13	50,50,50,50	0
2	MG	A	302	1/1	0.97	0.06	69,69,69,69	0
3	NCO	B	303	7/7	0.97	0.08	81,88,90,91	0
2	MG	E	301	1/1	0.97	0.08	55,55,55,55	0
2	MG	C	301	1/1	0.98	0.04	54,54,54,54	0

6.5 Other polymers (i)

There are no such residues in this entry.