wwPDB X-ray Structure Validation Summary Report (i)

Aug 4, 2026 – 05:37 PM EDT

PDB ID	:	3NAK / pdb_00003nak
Title	:	Crystal structure of the complex of goat lactoperoxidase with hypothiocyanite at 3.3 A resolution
Authors	:	Vikram, G.; Singh, R.P.; Singh, A.K.; Sinha, M.; Bhushan, A.; Kaur, P.; Sharma, S.; Singh, T.P.
Deposited on	:	2010-06-02
Resolution	:	3.30 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at *validation@mail.wwpdb.org*
A user guide is available at
https://www.wwpdb.org/validation/2017/XrayValidationReportHelp
with specific help available everywhere you see the (i) symbol.

The types of validation reports are described at
http://www.wwpdb.org/validation/2017/FAQs#types.

The following versions of software and data (see references (1)) were used in the production of this report:

MolProbity	:	4-5-2 with Phenix2.0
Mogul	:	2022.3.0, CSD as543be (2022)
Xtriage (Phenix)	:	2.0
EDS	:	3.0
Buster-report	:	wwPDB partial adaption of 1.1.7 (2018)
Percentile statistics	:	20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4	:	9.0.015 (Gargrove)
Density-Fitness	:	1.0.12
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.50

1 Overall quality at a glance (i)

The following experimental techniques were used to determine the structure:
X-RAY DIFFRACTION

The reported resolution of this entry is 3.30 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	1169 (3.32-3.28)
Clashscore	190562	1209 (3.32-3.28)
Ramachandran outliers	187476	1188 (3.32-3.28)
Sidechain outliers	187428	1187 (3.32-3.28)
RSRZ outliers	180081	1169 (3.32-3.28)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for >=3, 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions <=5% The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	595	5% 78% 20% •
1	B	595	4% 79% 17% • •
2	C	3	33% 67%
2	D	3	33% 67%

Continued on next page...

Continued from previous page...

Mol	Chain	Length	Quality of chain
2	G	3	67% 33%
3	E	3	100%
3	H	3	33% 67%
3	I	3	100%
4	F	2	100%
4	J	2	100%

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
7	OSM	A	3333	-	-	X	-
7	OSM	B	3333	-	-	X	-
8	FMT	A	2001	-	-	X	-
8	FMT	A	2002	-	-	X	-

2 Entry composition (i)

There are 9 unique types of molecules in this entry. The entry contains 10098 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Lactoperoxidase.

Mol	Chain	Residues	Atoms						ZeroOcc	AltConf	Trace
			Total	C	N	O	P	S			
1	A	595	4757	3021	844	865	1	26	0	0	0
1	B	595	4757	3021	844	865	1	26	0	0	0

- Molecule 2 is an oligosaccharide called alpha-D-mannopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose.



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	Trace
			Total	C	N	O			
2	C	3	39	22	2	15	0	0	0
2	D	3	39	22	2	15	0	0	0
2	G	3	39	22	2	15	0	0	0

- Molecule 3 is an oligosaccharide called beta-D-mannopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose.



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	Trace
			Total	C	N	O			
3	E	3	39	22	2	15	0	0	0
3	H	3	39	22	2	15	0	0	0

Continued on next page...

Continued from previous page...

Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	Trace
			Total	C	N	O			
3	I	3	39	22	2	15	0	0	0

- Molecule 4 is an oligosaccharide called 2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose.

Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	Trace
			Total	C	N	O			
4	F	2	28	16	2	10	0	0	0
			Total	C	N	O			
4	J	2	28	16	2	10	0	0	0

- Molecule 5 is CALCIUM ION (CCD ID: CA) (formula: Ca).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
			Total	Ca		
5	A	1	1	1	0	0
			Total	Ca		
5	B	1	1	1	0	0

- Molecule 6 is PROTOPORPHYRIN IX CONTAINING FE (CCD ID: HEM) (formula: $C_{34}H_{32}FeN_4O_4$).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
			Total	C	Fe	N	O		
6	A	1	43	34	1	4	4	0	0
6	B	1	43	34	1	4	4	0	0

- Molecule 7 is 1-(OXIDOSULFANYL)METHANAMINE (CCD ID: OSM) (formula: CH_5NOS).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
			Total	C	N	O	S		
7	A	1	4	1	1	1	1	0	0
7	B	1	4	1	1	1	1	0	0

- Molecule 8 is FORMIC ACID (CCD ID: FMT) (formula: CH_2O_2).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
			Total	C	O		
8	A	1	3	1	2	0	0
			Total	C	O		
8	A	1	3	1	2	0	0
			Total	C	O		
8	B	1	3	1	2	0	0
			Total	C	O		
8	B	1	3	1	2	0	0

- Molecule 9 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
			Total	O		
9	A	98	98	98	0	0
			Total	O		
9	B	88	88	88	0	0

3 Residue-property plots (i)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density (RSRZ > 2). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

• Molecule 1: Lactoperoxidase

Chain A:



• Molecule 1: Lactoperoxidase

Chain B:



• Molecule 2: alpha-D-mannopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose

Chain C: 33% 67%

NAG1 NAG2 MAN3

• Molecule 2: alpha-D-mannopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose

Chain D: 33% 67%

NAG1 NAG2 MAN3

• Molecule 2: alpha-D-mannopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose

Chain G: 67% 33%

NAG1 NAG2 MAN3

• Molecule 3: beta-D-mannopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose

Chain E: 100%

NAG1 NAG2 BMA3

• Molecule 3: beta-D-mannopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose

Chain H: 33% 67%

NAG1 NAG2 BMA3

• Molecule 3: beta-D-mannopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose

Chain I: 100%

NAG1 NAG2 BMA3

• Molecule 4: 2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose

Chain F: 100%

NAG1 NAG2

• Molecule 4: 2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose

Chain J: 100%

NAG1
NAG2

4 Data and refinement statistics (i)

Property	Value	Source
Space group	P 1	Depositor
Cell constants a, b, c, α, β, γ	57.99Å 72.27Å 83.65Å 85.45° 84.04° 75.89°	Depositor
Resolution (Å)	19.87 – 3.30 19.87 – 3.30	Depositor EDS
% Data completeness (in resolution range)	94.5 (19.87-3.30) 94.5 (19.87-3.30)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	(Not available)	Depositor
$< I/\sigma(I) >$ [1]	1.70 (at 3.29Å)	Xtriage
Refinement program	REFMAC 5.2.0019	Depositor
R, R_{free}	0.205 , 0.243 0.208 , 0.281	Depositor DCC
R_{free} test set	950 reflections (5.11%)	wwPDB-VP
Wilson B-factor (Å^2)	43.6	Xtriage
Anisotropy	0.448	Xtriage
Bulk solvent k_{sol}(e/Å^3), B_{sol}(Å^2)	0.44 , 82.6	EDS
L-test for twinning[2]	$< \|L\| > = 0.48$, $< L^2 > = 0.31$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o,F_c correlation	0.89	EDS
Total number of atoms	10098	wwPDB-VP
Average B, all atoms (Å^2)	25.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 6.40% of the height of the origin peak. No significant pseudotranslation is detected.*

[1]Intensities estimated from amplitudes.

[2]Theoretical values of $< |L| >$, $< L^2 >$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality (i)

5.1 Standard geometry (i)

Bond lengths and bond angles in the following residue types are not validated in this section: NAG, BMA, MAN, HEM, CA, SEP, OSM, FMT

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\#\|Z\|>5$	RMSZ	$\#\|Z\|>5$
1	A	0.51	2/4875 (0.0%)	0.87	8/6621 (0.1%)
1	B	0.53	1/4875 (0.0%)	0.91	16/6621 (0.2%)
All	All	0.52	3/9750 (0.0%)	0.89	24/13242 (0.2%)

All (3) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	B	332	ASN	CG-OD1	5.18	1.33	1.23
1	A	258	GLU	C-O	-5.08	1.18	1.24
1	A	258	GLU	CA-C	-5.05	1.46	1.52

The worst 5 of 24 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(^{o})	Ideal(^{o})
1	B	173	GLN	OE1-CD-NE2	-9.25	113.35	122.60
1	B	174	SER	N-CA-CB	-7.40	106.70	114.10
1	B	11	PRO	N-CA-C	7.08	123.63	111.03
1	B	173	GLN	CG-CD-NE2	6.80	126.61	116.40
1	B	118	GLU	N-CA-C	6.57	118.86	108.67

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts (i)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	4757	0	4644	115	0
1	B	4757	0	4644	115	0
2	C	39	0	34	0	0
2	D	39	0	34	0	0
2	G	39	0	34	1	0
3	E	39	0	34	0	0
3	H	39	0	34	1	0
3	I	39	0	34	0	0
4	F	28	0	25	0	0
4	J	28	0	25	0	0
5	A	1	0	0	0	0
5	B	1	0	0	0	0
6	A	43	0	30	11	0
6	B	43	0	30	17	0
7	A	4	0	5	3	0
7	B	4	0	5	4	0
8	A	6	0	2	6	0
8	B	6	0	2	0	0
9	A	98	0	0	1	0
9	B	88	0	0	2	0
All	All	10098	0	9616	237	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 12.

The worst 5 of 237 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:258:GLU:OE2	6:B:1003:HEM:HMB3	1.16	1.25
1:B:258:GLU:OE2	6:B:1003:HEM:CMB	1.85	1.23
1:B:167:CYS:HB3	1:B:168:PRO:HD3	1.21	1.19
1:B:167:CYS:CB	1:B:168:PRO:HD3	1.78	1.13
1:A:96:ARG:HG3	1:A:96:ARG:HH11	1.10	1.12

There are no symmetry-related clashes.

5.3 Torsion angles (i)

5.3.1 Protein backbone (i)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
1	A	592/595 (100%)	551 (93%)	36 (6%)	5 (1%)	16 45
1	B	592/595 (100%)	555 (94%)	28 (5%)	9 (2%)	8 32
All	All	1184/1190 (100%)	1106 (93%)	64 (5%)	14 (1%)	10 37

5 of 14 Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	9	PRO
1	A	167	CYS
1	A	171	PRO
1	B	167	CYS
1	B	171	PRO

5.3.2 Protein sidechains (i)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles
1	A	516/516 (100%)	501 (97%)	15 (3%)	37 62
1	B	516/516 (100%)	499 (97%)	17 (3%)	33 59
All	All	1032/1032 (100%)	1000 (97%)	32 (3%)	35 60

5 of 32 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	B	520	GLN

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	B	523	ARG
1	A	423	GLN
1	A	383	THR
1	B	551	ARG

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 18 such sidechains are listed below:

Mol	Chain	Res	Type
1	B	437	ASN
1	B	556	ASN
1	B	497	ASN
1	A	556	ASN
1	B	333	ASN

5.3.3 RNA (i)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains (i)

2 non-standard protein/DNA/RNA residues are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\#\|Z\| > 2$	Counts	RMSZ	$\#\|Z\| > 2$
1	SEP	A	198	1	8,9,10	1.76	2 (25%)	7,12,14	1.85	2 (28%)
1	SEP	B	198	1	8,9,10	1.74	1 (12%)	7,12,14	1.37	1 (14%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
1	SEP	A	198	1	-	6/6/8/10	-
1	SEP	B	198	1	-	1/6/8/10	-

All (3) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	B	198	SEP	P-O1P	3.91	1.62	1.50
1	A	198	SEP	P-O1P	3.78	1.62	1.50
1	A	198	SEP	P-O2P	2.05	1.62	1.54

All (3) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(^{o})	Ideal(^{o})
1	A	198	SEP	OG-CB-CA	3.60	111.65	108.14
1	A	198	SEP	O2P-P-OG	2.58	113.40	106.67
1	B	198	SEP	OG-CB-CA	2.53	110.60	108.14

There are no chirality outliers.

5 of 7 torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
1	A	198	SEP	C-CA-CB-OG
1	A	198	SEP	CB-OG-P-O1P
1	A	198	SEP	CB-OG-P-O2P
1	A	198	SEP	CB-OG-P-O3P
1	B	198	SEP	CA-CB-OG-P

There are no ring outliers.

2 monomers are involved in 4 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
1	A	198	SEP	2	0
1	B	198	SEP	2	0

5.5 Carbohydrates (i)

22 monosaccharides are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond

length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\#\|Z\| > 2$	Counts	RMSZ	$\#\|Z\| > 2$
2	NAG	C	1	2,1	14,14,15	0.62	0	17,19,21	0.55	0
2	NAG	C	2	2	14,14,15	0.61	0	17,19,21	1.75	5 (29%)
2	MAN	C	3	2	11,11,12	0.58	0	15,15,17	1.64	1 (6%)
2	NAG	D	1	2,1	14,14,15	0.59	0	17,19,21	0.80	1 (5%)
2	NAG	D	2	2	14,14,15	0.63	0	17,19,21	0.97	1 (5%)
2	MAN	D	3	2	11,11,12	0.56	0	15,15,17	0.75	0
3	NAG	E	1	1,3	14,14,15	0.52	0	17,19,21	1.42	3 (17%)
3	NAG	E	2	3	14,14,15	0.48	0	17,19,21	1.50	2 (11%)
3	BMA	E	3	3	11,11,12	0.63	0	15,15,17	1.65	1 (6%)
4	NAG	F	1	4,1	14,14,15	0.54	0	17,19,21	1.24	2 (11%)
4	NAG	F	2	4	14,14,15	0.80	0	17,19,21	1.27	1 (5%)
2	NAG	G	1	2,1	14,14,15	0.61	0	17,19,21	1.28	2 (11%)
2	NAG	G	2	2	14,14,15	0.83	0	17,19,21	1.79	4 (23%)
2	MAN	G	3	2	11,11,12	0.67	0	15,15,17	0.98	1 (6%)
3	NAG	H	1	1,3	14,14,15	0.52	0	17,19,21	0.87	1 (5%)
3	NAG	H	2	3	14,14,15	0.65	0	17,19,21	0.93	1 (5%)
3	BMA	H	3	3	11,11,12	0.74	0	15,15,17	1.43	1 (6%)
3	NAG	I	1	1,3	14,14,15	0.56	0	17,19,21	1.06	1 (5%)
3	NAG	I	2	3	14,14,15	0.47	0	17,19,21	1.18	2 (11%)
3	BMA	I	3	3	11,11,12	0.60	0	15,15,17	1.49	2 (13%)
4	NAG	J	1	4,1	14,14,15	0.54	0	17,19,21	1.23	3 (17%)
4	NAG	J	2	4	14,14,15	0.54	0	17,19,21	0.92	1 (5%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	NAG	C	1	2,1	-	2/6/23/26	0/1/1/1
2	NAG	C	2	2	-	4/6/23/26	0/1/1/1
2	MAN	C	3	2	-	2/2/19/22	0/1/1/1
2	NAG	D	1	2,1	-	2/6/23/26	0/1/1/1

Continued on next page...

Continued from previous page...

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	NAG	D	2	2	-	2/6/23/26	0/1/1/1
2	MAN	D	3	2	-	2/2/19/22	0/1/1/1
3	NAG	E	1	1,3	-	2/6/23/26	0/1/1/1
3	NAG	E	2	3	-	2/6/23/26	0/1/1/1
3	BMA	E	3	3	-	0/2/19/22	1/1/1/1
4	NAG	F	1	4,1	-	0/6/23/26	0/1/1/1
4	NAG	F	2	4	-	1/6/23/26	0/1/1/1
2	NAG	G	1	2,1	-	4/6/23/26	0/1/1/1
2	NAG	G	2	2	-	0/6/23/26	0/1/1/1
2	MAN	G	3	2	-	2/2/19/22	0/1/1/1
3	NAG	H	1	1,3	-	0/6/23/26	0/1/1/1
3	NAG	H	2	3	-	2/6/23/26	0/1/1/1
3	BMA	H	3	3	-	2/2/19/22	1/1/1/1
3	NAG	I	1	1,3	-	3/6/23/26	0/1/1/1
3	NAG	I	2	3	-	3/6/23/26	0/1/1/1
3	BMA	I	3	3	-	0/2/19/22	1/1/1/1
4	NAG	J	1	4,1	-	2/6/23/26	0/1/1/1
4	NAG	J	2	4	-	4/6/23/26	0/1/1/1

There are no bond length outliers.

The worst 5 of 36 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(^{o})	Ideal(^{o})
3	E	3	BMA	C1-O5-C5	5.59	119.67	112.19
2	C	3	MAN	C1-O5-C5	4.98	118.86	112.19
3	H	3	BMA	C1-O5-C5	4.55	118.28	112.19
2	G	2	NAG	C4-C3-C2	4.54	117.67	111.02
3	E	2	NAG	C1-O5-C5	4.52	118.24	112.19

There are no chirality outliers.

5 of 41 torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
2	C	1	NAG	C8-C7-N2-C2
2	C	1	NAG	O7-C7-N2-C2
2	C	2	NAG	C8-C7-N2-C2
2	C	2	NAG	O7-C7-N2-C2
3	E	2	NAG	C8-C7-N2-C2

All (3) ring outliers are listed below:

Mol	Chain	Res	Type	Atoms
3	E	3	BMA	C1-C2-C3-C4-C5-O5
3	I	3	BMA	C1-C2-C3-C4-C5-O5
3	H	3	BMA	C1-C2-C3-C4-C5-O5

3 monomers are involved in 2 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	G	1	NAG	1	0
3	H	2	NAG	1	0
3	H	1	NAG	1	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for oligosaccharide.

Oligosaccharide Chain C



Oligosaccharide Chain D



Oligosaccharide Chain G



Oligosaccharide Chain E



Oligosaccharide Chain H



Oligosaccharide Chain I



Oligosaccharide Chain F



Bond lengths



Bond angles



Torsions



Rings

Oligosaccharide Chain J



5.6 Ligand geometry (i)

Of 10 ligands modelled in this entry, 2 are monoatomic - leaving 8 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\#\|Z\| > 2$	Counts	RMSZ	$\#\|Z\| > 2$
7	OSM	A	3333	-	1,3,3	0.05	0	0,2,2	-	-
6	HEM	A	3003	1	50,50,50	1.89	10 (20%)	67,82,82	1.26	6 (8%)
8	FMT	B	2004	-	2,2,2	0.70	0	1,1,1	0.32	0

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\#\|Z\| > 2$	Counts	RMSZ	$\#\|Z\| > 2$
8	FMT	A	2002	-	2,2,2	0.69	0	1,1,1	0.31	0
6	HEM	B	1003	1	50,50,50	1.93	9 (18%)	67,82,82	1.28	8 (11%)
7	OSM	B	3333	-	1,3,3	0.13	0	0,2,2	-	-
8	FMT	B	2003	-	2,2,2	0.68	0	1,1,1	0.22	0
8	FMT	A	2001	-	2,2,2	0.71	0	1,1,1	0.30	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
7	OSM	A	3333	-	-	0/0/1/1	-
6	HEM	B	1003	1	-	4/14/54/54	-
6	HEM	A	3003	1	-	4/14/54/54	-
7	OSM	B	3333	-	-	0/0/1/1	-

The worst 5 of 19 bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
6	B	1003	HEM	C3D-C2D	7.64	1.53	1.36
6	A	3003	HEM	C3D-C2D	7.57	1.53	1.36
6	A	3003	HEM	FE-NB	5.35	2.11	1.94
6	B	1003	HEM	FE-NB	4.90	2.10	1.94
6	B	1003	HEM	FE-NC	4.41	2.09	1.95

The worst 5 of 14 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(^{o})	Ideal(^{o})
6	B	1003	HEM	C4D-ND-C1D	5.37	111.56	105.21
6	A	3003	HEM	C4D-ND-C1D	5.33	111.52	105.21
6	B	1003	HEM	CBD-CAD-C3D	-2.67	105.14	112.53
6	B	1003	HEM	C3B-C2B-C1B	2.67	108.41	106.41
6	A	3003	HEM	CMD-C2D-C1D	2.47	128.89	125.03

There are no chirality outliers.

5 of 8 torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
6	B	1003	HEM	CAD-CBD-CGD-O2D
6	B	1003	HEM	CAD-CBD-CGD-O1D

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms
6	B	1003	HEM	CAA-CBA-CGA-O2A
6	B	1003	HEM	CAA-CBA-CGA-O1A
6	A	3003	HEM	CAD-CBD-CGD-O2D

There are no ring outliers.

6 monomers are involved in 38 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
7	A	3333	OSM	3	0
6	A	3003	HEM	11	0
8	A	2002	FMT	4	0
6	B	1003	HEM	17	0
7	B	3333	OSM	4	0
8	A	2001	FMT	2	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less then 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.

Ligand HEM A 3003



Bond lengths



Bond angles



Torsions



Rings

Ligand HEM B 1003



5.7 Other polymers (i)

There are no such residues in this entry.

5.8 Polymer linkage issues (i)

There are no chain breaks in this entry.

6 Fit of model and data (i)

6.1 Protein, DNA and RNA chains (i)

In the following table, the column labelled '#RSRZ> 2' contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled 'Q< 0.9' lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2			OWAB(Å^2)	Q<0.9
1	A	594/595 (99%)	0.15	28 (4%)	36	24	8, 24, 47, 72	0
1	B	594/595 (99%)	0.13	21 (3%)	47	31	4, 23, 47, 76	0
All	All	1188/1190 (99%)	0.14	49 (4%)	41	27	4, 23, 47, 76	0

The worst 5 of 49 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	6	CYS	5.8
1	A	120	GLY	4.8
1	B	172	TYR	4.6
1	B	4	VAL	4.3
1	A	121	SER	3.9

6.2 Non-standard residues in protein, DNA, RNA chains (i)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å^2)	Q<0.9
1	SEP	B	198	10/11	0.61	0.30	33,34,41,41	0
1	SEP	A	198	10/11	0.76	0.20	33,34,39,39	0

6.3 Carbohydrates (i)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å^2)	Q<0.9
2	MAN	C	3	11/12	0.37	0.17	67,68,69,69	0
2	NAG	C	1	14/15	0.45	0.19	48,53,55,59	0
3	BMA	E	3	11/12	0.46	0.17	59,60,61,61	0
4	NAG	J	2	14/15	0.46	0.16	62,63,64,64	0
2	NAG	C	2	14/15	0.47	0.16	62,65,65,67	0
2	MAN	G	3	11/12	0.49	0.17	71,71,72,72	0
3	NAG	E	2	14/15	0.50	0.20	53,54,56,58	0
3	BMA	H	3	11/12	0.51	0.20	66,68,68,68	0
4	NAG	F	2	14/15	0.52	0.18	55,57,57,57	0
2	MAN	D	3	11/12	0.53	0.22	59,60,61,61	0
4	NAG	F	1	14/15	0.56	0.17	43,46,48,51	0
3	BMA	I	3	11/12	0.57	0.19	65,66,66,67	0
3	NAG	H	2	14/15	0.61	0.16	57,58,61,64	0
3	NAG	I	2	14/15	0.66	0.15	60,62,63,64	0
4	NAG	J	1	14/15	0.67	0.16	50,55,56,59	0
2	NAG	G	2	14/15	0.70	0.16	63,66,67,69	0
2	NAG	G	1	14/15	0.73	0.14	46,51,54,58	0
2	NAG	D	2	14/15	0.76	0.13	51,52,55,57	0
3	NAG	E	1	14/15	0.83	0.14	42,44,46,50	0
3	NAG	I	1	14/15	0.83	0.14	48,50,53,56	0
3	NAG	H	1	14/15	0.85	0.13	44,46,49,53	0
2	NAG	D	1	14/15	0.91	0.12	38,41,43,47	0

The following is a graphical depiction of the model fit to experimental electron density for oligosaccharide. Each fit is shown from different orientation to approximate a three-dimensional view.

Electron density around Chain C:

$2mF_o$-DF_c (at 0.7 rmsd) in gray
mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)







Electron density around Chain D:

$2mF_o$-DF_c (at 0.7 rmsd) in gray
mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)







Electron density around Chain G:

$2mF_o$-DF_c (at 0.7 rmsd) in gray
mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)







Electron density around Chain E:

$2mF_o$-DF_c (at 0.7 rmsd) in gray
mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)



Electron density around Chain H:

$2mF_o$-DF_c (at 0.7 rmsd) in gray
mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)



Electron density around Chain I:

$2mF_o$-DF_c (at 0.7 rmsd) in gray
mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)



Electron density around Chain F:

$2mF_o$-DF_c (at 0.7 rmsd) in gray
mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)







Electron density around Chain J:

$2mF_o$-DF_c (at 0.7 rmsd) in gray
mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)



6.4 Ligands (i)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å^2)	Q<0.9
7	OSM	B	3333	4/4	0.68	0.27	54,54,54,54	0
7	OSM	A	3333	4/4	0.73	0.25	48,48,48,48	0
6	HEM	B	1003	43/43	0.79	0.21	12,13,14,15	0
6	HEM	A	3003	43/43	0.82	0.19	17,18,22,23	0
5	CA	A	607	1/1	0.90	0.07	25,25,25,25	0
8	FMT	A	2002	3/3	0.92	0.17	32,32,32,32	0
8	FMT	A	2001	3/3	0.94	0.15	42,42,42,42	0
8	FMT	B	2003	3/3	0.94	0.12	46,46,46,46	0
8	FMT	B	2004	3/3	0.94	0.14	29,29,29,29	0
5	CA	B	607	1/1	0.97	0.06	34,34,34,34	0

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.

Electron density around HEM B 1003:

$2mF_o$-DF_c (at 0.7 rmsd) in gray
mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)



Electron density around HEM A 3003: 2mF$_o$-DF$_c$ (at 0.7 rmsd) in gray mF$_o$-DF$_c$ (at 3 rmsd) in purple (negative) and green (positive)	
	

6.5 Other polymers (i)

There are no such residues in this entry.