



wwPDB X-ray Structure Validation Summary Report ⓘ

Mar 5, 2026 – 12:32 PM UTC

PDB ID : 1NMT / pdb_00001nmt
Title : N-MYRISTOYL TRANSFERASE FROM CANDIDA ALBICANS AT 2.45 Å
Authors : Weston, S.A.; Pauptit, R.A.
Deposited on : 1997-12-11
Resolution : 2.45 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4-5-2 with Phenix2.0
Mogul : 2022.3.0, CSD as543be (2022)
Xtriage (Phenix) : 2.0
EDS : 3.0
Percentile statistics : 20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4 : 9.0.010 (Gargrove)
Density-Fitness : 1.0.12
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.45 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	1190 (2.46-2.46)
Clashscore	190562	1229 (2.46-2.46)
Ramachandran outliers	187476	1218 (2.46-2.46)
Sidechain outliers	187428	1218 (2.46-2.46)
RSRZ outliers	180081	1190 (2.46-2.46)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	392	 14% 70% 27% 5% 2%
1	B	392	 45% 71% 23% 5% 2%
1	C	392	 49% 69% 27% 5% 2%

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
2	GOL	B	12	-	-	X	-

2 Entry composition [i](#)


There are 3 unique types of molecules in this entry. The entry contains 10001 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called N-MYRISTOYL TRANSFERASE.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	392	Total 3210	C 2079	N 524	O 598	S 9	0	0	0
1	B	392	Total 3207	C 2078	N 524	O 596	S 9	0	0	0
1	C	389	Total 3185	C 2064	N 521	O 591	S 9	0	0	0

- Molecule 2 is GLYCEROL (CCD ID: GOL) (formula: C₃H₈O₃).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
			Total	C	O		
2	A	1	Total 6	C 3	O 3	0	0
2	A	1	Total 6	C 3	O 3	0	0
2	A	1	Total 6	C 3	O 3	0	0


Continued on next page...

Continued from previous page...


Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
2	A	1	Total C O 6 3 3	0	0
2	A	1	Total C O 6 3 3	0	0
2	A	1	Total C O 6 3 3	0	0
2	A	1	Total C O 6 3 3	0	0
2	A	1	Total C O 6 3 3	0	0
2	A	1	Total C O 6 3 3	0	0
2	B	1	Total C O 6 3 3	0	0
2	B	1	Total C O 6 3 3	0	0
2	B	1	Total C O 6 3 3	0	0
2	B	1	Total C O 6 3 3	0	0
2	B	1	Total C O 6 3 3	0	0
2	C	1	Total C O 6 3 3	0	0
2	C	1	Total C O 6 3 3	0	0
2	C	1	Total C O 6 3 3	0	0
2	C	1	Total C O 6 3 3	0	0

- Molecule 3 is water.

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
3	A	105	Total O 105 105	0	0
3	B	150	Total O 150 150	0	0
3	C	36	Total O 36 36	0	0



● Molecule 1: N-MYRISTOYL TRANSFERASE



4 Data and refinement statistics

Property	Value	Source
Space group	C 2 2 21	Depositor
Cell constants a, b, c, α , β , γ	132.00Å 166.50Å 179.40Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	20.85 – 2.45 20.85 – 2.45	Depositor EDS
% Data completeness (in resolution range)	98.3 (20.85-2.45) 98.6 (20.85-2.45)	Depositor EDS
R_{merge}	0.05	Depositor
R_{sym}	0.05	Depositor
$\langle I/\sigma(I) \rangle$ ¹	2.68 (at 2.44Å)	Xtrriage
Refinement program	X-PLOR 3.98	Depositor
R, R_{free}	0.214 , 0.251 0.299 , 0.316	Depositor DCC
R_{free} test set	3703 reflections (5.10%)	wwPDB-VP
Wilson B-factor (Å ²)	36.9	Xtrriage
Anisotropy	0.218	Xtrriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.35 , 41.4	EDS
L-test for twinning ²	$\langle L \rangle = 0.49$, $\langle L^2 \rangle = 0.32$	Xtrriage
Estimated twinning fraction	No twinning to report.	Xtrriage
F_o, F_c correlation	0.86	EDS
Total number of atoms	10001	wwPDB-VP
Average B, all atoms (Å ²)	38.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 2.99% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: GOL

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# $ Z > 5$	RMSZ	# $ Z > 5$
1	A	0.63	0/3296	1.04	20/4473 (0.4%)
1	B	0.70	1/3293 (0.0%)	1.05	20/4469 (0.4%)
1	C	0.56	0/3270	1.01	16/4436 (0.4%)
All	All	0.63	1/9859 (0.0%)	1.03	56/13378 (0.4%)

All (1) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	B	218	THR	CA-CB	7.34	1.57	1.52

The worst 5 of 56 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	296	THR	N-CA-C	-9.90	95.86	110.52
1	C	296	THR	N-CA-C	-9.22	97.21	110.59
1	B	296	THR	N-CA-C	-9.15	97.32	110.59
1	C	230	ILE	N-CA-C	-7.97	105.44	111.90
1	B	139	VAL	N-CA-C	7.91	120.08	107.73

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	3210	0	3167	70	6
1	B	3207	0	3165	87	5
1	C	3185	0	3145	77	4
2	A	54	0	72	12	0
2	B	30	0	40	8	0
2	C	24	0	32	6	0
3	A	105	0	0	3	0
3	B	150	0	0	3	1
3	C	36	0	0	0	0
All	All	10001	0	9621	228	10

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 12.

The worst 5 of 228 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:C:265:LYS:H	1:C:265:LYS:HE2	1.26	1.00
1:B:215:ILE:HD12	1:C:276:SER:HB2	1.44	0.98
1:B:265:LYS:HE2	1:B:265:LYS:H	1.27	0.97
1:A:235:LEU:HD21	1:A:390:VAL:HG22	1.49	0.94
1:A:265:LYS:H	1:A:265:LYS:HE2	1.30	0.94

The worst 5 of 10 symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:148:LYS:NZ	1:B:348:ASP:OD1[3_655]	1.35	0.85
1:A:145:SER:OG	1:C:346:GLN:OE1[7_555]	1.76	0.44
3:B:455:HOH:O	3:B:556:HOH:O[3_655]	1.78	0.42
1:B:148:LYS:CE	1:B:348:ASP:OD1[3_655]	1.94	0.26
1:A:348:ASP:OD1	1:C:148:LYS:NZ[7_555]	2.04	0.16

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	390/392 (100%)	367 (94%)	22 (6%)	1 (0%)	36	45
1	B	390/392 (100%)	368 (94%)	18 (5%)	4 (1%)	12	14
1	C	385/392 (98%)	364 (94%)	18 (5%)	3 (1%)	16	20
All	All	1165/1176 (99%)	1099 (94%)	58 (5%)	8 (1%)	18	24

5 of 8 Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	B	109	GLU
1	A	245	PRO
1	C	362	SER
1	B	65	LYS
1	B	245	PRO

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	360/360 (100%)	332 (92%)	28 (8%)	11	14
1	B	359/360 (100%)	332 (92%)	27 (8%)	12	16
1	C	357/360 (99%)	326 (91%)	31 (9%)	9	11
All	All	1076/1080 (100%)	990 (92%)	86 (8%)	11	13

5 of 86 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	C	149	LEU
1	C	284	LYS
1	C	172	VAL
1	C	249	LYS
1	C	331	TYR

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 36 such sidechains are listed below:

Mol	Chain	Res	Type
1	C	260	ASN
1	C	421	ASN
1	C	286	GLN
1	C	367	ASN
1	B	95	GLN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

18 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
2	GOL	B	17	-	5,5,5	0.49	0	5,5,5	0.93	0
2	GOL	C	6	-	5,5,5	0.67	0	5,5,5	1.01	0
2	GOL	C	10	-	5,5,5	0.61	0	5,5,5	0.95	0
2	GOL	B	12	-	5,5,5	0.71	0	5,5,5	1.11	0
2	GOL	A	9	-	5,5,5	0.79	0	5,5,5	1.00	0
2	GOL	A	16	-	5,5,5	0.51	0	5,5,5	0.91	0
2	GOL	B	2	-	5,5,5	0.44	0	5,5,5	0.89	0

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
2	GOL	B	11	-	5,5,5	0.61	0	5,5,5	0.96	0
2	GOL	A	15	-	5,5,5	0.42	0	5,5,5	0.84	0
2	GOL	C	14	-	5,5,5	0.67	0	5,5,5	1.05	0
2	GOL	A	8	-	5,5,5	0.55	0	5,5,5	0.91	0
2	GOL	B	1	-	5,5,5	0.40	0	5,5,5	0.83	0
2	GOL	A	3	-	5,5,5	0.49	0	5,5,5	0.95	0
2	GOL	A	4	-	5,5,5	0.19	0	5,5,5	0.84	0
2	GOL	A	18	-	5,5,5	0.47	0	5,5,5	0.95	0
2	GOL	C	5	-	5,5,5	0.50	0	5,5,5	0.83	0
2	GOL	A	7	-	5,5,5	0.48	0	5,5,5	0.85	0
2	GOL	A	13	-	5,5,5	0.56	0	5,5,5	1.06	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	GOL	B	17	-	-	4/4/4/4	-
2	GOL	C	6	-	-	0/4/4/4	-
2	GOL	C	10	-	-	2/4/4/4	-
2	GOL	B	12	-	-	2/4/4/4	-
2	GOL	A	9	-	-	2/4/4/4	-
2	GOL	A	16	-	-	1/4/4/4	-
2	GOL	B	2	-	-	0/4/4/4	-
2	GOL	B	11	-	-	2/4/4/4	-
2	GOL	A	15	-	-	1/4/4/4	-
2	GOL	C	14	-	-	4/4/4/4	-
2	GOL	A	8	-	-	2/4/4/4	-
2	GOL	B	1	-	-	3/4/4/4	-
2	GOL	A	3	-	-	4/4/4/4	-
2	GOL	A	4	-	-	2/4/4/4	-
2	GOL	A	18	-	-	2/4/4/4	-
2	GOL	C	5	-	-	2/4/4/4	-
2	GOL	A	7	-	-	2/4/4/4	-
2	GOL	A	13	-	-	0/4/4/4	-

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

5 of 35 torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
2	A	3	GOL	O1-C1-C2-C3
2	A	3	GOL	C1-C2-C3-O3
2	A	4	GOL	C1-C2-C3-O3
2	A	7	GOL	O1-C1-C2-C3
2	A	8	GOL	C1-C2-C3-O3

There are no ring outliers.

15 monomers are involved in 26 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	B	17	GOL	2	0
2	C	6	GOL	1	0
2	C	10	GOL	2	0
2	B	12	GOL	5	0
2	A	9	GOL	2	0
2	A	16	GOL	3	0
2	B	11	GOL	1	0
2	A	15	GOL	1	0
2	C	14	GOL	2	0
2	A	8	GOL	2	0
2	A	4	GOL	2	0
2	A	18	GOL	1	0
2	C	5	GOL	1	0
2	A	7	GOL	1	0
2	A	13	GOL	2	0

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	392/392 (100%)	1.11	55 (14%) 6 5	20, 34, 59, 88	0
1	B	392/392 (100%)	1.97	175 (44%) 0 0	15, 28, 55, 90	0
1	C	389/392 (99%)	2.15	194 (49%) 0 0	27, 44, 69, 89	0
All	All	1173/1176 (99%)	1.74	424 (36%) 1 0	15, 35, 64, 90	0

The worst 5 of 424 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	C	66	LEU	7.0
1	B	244	PRO	6.1
1	C	108	VAL	5.9
1	C	345	ALA	5.9
1	A	66	LEU	5.4

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled ‘Q < 0.9’ lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(\AA^2)	Q<0.9
2	GOL	C	14	6/6	0.57	0.34	79,81,83,84	0
2	GOL	B	11	6/6	0.65	0.24	52,55,59,66	0
2	GOL	B	12	6/6	0.66	0.29	61,69,71,74	0
2	GOL	A	16	6/6	0.67	0.22	78,79,80,80	0
2	GOL	A	9	6/6	0.70	0.20	36,49,55,59	0
2	GOL	C	6	6/6	0.71	0.21	61,63,71,73	0
2	GOL	B	17	6/6	0.72	0.35	75,80,82,83	0
2	GOL	A	15	6/6	0.73	0.24	63,72,75,86	0
2	GOL	A	13	6/6	0.74	0.24	61,67,67,70	0
2	GOL	C	10	6/6	0.75	0.18	41,51,55,56	0
2	GOL	A	7	6/6	0.75	0.19	45,46,50,59	0
2	GOL	B	1	6/6	0.80	0.15	32,34,38,42	0
2	GOL	C	5	6/6	0.81	0.19	52,55,57,61	0
2	GOL	A	8	6/6	0.81	0.16	32,43,46,50	0
2	GOL	B	2	6/6	0.82	0.13	40,49,54,56	0
2	GOL	A	3	6/6	0.84	0.17	39,46,50,52	0
2	GOL	A	18	6/6	0.88	0.13	32,46,47,59	0
2	GOL	A	4	6/6	0.89	0.14	32,36,45,46	0

6.5 Other polymers [i](#)

There are no such residues in this entry.