



Full wwPDB X-ray Structure Validation Report ⓘ

Mar 10, 2026 – 08:52 PM UTC

PDB ID : 3NMO / pdb_00003nmo
Title : Crystal structure of an engineered monomeric CLC-ec1 Cl⁻/H⁺ transporter
Authors : Robertson, J.L.; Kolmakova-Partensky, L.; Miller, C.
Deposited on : 2010-06-22
Resolution : 3.10 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4-5-2 with Phenix2.0
Xtrriage (Phenix) : 2.0
EDS : 3.0
Percentile statistics : 20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4 : 9.0.010 (Gargrove)
Density-Fitness : 1.0.12
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 3.10 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	1456 (3.10-3.10)
Clashscore	190562	1539 (3.10-3.10)
Ramachandran outliers	187476	1467 (3.10-3.10)
Sidechain outliers	187428	1467 (3.10-3.10)
RSRZ outliers	180081	1456 (3.10-3.10)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	465	

2 Entry composition

There are 2 unique types of molecules in this entry. The entry contains 3293 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called H(+)/Cl(-) exchange transporter clcA.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	439	3292	2168	547	557	20	0	0	0

There are 2 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	201	TRP	ILE	engineered mutation	UNP P37019
A	422	TRP	ILE	engineered mutation	UNP P37019


- Molecule 2 is CHLORIDE ION (CCD ID: CL) (formula: Cl).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
2	A	1	Total	Cl	0	0
			1	1		

3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: H(+)/Cl(-) exchange transporter clcA



4 Data and refinement statistics

Property	Value	Source
Space group	C 1 2 1	Depositor
Cell constants a, b, c, α , β , γ	200.75Å 60.33Å 70.03Å 90.00° 94.27° 90.00°	Depositor
Resolution (Å)	44.99 – 3.10 44.99 – 3.10	Depositor EDS
% Data completeness (in resolution range)	98.3 (44.99-3.10) 98.3 (44.99-3.10)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	2.69 (at 3.12Å)	Xtrriage
Refinement program	REFMAC 5.5.0109	Depositor
R, R_{free}	0.263 , 0.303 (Not available) , 0.290	Depositor DCC
R_{free} test set	777 reflections (5.03%)	wwPDB-VP
Wilson B-factor (Å ²)	96.7	Xtrriage
Anisotropy	0.092	Xtrriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.28 , 54.6	EDS
L-test for twinning ²	$\langle L \rangle = 0.50$, $\langle L^2 \rangle = 0.34$	Xtrriage
Estimated twinning fraction	No twinning to report.	Xtrriage
F_o, F_c correlation	0.87	EDS
Total number of atoms	3293	wwPDB-VP
Average B, all atoms (Å ²)	93.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 7.07% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: CL

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.59	1/3368 (0.0%)	0.94	1/4576 (0.0%)

All (1) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	320	ILE	CA-CB	5.25	1.57	1.54

All (1) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	30	LYS	N-CA-C	5.26	117.85	109.96

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	3292	0	3415	51	0
2	A	1	0	0	1	0
All	All	3293	0	3415	51	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 8.

All (51) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:107:SER:OG	2:A:474:CL:CL	2.00	1.14
1:A:148:GLU:H	1:A:148:GLU:CD	1.92	0.76
1:A:28:ARG:O	1:A:28:ARG:HG3	1.85	0.75
1:A:100:TYR:O	1:A:126:ARG:HD3	1.92	0.69
1:A:147:ARG:NH2	1:A:189:ALA:O	2.26	0.68
1:A:148:GLU:HG2	1:A:357:PHE:HB3	1.78	0.65
1:A:38:MET:HG3	1:A:168:LEU:HD11	1.81	0.63
1:A:28:ARG:C	1:A:29:ASP:OD1	2.48	0.57
1:A:94:TYR:CZ	1:A:352:ALA:HB2	2.42	0.54
1:A:38:MET:HA	1:A:38:MET:HE2	1.88	0.54
1:A:47:GLY:HA3	1:A:228:MET:HG3	1.89	0.54
1:A:229:TYR:C	1:A:229:TYR:CD2	2.86	0.54
1:A:28:ARG:O	1:A:29:ASP:OD1	2.26	0.53
1:A:278:ASP:CG	1:A:451:ARG:HH22	2.16	0.53
1:A:95:PHE:CZ	1:A:99:LYS:HG3	2.44	0.53
1:A:54:ASP:OD1	1:A:147:ARG:HD2	2.07	0.52
1:A:228:MET:HA	1:A:228:MET:HE2	1.93	0.51
1:A:241:VAL:HG11	1:A:391:ILE:HD11	1.93	0.51
1:A:54:ASP:CG	1:A:147:ARG:HH11	2.20	0.50
1:A:123:ARG:HH21	1:A:126:ARG:HD2	1.77	0.49
1:A:413:LEU:HD13	1:A:425:MET:HE1	1.94	0.49
1:A:360:MET:HE1	1:A:401:SER:HB3	1.95	0.49
1:A:394:MET:HE2	1:A:412:VAL:HG13	1.96	0.48
1:A:305:LEU:HB3	1:A:313:SER:HB2	1.95	0.48
1:A:408:GLY:O	1:A:412:VAL:HG23	2.14	0.47
1:A:138:THR:HG21	1:A:352:ALA:HB1	1.96	0.47
1:A:382:TYR:O	1:A:383:HIS:HB2	2.15	0.47
1:A:118:ASP:CG	1:A:174:ARG:HH21	2.23	0.45
1:A:443:PRO:HB2	1:A:446:SER:HB2	1.97	0.45
1:A:78:LEU:HD21	1:A:307:PHE:CZ	2.52	0.45
1:A:86:SER:OG	1:A:303:GLY:HA3	2.17	0.45
1:A:163:LEU:HD12	1:A:168:LEU:HB2	1.99	0.44
1:A:122:VAL:O	1:A:122:VAL:HG12	2.18	0.44
1:A:97:VAL:HG21	1:A:353:PRO:HD3	1.99	0.44
1:A:190:PHE:HE2	1:A:317:PHE:HZ	1.65	0.44
1:A:420:GLN:NE2	1:A:421:LEU:HG	2.33	0.44
1:A:59:TRP:O	1:A:63:GLN:HG2	2.19	0.43
1:A:197:ILE:O	1:A:201:TRP:HD1	2.01	0.43
1:A:54:ASP:OD2	1:A:147:ARG:NH1	2.51	0.43
1:A:61:GLN:HG2	1:A:64:ARG:HH21	1.83	0.43

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:109:ILE:N	1:A:110:PRO:CD	2.81	0.42
1:A:28:ARG:O	1:A:28:ARG:CG	2.58	0.42
1:A:98:ARG:HD2	1:A:291:TRP:CD2	2.54	0.42
1:A:251:THR:HG23	1:A:254:LEU:HD12	2.02	0.42
1:A:225:SER:O	1:A:228:MET:HB2	2.19	0.42
1:A:274:LEU:HD22	1:A:451:ARG:HD2	2.02	0.41
1:A:183:ALA:HB2	1:A:200:ILE:CG1	2.50	0.41
1:A:108:GLY:HA3	1:A:153:GLN:OE1	2.20	0.41
1:A:64:ARG:NH1	1:A:141:GLY:O	2.54	0.41
1:A:46:VAL:HG22	1:A:155:GLY:HA2	2.03	0.40
1:A:131:LYS:HD3	1:A:153:GLN:OE1	2.21	0.40

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
1	A	437/465 (94%)	414 (95%)	19 (4%)	4 (1%)	14 44

All (4) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	235	GLU
1	A	234	HIS
1	A	383	HIS
1	A	222	VAL

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar

resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles
1	A	131/353 (37%)	127 (97%)	4 (3%)	35 64

All (4) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	98	ARG
1	A	99	LYS
1	A	107	SER
1	A	413	LEU

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. There are no such sidechains identified.

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

Of 1 ligands modelled in this entry, 1 is monoatomic - leaving 0 for Mogul analysis.

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	439/465 (94%)	1.17	96 (21%) 2 1	67, 91, 117, 128	0

All (96) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	351	GLY	9.5
1	A	348	PHE	7.2
1	A	356	ILE	6.5
1	A	212	LEU	6.5
1	A	385	GLU	6.0
1	A	388	THR	6.0
1	A	452	THR	5.7
1	A	347	CYS	5.5
1	A	246	ASP	5.5
1	A	318	ASN	5.4
1	A	171	ASP	5.4
1	A	172	GLU	5.3
1	A	235	GLU	5.1
1	A	354	GLY	5.0
1	A	445	TYR	4.8
1	A	76	PRO	4.8
1	A	245	SER	4.7
1	A	386	ALA	4.7
1	A	373	MET	4.6
1	A	355	GLY	4.6
1	A	107	SER	4.6
1	A	302	CYS	4.5
1	A	383	HIS	4.4
1	A	213	ILE	4.4
1	A	421	LEU	4.3
1	A	387	GLY	4.2
1	A	316	GLY	4.1

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	A	392	ALA	4.1
1	A	211	THR	4.0
1	A	330	MET	4.0
1	A	247	ALA	3.9
1	A	201	TRP	3.9
1	A	357	PHE	3.9
1	A	420	GLN	3.9
1	A	34	ALA	3.8
1	A	234	HIS	3.7
1	A	438	PHE	3.7
1	A	294	MET	3.5
1	A	424	PRO	3.5
1	A	85	CYS	3.4
1	A	237	ALA	3.4
1	A	448	ILE	3.4
1	A	360	MET	3.3
1	A	166	PHE	3.3
1	A	106	GLY	3.3
1	A	400	ALA	3.3
1	A	352	ALA	3.3
1	A	236	VAL	3.2
1	A	337	PHE	3.2
1	A	29	ASP	3.1
1	A	349	SER	3.1
1	A	91	MET	3.1
1	A	204	MET	3.1
1	A	161	MET	3.1
1	A	108	GLY	3.1
1	A	65	MET	3.0
1	A	148	GLU	2.9
1	A	432	ALA	2.9
1	A	267	PRO	2.9
1	A	317	PHE	2.8
1	A	394	MET	2.8
1	A	350	SER	2.8
1	A	36	LEU	2.7
1	A	79	LEU	2.7
1	A	208	PHE	2.6
1	A	105	GLY	2.6
1	A	459	GLU	2.6
1	A	391	ILE	2.5
1	A	238	LEU	2.5

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	A	332	MET	2.5
1	A	30	LYS	2.5
1	A	32	PRO	2.5
1	A	33	LEU	2.5
1	A	276	MET	2.5
1	A	328	PHE	2.5
1	A	227	ILE	2.4
1	A	239	ILE	2.4
1	A	456	GLN	2.3
1	A	252	LEU	2.3
1	A	250	ASN	2.2
1	A	407	THR	2.2
1	A	206	PRO	2.2
1	A	256	LEU	2.2
1	A	460	GLN	2.2
1	A	209	ARG	2.2
1	A	228	MET	2.2
1	A	425	MET	2.2
1	A	210	TYR	2.1
1	A	345	LEU	2.1
1	A	165	ILE	2.1
1	A	25	LEU	2.1
1	A	80	THR	2.1
1	A	320	ILE	2.1
1	A	22	ILE	2.1
1	A	31	THR	2.0
1	A	109	ILE	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum,

median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
2	CL	A	474	1/1	0.77	0.40	72,72,72,72	0

6.5 Other polymers [i](#)

There are no such residues in this entry.