



wwPDB X-ray Structure Validation Summary Report ⓘ

Mar 9, 2026 – 04:03 AM UTC

PDB ID : 1NTO / pdb_00001nto
Title : N249Y MUTANT OF ALCOHOL DEHYDROGENASE FROM THE ARCHAEON SULFOLOBUS SOLFATARICUS-MONOCLINIC CRYSTAL FORM
Authors : Esposito, L.; Bruno, I.; Sica, F.; Raia, C.A.; Giordano, A.; Rossi, M.; Mazzarella, L.; Zagari, A.
Deposited on : 2003-01-30
Resolution : 1.94 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4-5-2 with Phenix2.0
Xtriage (Phenix)	:	2.0
EDS	:	3.0
Percentile statistics	:	20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4	:	9.0.010 (Gargrove)
Density-Fitness	:	1.0.12
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 1.94 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	1452 (1.94-1.94)
Clashscore	190562	1494 (1.94-1.94)
Ramachandran outliers	187476	1479 (1.94-1.94)
Sidechain outliers	187428	1479 (1.94-1.94)
RSRZ outliers	180081	1453 (1.94-1.94)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	347	16% 78% 20%
1	B	347	18% 75% 24%
1	C	347	19% 79% 20%
1	D	347	15% 79% 19%
1	E	347	12% 78% 21%

Continued on next page...

Continued from previous page...

Mol	Chain	Length	Quality of chain
1	H	347	16% 78% 21%

2 Entry composition [i](#)

There are 3 unique types of molecules in this entry. The entry contains 16354 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called NAD-dependent alcohol dehydrogenase.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	347	Total	C	N	O	S	0	0	0
			2620	1666	454	486	14			
1	B	347	Total	C	N	O	S	0	0	0
			2604	1656	448	486	14			
1	C	347	Total	C	N	O	S	0	0	0
			2617	1665	450	488	14			
1	D	347	Total	C	N	O	S	0	0	0
			2620	1668	452	486	14			
1	E	347	Total	C	N	O	S	0	0	0
			2628	1671	455	488	14			
1	H	347	Total	C	N	O	S	0	0	0
			2620	1667	451	488	14			

There are 6 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	249	TYR	ASN	engineered mutation	UNP P39462
B	249	TYR	ASN	engineered mutation	UNP P39462
C	249	TYR	ASN	engineered mutation	UNP P39462
D	249	TYR	ASN	engineered mutation	UNP P39462
E	249	TYR	ASN	engineered mutation	UNP P39462
H	249	TYR	ASN	engineered mutation	UNP P39462

- Molecule 2 is ZINC ION (CCD ID: ZN) (formula: Zn).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
2	A	2	Total	Zn	0	0
			2	2		
2	B	2	Total	Zn	0	0
			2	2		
2	C	2	Total	Zn	0	0
			2	2		

Continued on next page...

Continued from previous page...

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
2	D	2	Total 2	Zn 2	0	0
2	E	2	Total 2	Zn 2	0	0
2	H	2	Total 2	Zn 2	0	0


- Molecule 3 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	A	142	Total 142	O 142	0	0
3	B	83	Total 83	O 83	0	0
3	C	92	Total 92	O 92	0	0
3	D	138	Total 138	O 138	0	0
3	E	101	Total 101	O 101	0	0
3	H	77	Total 77	O 77	0	0


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: NAD-dependent alcohol dehydrogenase




- Molecule 1: NAD-dependent alcohol dehydrogenase




- Molecule 1: NAD-dependent alcohol dehydrogenase







● Molecule 1: NAD-dependent alcohol dehydrogenase



● Molecule 1: NAD-dependent alcohol dehydrogenase



● Molecule 1: NAD-dependent alcohol dehydrogenase



4 Data and refinement statistics

Property	Value	Source
Space group	C 1 2 1	Depositor
Cell constants a, b, c, α , β , γ	215.33Å 91.59Å 118.24Å 90.00° 99.20° 90.00°	Depositor
Resolution (Å)	15.00 – 1.94 15.00 – 1.94	Depositor EDS
% Data completeness (in resolution range)	86.9 (15.00-1.94) 86.8 (15.00-1.94)	Depositor EDS
R_{merge}	0.03	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	9.52 (at 1.94Å)	Xtriage
Refinement program	CNS 1.0	Depositor
R, R_{free}	0.210 , 0.244 0.210 , 0.244	Depositor DCC
R_{free} test set	7421 reflections (5.06%)	wwPDB-VP
Wilson B-factor (Å ²)	23.5	Xtriage
Anisotropy	0.241	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.43 , 54.4	EDS
L-test for twinning ²	$\langle L \rangle = 0.49$, $\langle L^2 \rangle = 0.32$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.93	EDS
Total number of atoms	16354	wwPDB-VP
Average B, all atoms (Å ²)	36.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 3.49% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: ZN

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# $ Z > 5$	RMSZ	# $ Z > 5$
1	A	0.63	1/2665 (0.0%)	0.94	2/3606 (0.1%)
1	B	0.55	0/2648	0.90	5/3585 (0.1%)
1	C	0.55	1/2662 (0.0%)	0.93	7/3604 (0.2%)
1	D	0.62	0/2665	0.93	4/3605 (0.1%)
1	E	0.54	0/2673	0.91	3/3615 (0.1%)
1	H	0.55	0/2665	0.90	3/3606 (0.1%)
All	All	0.57	2/15978 (0.0%)	0.92	24/21621 (0.1%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	E	0	1

All (2) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	269	MET	SD-CE	-6.57	1.63	1.79
1	C	269	MET	SD-CE	-5.13	1.66	1.79

The worst 5 of 24 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	195	SER	N-CA-C	7.47	123.30	114.04
1	B	194	VAL	N-CA-C	7.33	121.42	113.43
1	C	195	SER	N-CA-C	7.08	123.03	113.97
1	H	194	VAL	N-CA-C	7.04	119.93	113.10
1	E	70	ILE	N-CA-C	7.00	118.20	108.12

There are no chirality outliers.

All (1) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	E	103	TYR	Sidechain

5.2 Too-close contacts ⓘ

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2620	0	2670	59	0
1	B	2604	0	2645	72	0
1	C	2617	0	2657	61	0
1	D	2620	0	2674	66	0
1	E	2628	0	2685	59	0
1	H	2620	0	2667	54	0
2	A	2	0	0	0	0
2	B	2	0	0	0	0
2	C	2	0	0	0	0
2	D	2	0	0	0	0
2	E	2	0	0	0	0
2	H	2	0	0	0	0
3	A	142	0	0	4	0
3	B	83	0	0	2	0
3	C	92	0	0	2	0
3	D	138	0	0	1	0
3	E	101	0	0	1	0
3	H	77	0	0	1	0
All	All	16354	0	15998	360	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 11.

The worst 5 of 360 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:E:8:GLU:HB3	1:E:11:LYS:HE2	1.51	0.93
1:B:42:VAL:HG13	1:B:335:GLU:HG2	1.55	0.88

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:D:54:ILE:HB	1:D:60:VAL:HG21	1.58	0.86
1:E:192:LYS:HE3	1:E:216:GLY:O	1.78	0.84
1:B:54:ILE:HB	1:B:60:VAL:HG21	1.58	0.83

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	345/347 (99%)	333 (96%)	10 (3%)	2 (1%)	21	11
1	B	345/347 (99%)	325 (94%)	18 (5%)	2 (1%)	21	11
1	C	345/347 (99%)	330 (96%)	12 (4%)	3 (1%)	14	6
1	D	345/347 (99%)	330 (96%)	13 (4%)	2 (1%)	21	11
1	E	345/347 (99%)	333 (96%)	10 (3%)	2 (1%)	21	11
1	H	345/347 (99%)	327 (95%)	16 (5%)	2 (1%)	21	11
All	All	2070/2082 (99%)	1978 (96%)	79 (4%)	13 (1%)	21	11

5 of 13 Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	B	51	ASN
1	C	51	ASN
1	H	51	ASN
1	A	51	ASN
1	A	296	VAL

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	273/278 (98%)	268 (98%)	5 (2%)	51	43
1	B	270/278 (97%)	263 (97%)	7 (3%)	40	28
1	C	272/278 (98%)	268 (98%)	4 (2%)	57	49
1	D	273/278 (98%)	267 (98%)	6 (2%)	45	35
1	E	275/278 (99%)	267 (97%)	8 (3%)	37	24
1	H	273/278 (98%)	267 (98%)	6 (2%)	45	35
All	All	1636/1668 (98%)	1600 (98%)	36 (2%)	45	35

5 of 36 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	E	292	VAL
1	H	338	LYS
1	E	338	LYS
1	H	183	LEU
1	C	183	LEU

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 20 such sidechains are listed below:

Mol	Chain	Res	Type
1	E	278	HIS
1	H	16	GLN
1	H	336	ASN
1	H	290	GLN
1	C	333	ASN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

Of 12 ligands modelled in this entry, 12 are monoatomic - leaving 0 for Mogul analysis.

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	347/347 (100%)	0.60	55 (15%) 5 5	12, 26, 72, 87	0
1	B	347/347 (100%)	0.90	62 (17%) 3 4	17, 35, 75, 88	0
1	C	347/347 (100%)	0.87	65 (18%) 3 3	16, 32, 74, 89	0
1	D	347/347 (100%)	0.59	51 (14%) 6 6	12, 27, 73, 88	0
1	E	347/347 (100%)	0.62	42 (12%) 8 10	17, 31, 74, 88	0
1	H	347/347 (100%)	0.74	56 (16%) 4 5	16, 33, 75, 89	0
All	All	2082/2082 (100%)	0.72	331 (15%) 5 5	12, 31, 75, 89	0

The worst 5 of 331 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	C	47	GLY	11.6
1	E	340	ILE	10.3
1	A	338	LYS	7.6
1	C	48	ARG	7.6
1	B	59	GLY	7.4

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
2	ZN	B	500	1/1	0.94	0.06	41,41,41,41	0
2	ZN	A	500	1/1	0.98	0.05	28,28,28,28	0
2	ZN	C	500	1/1	0.98	0.05	33,33,33,33	0
2	ZN	E	500	1/1	0.98	0.04	32,32,32,32	0
2	ZN	H	500	1/1	0.98	0.04	35,35,35,35	0
2	ZN	B	400	1/1	0.99	0.03	24,24,24,24	0
2	ZN	D	500	1/1	0.99	0.02	26,26,26,26	0
2	ZN	E	400	1/1	0.99	0.02	23,23,23,23	0
2	ZN	A	400	1/1	0.99	0.02	20,20,20,20	0
2	ZN	H	400	1/1	0.99	0.02	24,24,24,24	0
2	ZN	C	400	1/1	0.99	0.01	21,21,21,21	0
2	ZN	D	400	1/1	1.00	0.02	22,22,22,22	0

6.5 Other polymers [i](#)

There are no such residues in this entry.