



wwPDB X-ray Structure Validation Summary Report ⓘ

Mar 5, 2026 – 01:51 PM UTC

PDB ID : 2NYA / pdb_00002nya
Title : Crystal structure of the periplasmic nitrate reductase (NAP) from Escherichia coli
Authors : Jepson, B.J.N.; Richardson, D.J.; Hemmings, A.M.
Deposited on : 2006-11-20
Resolution : 2.50 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4-5-2 with Phenix2.0
Mogul : 2022.3.0, CSD as543be (2022)
Xtriage (Phenix) : 2.0
EDS : 3.0
Buster-report : wwPDB partial adaption of 1.1.7 (2018)
Percentile statistics : 20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4 : 9.0.010 (Gargrove)
Density-Fitness : 1.0.12
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.50 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	5829 (2.50-2.50)
Clashscore	190562	6492 (2.50-2.50)
Ramachandran outliers	187476	6378 (2.50-2.50)
Sidechain outliers	187428	6380 (2.50-2.50)
RSRZ outliers	180081	5833 (2.50-2.50)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	792	 66% 28% 6%
1	F	792	 63% 31% 6%

2 Entry composition [i](#)

There are 5 unique types of molecules in this entry. The entry contains 13733 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Periplasmic nitrate reductase.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	791	Total	C	N	O	S	0	0	0
			6301	4012	1100	1157	32			
1	F	791	Total	C	N	O	S	0	0	0
			6301	4012	1100	1157	32			

- Molecule 2 is IRON/SULFUR CLUSTER (CCD ID: SF4) (formula: Fe₄S₄).




Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
2	A	1	Total	Fe	S	0	0
			8	4	4		
2	F	1	Total	Fe	S	0	0
			8	4	4		

- Molecule 3 is MOLYBDENUM(VI) ION (CCD ID: 6MO) (formula: Mo).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	A	1	Total	Mo	0	0
			1	1		
3	F	1	Total	Mo	0	0
			1	1		

- Molecule 4 is 2-AMINO-5,6-DIMERCAPTO-7-METHYL-3,7,8A,9-TETRAHYDRO-8-OXA-1,3,9,10-TETRAAZA-ANTHRACEN-4-ONE GUANOSINE DINUCLEOTIDE (CCD ID: MGD) (formula: $C_{20}H_{26}N_{10}O_{13}P_2S_2$).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	
4	A	1	Total 47	C 20	N 10	O 13	P 2	S 2	0	0
4	A	1	Total 47	C 20	N 10	O 13	P 2	S 2	0	0
4	F	1	Total 47	C 20	N 10	O 13	P 2	S 2	0	0
4	F	1	Total 47	C 20	N 10	O 13	P 2	S 2	0	0


- Molecule 5 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
5	A	489	Total	O	0	0
			489	489		
5	F	436	Total	O	0	0
			436	436		


3 Residue-property plots

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

• Molecule 1: Periplasmic nitrate reductase



• Molecule 1: Periplasmic nitrate reductase



K197	M308	K408	A498	A598	K718
M198	T309	K409	Y499	W599	A719
Q199	G310	G311	G500	F600	R720
	V311	I412	N501	G601	D721
T202	K312	P413	A502	L722	L722
V203	D314	T416	E503	H606	R723
	Q315		R504		
L206	L316		R505	Y614	
S207	E317	I421	T506	G619	D726
T208	Q318	G422	F507	L620	V730
Y209	L319	L423	W509	R621	V731
Q210		H424	R510		S732
H211	I331	A427	Q511	K628	R733
E212	S332		Q512	Q631	R734
F213	M336	L432	A515	Y634	G735
E215	G337	D434	E518		
L216	F338	L437	A519	D639	T740
A217	Q340	T442	K520	V642	V741
D218	H341	M443	Q525	K643	N747
	R343	C444	Q528	A644	R748
I222	W346	M446	R531	G652	P749
F223	N353	M447	R532	V659	P750
T224	L354	Q449	F533	I660	
P225	H355	P452	V538	A668	L753
Q226	K360	E456	W539	D673	V754
S227	C366	E457	L543	B674	Y755
V230	G373	N457	L544	R684	N756
I231	Q374	I470	R551	V685	F759
L232	P375	V471	T554	L686	
N233	S376	S472	L555	B687	L764
Y234	R382	D473	Y556	H690	L768
I235	V384	P474	A561	T691	T769
	R390	S479	E572	R696	L770
Q241	L391	A483	L573	R697	P775
	M395	D484	A574	E700	E779
I245	V396	I486	Q577	L701	T780
D248	V397	L487	L578	H702	K783
S251	T398	F488	N579	R703	K784
K252	N399	T489	D580	A704	
H253	K401	A490	E581	P706	V787
L256	H402	M491	S582	V709	K791
I263	R403	W492	F587	L710	VAL
R268	L405	E494	Y588	F711	
G281	I406	E496	L589	I712	
T301	C406	G497	Y597	H713	
L302	E407				
E303					
R304					
T305					

4 Data and refinement statistics

Property	Value	Source
Space group	P 1 21 1	Depositor
Cell constants a, b, c, α , β , γ	69.45Å 94.60Å 131.21Å 90.00° 96.34° 90.00°	Depositor
Resolution (Å)	50.00 – 2.50 50.00 – 2.50	Depositor EDS
% Data completeness (in resolution range)	97.1 (50.00-2.50) 97.1 (50.00-2.50)	Depositor EDS
R_{merge}	0.11	Depositor
R_{sym}	0.11	Depositor
$\langle I/\sigma(I) \rangle$ ¹	2.50 (at 2.51Å)	Xtriage
Refinement program	REFMAC 5.2.0005	Depositor
R, R_{free}	0.181 , 0.243 0.181 , 0.241	Depositor DCC
R_{free} test set	2860 reflections (4.89%)	wwPDB-VP
Wilson B-factor (Å ²)	27.6	Xtriage
Anisotropy	0.729	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.34 , 43.5	EDS
L-test for twinning ²	$\langle L \rangle = 0.49$, $\langle L^2 \rangle = 0.32$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.95	EDS
Total number of atoms	13733	wwPDB-VP
Average B, all atoms (Å ²)	28.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 11.11% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: SF4, 6MO, MGD

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.88	2/6470 (0.0%)	1.07	13/8770 (0.1%)
1	F	0.83	0/6470	1.03	9/8770 (0.1%)
All	All	0.86	2/12940 (0.0%)	1.05	22/17540 (0.1%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	A	0	1
1	F	0	2
All	All	0	3

All (2) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	224	THR	CA-CB	5.12	1.63	1.53
1	A	80	THR	CA-CB	5.05	1.62	1.54

The worst 5 of 22 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	57	GLY	O-C-N	-7.37	113.12	122.70
1	F	263	ILE	N-CA-C	7.33	118.44	111.91
1	A	103	GLY	CA-C-N	7.23	127.85	119.47
1	A	103	GLY	C-N-CA	7.23	127.85	119.47
1	F	126	LEU	N-CA-C	-6.48	104.13	111.07

There are no chirality outliers.

All (3) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	A	138	PRO	Peptide
1	F	313	LYS	Peptide
1	F	57	GLY	Peptide

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	6301	0	6155	227	0
1	F	6301	0	6155	258	0
2	A	8	0	0	0	0
2	F	8	0	0	0	0
3	A	1	0	0	0	0
3	F	1	0	0	0	0
4	A	94	0	44	14	0
4	F	94	0	44	17	0
5	A	489	0	0	15	0
5	F	436	0	0	29	1
All	All	13733	0	12398	481	1

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 19.

The worst 5 of 481 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:F:783:LYS:HG3	5:F:7033:HOH:O	1.43	1.18
1:F:109:MET:CE	1:F:123:ALA:HB1	1.95	0.96
1:F:684:ARG:NH2	4:F:6001:MGD:H15	1.63	0.95
1:F:193:ASN:HB3	5:F:7024:HOH:O	1.68	0.93
1:F:668:ALA:HA	5:F:7032:HOH:O	1.67	0.93

All (1) symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
5:F:7379:HOH:O	5:F:7409:HOH:O[2_655]	1.98	0.22

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	789/792 (100%)	736 (93%)	47 (6%)	6 (1%)	16	31
1	F	789/792 (100%)	725 (92%)	55 (7%)	9 (1%)	11	22
All	All	1578/1584 (100%)	1461 (93%)	102 (6%)	15 (1%)	12	24

5 of 15 Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	139	ASN
1	F	70	ASN
1	F	215	GLU
1	F	314	ASP
1	F	241	GLN

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	661/663 (100%)	595 (90%)	66 (10%)	7	15
1	F	661/663 (100%)	604 (91%)	57 (9%)	10	21
All	All	1322/1326 (100%)	1199 (91%)	123 (9%)	8	18

5 of 123 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	700	GLU
1	F	518	GLU
1	F	51	LEU
1	F	512	GLN
1	F	700	GLU

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 57 such sidechains are listed below:

Mol	Chain	Res	Type
1	F	135	ASN
1	F	606	HIS
1	F	233	ASN
1	F	604	HIS
1	F	501	ASN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

Of 8 ligands modelled in this entry, 2 are monoatomic - leaving 6 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
4	MGD	A	4001	3	47,52,52	1.46	7 (14%)	58,81,81	1.97	13 (22%)
4	MGD	F	7001	3	47,52,52	1.47	7 (14%)	58,81,81	2.22	16 (27%)
2	SF4	F	5001	1	0,12,12	-	-	-	-	-
2	SF4	A	2001	1	0,12,12	-	-	-	-	-
4	MGD	A	3001	3	47,52,52	1.33	4 (8%)	58,81,81	2.16	18 (31%)
4	MGD	F	6001	3	47,52,52	1.45	8 (17%)	58,81,81	2.26	21 (36%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
4	MGD	A	4001	3	-	7/22/66/66	0/6/6/6
4	MGD	F	7001	3	-	6/22/66/66	0/6/6/6
2	SF4	F	5001	1	-	-	0/6/5/5
2	SF4	A	2001	1	-	-	0/6/5/5
4	MGD	A	3001	3	-	6/22/66/66	0/6/6/6
4	MGD	F	6001	3	-	4/22/66/66	0/6/6/6

The worst 5 of 26 bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
4	A	4001	MGD	C16-C21	5.58	1.48	1.38
4	F	7001	MGD	C16-C21	5.58	1.48	1.38
4	F	6001	MGD	C16-C21	5.01	1.47	1.38
4	A	3001	MGD	C16-C21	4.72	1.46	1.38
4	A	4001	MGD	C17-N18	-3.40	1.32	1.38

The worst 5 of 68 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
4	F	7001	MGD	C23-C14-N15	7.73	115.46	107.87
4	A	4001	MGD	C5-C4-N3	-5.92	118.97	128.39
4	F	7001	MGD	C5-C4-N3	-5.68	119.35	128.39
4	F	7001	MGD	C2-N3-C4	5.59	121.93	112.30
4	A	3001	MGD	C19-N20-C21	5.42	122.92	113.36

There are no chirality outliers.

5 of 23 torsion outliers are listed below:


Mol	Chain	Res	Type	Atoms
4	A	3001	MGD	PA-O3B-PB-O5'
4	A	3001	MGD	C5'-O5'-PB-O2B
4	A	3001	MGD	C5'-O5'-PB-O3B
4	A	4001	MGD	C5'-O5'-PB-O1B
4	A	4001	MGD	C5'-O5'-PB-O2B


There are no ring outliers.


4 monomers are involved in 31 short contacts:


Mol	Chain	Res	Type	Clashes	Symm-Clashes
4	A	4001	MGD	7	0
4	F	7001	MGD	8	0
4	A	3001	MGD	7	0
4	F	6001	MGD	9	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.









5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	791/792 (99%)	-0.47	0 100 100	13, 24, 37, 46	0
1	F	791/792 (99%)	-0.30	0 100 100	17, 30, 45, 55	0
All	All	1582/1584 (99%)	-0.38	0 100 100	13, 27, 42, 55	0

There are no RSRZ outliers to report.

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled ‘Q < 0.9’ lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
4	MGD	F	7001	47/47	0.97	0.06	18,22,27,31	0
4	MGD	A	3001	47/47	0.98	0.05	11,16,17,20	0
4	MGD	A	4001	47/47	0.98	0.06	13,21,26,30	0
4	MGD	F	6001	47/47	0.98	0.05	15,19,21,22	0
2	SF4	F	5001	8/8	0.98	0.03	25,26,29,30	0
2	SF4	A	2001	8/8	0.99	0.03	24,26,28,28	0
3	6MO	A	2002	1/1	0.99	0.03	20,20,20,20	0

Continued on next page...




Continued from previous page...

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors($\text{\AA}^2$)	Q<0.9
3	6MO	F	5002	1/1	1.00	0.02	24,24,24,24	0

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.



Electron density around MGD F 7001:

2mF_o-DF_c (at 0.7 rmsd) in gray
mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)





Electron density around MGD A 3001:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)




**Electron density around MGD A 4001:**

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)



Electron density around MGD F 6001:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)



6.5 Other polymers [i](#)

There are no such residues in this entry.