wwPDB X-ray Structure Validation Summary Report (i)

Aug 5, 2026 – 03:37 PM EDT

PDB ID	:	3O0O / pdb_00003o0o
Title	:	Thermotoga maritima Ribonucleotide Reductase, NrdJ, in complex with dTTP, GDP and Adenosylcobalamin
Authors	:	Larsson, K.-M.; Logan, D.T.; Nordlund, P.
Deposited on	:	2010-07-19
Resolution	:	1.90 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at *validation@mail.wwpdb.org*
A user guide is available at
https://www.wwpdb.org/validation/2017/XrayValidationReportHelp
with specific help available everywhere you see the (i) symbol.

The types of validation reports are described at
http://www.wwpdb.org/validation/2017/FAQs#types.

The following versions of software and data (see references (i)) were used in the production of this report:

MolProbity	:	4-5-2 with Phenix2.0
Mogul	:	2022.3.0, CSD as543be (2022)
Xtriage (Phenix)	:	2.0
EDS	:	3.0
Buster-report	:	wwPDB partial adaption of 1.1.7 (2018)
Percentile statistics	:	20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4	:	9.0.015 (Gargrove)
Density-Fitness	:	1.0.12
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.50

1 Overall quality at a glance (i)

The following experimental techniques were used to determine the structure:
X-RAY DIFFRACTION

The reported resolution of this entry is 1.90 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	7789 (1.90-1.90)
Clashscore	190562	8410 (1.90-1.90)
Ramachandran outliers	187476	8333 (1.90-1.90)
Sidechain outliers	187428	8333 (1.90-1.90)
RSRZ outliers	180081	7790 (1.90-1.90)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for >=3, 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions <=5% The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	644	6% 71% 22% • 5%
1	B	644	2% 82% 13% • •

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
5	B12	A	1004	X	-	-	-
5	B12	B	1004	X	-	X	-

2 Entry composition (i)

There are 7 unique types of molecules in this entry. The entry contains 10787 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Ribonucleoside-diphosphate reductase.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	612	4916	3152	840	904	20	0	1	0
1	B	618	5018	3216	858	921	23	0	8	0

There are 2 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	205	SER	TYR	SEE REMARK 999	UNP O33839
B	205	SER	TYR	SEE REMARK 999	UNP O33839

- Molecule 2 is THYMIDINE-5'-TRIPHOSPHATE (CCD ID: TTP) (formula: $C_{10}H_{17}N_2O_{14}P_3$).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
			Total	C	N	O	P		
2	A	1	29	10	2	14	3	0	0

Continued on next page...

Continued from previous page...

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
			Total	C	N	O	P		
2	B	1	29	10	2	14	3	0	0

- Molecule 3 is MAGNESIUM ION (CCD ID: MG) (formula: Mg).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
			Total	Mg		
3	A	1	1	1	0	0
			Total	Mg		
3	B	1	1	1	0	0

- Molecule 4 is GUANOSINE-5'-DIPHOSPHATE (CCD ID: GDP) (formula: $C_{10}H_{15}N_5O_{11}P_2$).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
			Total	C	N	O	P		
4	A	1	56	20	10	22	4	0	1
			Total	C	N	O	P		
4	B	1	56	20	10	22	4	0	1

- Molecule 5 is COBALAMIN (CCD ID: B12) (formula: $C_{62}H_{89}CoN_{13}O_{14}P$).



Mol	Chain	Residues	Atoms						ZeroOcc	AltConf
			Total	C	Co	N	O	P		
5	A	1	91	62	1	13	14	1	0	0
5	B	1	91	62	1	13	14	1	0	0

- Molecule 6 is 5'-DEOXYADENOSINE (CCD ID: 5AD) (formula: $C_{10}H_{13}N_5O_3$).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
			Total	C	N	O		
6	A	1	18	10	5	3	0	0
6	B	1	18	10	5	3	0	0

- Molecule 7 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
			Total	O		
7	A	151	151	151	0	0
			Total	O		
7	B	312	312	312	0	0

3 Residue-property plots (i)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density (RSRZ > 2). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

• Molecule 1: Ribonucleoside-diphosphate reductase

Chain A: 6% 71% 22% • 5%



• Molecule 1: Ribonucleoside-diphosphate reductase

Chain B: 2% 82% 13% • •



THR
GLN
VAL
LEU
THR
LYS
ALA
LEU
LYS
THR

4 Data and refinement statistics (i)

Property	Value	Source
Space group	C 1 2 1	Depositor
Cell constants a, b, c, α, β, γ	120.02Å 124.23Å 107.17Å 90.00° 102.57° 90.00°	Depositor
Resolution (Å)	46.94 – 1.90 46.94 – 1.90	Depositor EDS
% Data completeness (in resolution range)	99.5 (46.94-1.90) 99.4 (46.94-1.90)	Depositor EDS
R_{merge}	0.06	Depositor
R_{sym}	(Not available)	Depositor
$< I/\sigma(I) >$ [1]	1.41 (at 1.90Å)	Xtriage
Refinement program	PHENIX (phenix.refine: 1.6.2_432), REFMAC	Depositor
R, R_{free}	0.202 , 0.238 0.201 , 0.237	Depositor DCC
R_{free} test set	5994 reflections (4.98%)	wwPDB-VP
Wilson B-factor (Å^2)	31.5	Xtriage
Anisotropy	0.404	Xtriage
Bulk solvent k_{sol}(e/Å^3), B_{sol}(Å^2)	0.35 , 65.2	EDS
L-test for twinning[2]	$< \|L\| > = 0.48$, $< L^2 > = 0.31$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o,F_c correlation	0.97	EDS
Total number of atoms	10787	wwPDB-VP
Average B, all atoms (Å^2)	59.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 4.88% of the height of the origin peak. No significant pseudotranslation is detected.*

[1]Intensities estimated from amplitudes.

[2]Theoretical values of $< |L| >$, $< L^2 >$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality (i)

5.1 Standard geometry (i)

Bond lengths and bond angles in the following residue types are not validated in this section: B12, 5AD, MG, TTP, GDP

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\#\|Z\| >5$	RMSZ	$\#\|Z\| >5$
1	A	0.42	0/5015	0.78	3/6766 (0.0%)
1	B	0.52	0/5129	0.78	1/6918 (0.0%)
All	All	0.47	0/10144	0.78	4/13684 (0.0%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand.A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	B	0	1

There are no bond length outliers.

All (4) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(^{o})	Ideal(^{o})
1	B	134	CYS	N-CA-C	5.13	116.00	107.73
1	A	171	ARG	CA-C-N	5.06	124.99	119.78
1	A	171	ARG	C-N-CA	5.06	124.99	119.78
1	A	32	LEU	N-CA-C	-5.01	107.00	113.01

There are no chirality outliers.

All (1) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	B	241	PHE	Peptide

5.2 Too-close contacts (i)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	4916	0	4979	121	0
1	B	5018	0	5091	70	0
2	A	29	0	13	1	0
2	B	29	0	13	0	0
3	A	1	0	0	0	0
3	B	1	0	0	0	0
4	A	56	0	24	3	0
4	B	56	0	24	6	0
5	A	91	0	86	14	0
5	B	91	0	86	21	0
6	A	18	0	13	1	0
6	B	18	0	13	3	0
7	A	151	0	0	4	0
7	B	312	0	0	4	0
All	All	10787	0	10342	224	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 11.

The worst 5 of 224 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:617:ALA:CB	1:A:625:ILE:HD11	1.86	1.05
1:A:583:MET:HE3	1:A:583:MET:HA	1.40	1.01
5:A:1004:B12:H601	5:A:1004:B12:H262	1.42	0.99
5:B:1004:B12:H351	5:B:1004:B12:H362	1.51	0.93
1:B:428:HIS:HD2	1:B:480:ARG:HH22	1.20	0.90

There are no symmetry-related clashes.

5.3 Torsion angles (i)

5.3.1 Protein backbone (i)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	603/644 (94%)	578 (96%)	23 (4%)	2 (0%)	36	29
1	B	620/644 (96%)	600 (97%)	20 (3%)	0	100	100
All	All	1223/1288 (95%)	1178 (96%)	43 (4%)	2 (0%)	43	36

All (2) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	546	GLU
1	A	2	LYS

5.3.2 Protein sidechains (i)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	539/566 (95%)	515 (96%)	24 (4%)	24	17
1	B	551/566 (97%)	541 (98%)	10 (2%)	51	50
All	All	1090/1132 (96%)	1056 (97%)	34 (3%)	35	29

5 of 34 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	B	418	ASN
1	B	445	GLU
1	B	564	LYS
1	A	333	CYS

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	A	325	ILE

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 20 such sidechains are listed below:

Mol	Chain	Res	Type
1	B	96	ASN
1	B	359	GLN
1	B	507	ASN
1	B	428	HIS
1	A	418	ASN

5.3.3 RNA (i)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains (i)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates (i)

There are no oligosaccharides in this entry.

5.6 Ligand geometry (i)

Of 12 ligands modelled in this entry, 2 are monoatomic - leaving 10 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\#\|Z\| > 2$	Counts	RMSZ	$\#\|Z\| > 2$
6	5AD	A	1005	-	20,20,20	1.78	7 (35%)	28,30,30	2.69	11 (39%)
6	5AD	B	1005	-	20,20,20	1.77	7 (35%)	28,30,30	2.86	11 (39%)

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\#\|Z\| > 2$	Counts	RMSZ	$\#\|Z\| > 2$
2	TTP	A	1001	3	29,30,30	1.41	6 (20%)	43,47,47	2.02	11 (25%)
4	GDP	B	1003[A]	-	29,30,30	1.22	4 (13%)	45,47,47	1.98	9 (20%)
2	TTP	B	1001	3	29,30,30	1.41	7 (24%)	43,47,47	1.85	10 (23%)
4	GDP	B	1003[B]	-	29,30,30	1.18	4 (13%)	45,47,47	1.66	8 (17%)
4	GDP	A	1003[A]	-	29,30,30	1.19	3 (10%)	45,47,47	1.79	8 (17%)
5	B12	A	1004	-	94,101,101	1.02	5 (5%)	149,166,166	1.39	9 (6%)
5	B12	B	1004	-	94,101,101	1.07	5 (5%)	149,166,166	1.46	15 (10%)
4	GDP	A	1003[B]	-	29,30,30	1.15	3 (10%)	45,47,47	1.74	8 (17%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
6	5AD	A	1005	-	-	0/4/20/20	0/3/3/3
6	5AD	B	1005	-	-	0/4/20/20	0/3/3/3
2	TTP	A	1001	3	-	2/22/34/34	0/2/2/2
2	TTP	B	1001	3	-	6/22/34/34	0/2/2/2
4	GDP	B	1003[A]	-	-	7/16/32/32	0/3/3/3
4	GDP	B	1003[B]	-	-	7/16/32/32	0/3/3/3
4	GDP	A	1003[A]	-	-	7/16/32/32	0/3/3/3
5	B12	A	1004	-	1/1/36/38	12/56/223/223	0/3/11/11
5	B12	B	1004	-	1/1/36/38	4/56/223/223	0/3/11/11
4	GDP	A	1003[B]	-	-	3/16/32/32	0/3/3/3

The worst 5 of 51 bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
5	B	1004	B12	C5M-C5B	-3.82	1.43	1.51
5	B	1004	B12	C14-N23	3.79	1.40	1.35
5	B	1004	B12	C53-C15	3.78	1.58	1.50
5	A	1004	B12	C5M-C5B	-3.74	1.44	1.51
5	A	1004	B12	C53-C15	3.55	1.58	1.50

The worst 5 of 100 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(^{o})	Ideal(^{o})
6	B	1005	5AD	C5'-C4'-C3'	-8.87	106.39	115.70
5	A	1004	B12	C1-C19-C18	8.44	135.58	121.90
5	B	1004	B12	C1-C19-N24	8.11	115.27	106.25
5	B	1004	B12	C1-C19-C18	7.73	134.44	121.90
5	A	1004	B12	C1-C19-N24	7.55	114.65	106.25

All (2) chirality outliers are listed below:

Mol	Chain	Res	Type	Atom
5	A	1004	B12	C19
5	B	1004	B12	C19

5 of 48 torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
2	B	1001	TTP	C5'-O5'-PA-O1A
4	A	1003[A]	GDP	C5'-O5'-PA-O3A
4	A	1003[A]	GDP	C5'-O5'-PA-O1A
4	A	1003[A]	GDP	C5'-O5'-PA-O2A
4	A	1003[A]	GDP	O4'-C4'-C5'-O5'

There are no ring outliers.

8 monomers are involved in 43 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
6	A	1005	5AD	1	0
6	B	1005	5AD	3	0
2	A	1001	TTP	1	0
4	B	1003[A]	GDP	1	0
4	B	1003[B]	GDP	5	0
5	A	1004	B12	14	0
5	B	1004	B12	21	0
4	A	1003[B]	GDP	3	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less then 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and

any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.

Ligand 5AD A 1005



Bond lengths



Bond angles



Torsions



Rings

Ligand 5AD B 1005



Bond lengths



Bond angles



Torsions



Rings

Ligand TTP A 1001



Bond lengths



Bond angles



Torsions



Rings

Ligand GDP B 1003 (A)



Bond lengths



Bond angles



Torsions



Rings

Ligand TTP B 1001



Bond lengths



Bond angles



Torsions



Rings

Ligand GDP B 1003 (B)



Bond lengths



Bond angles



Torsions



Rings

Ligand GDP A 1003 (A)	
 Bond lengths	 Bond angles
 Torsions	 Rings

Ligand B12 A 1004



Bond lengths



Bond angles



Torsions



Rings

Ligand B12 B 1004



Bond lengths



Bond angles



Torsions



Rings



5.7 Other polymers (i)

There are no such residues in this entry.

5.8 Polymer linkage issues (i)

There are no chain breaks in this entry.

6 Fit of model and data (i)

6.1 Protein, DNA and RNA chains (i)

In the following table, the column labelled '#RSRZ> 2' contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled 'Q< 0.9' lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2			OWAB(Å^2)	Q<0.9
1	A	612/644 (95%)	0.61	39 (6%)	25	27	34, 68, 122, 153	1 (0%)
1	B	618/644 (95%)	-0.07	13 (2%)	63	67	14, 40, 82, 110	8 (1%)
All	All	1230/1288 (95%)	0.27	52 (4%)	40	43	14, 53, 113, 153	9 (0%)

The worst 5 of 52 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	237	VAL	4.7
1	A	204	GLY	4.7
1	B	209	GLY	4.4
1	B	204	GLY	4.1
1	B	237	VAL	4.1

6.2 Non-standard residues in protein, DNA, RNA chains (i)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates (i)

There are no oligosaccharides in this entry.

6.4 Ligands (i)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å^2)	Q<0.9
3	MG	B	1002	1/1	0.81	0.17	57,57,57,57	0
6	5AD	A	1005	18/18	0.83	0.18	54,69,71,72	16
5	B12	A	1004	91/91	0.89	0.15	33,71,88,99	81
4	GDP	A	1003[A]	28/28	0.90	0.16	36,64,72,73	28
4	GDP	A	1003[B]	28/28	0.90	0.16	31,51,65,68	28
4	GDP	B	1003[A]	28/28	0.91	0.13	13,35,46,51	28
4	GDP	B	1003[B]	28/28	0.91	0.13	15,28,49,54	28
6	5AD	B	1005	18/18	0.92	0.11	41,56,67,69	12
2	TTP	B	1001	29/29	0.94	0.08	31,40,60,63	0
5	B12	B	1004	91/91	0.94	0.11	27,52,72,90	65
2	TTP	A	1001	29/29	0.95	0.07	20,38,44,47	0
3	MG	A	1002	1/1	0.96	0.08	36,36,36,36	0

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.

Electron density around 5AD A 1005:

2mF$_o$-DF$_c$ (at 0.7 rmsd) in gray
mF$_o$-DF$_c$ (at 3 rmsd) in purple (negative) and green (positive)







Electron density around B12 A 1004:

$2mF_o$-DF_c (at 0.7 rmsd) in gray
mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)







Electron density around GDP A 1003 (A):

$2mF_o$-DF_c (at 0.7 rmsd) in gray
mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)







Electron density around GDP A 1003 (B):

$2mF_o$-DF_c (at 0.7 rmsd) in gray
mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)



Electron density around GDP B 1003 (A):

$2mF_o$-DF_c (at 0.7 rmsd) in gray
mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)



Electron density around GDP B 1003 (B):

$2mF_o$-DF_c (at 0.7 rmsd) in gray
mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)







Electron density around 5AD B 1005:

$2mF_o$-DF_c (at 0.7 rmsd) in gray
mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)







Electron density around TTP B 1001:

$2mF_o$-DF_c (at 0.7 rmsd) in gray
mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)



Electron density around B12 B 1004:

$2mF_o$-DF_c (at 0.7 rmsd) in gray
mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)



Electron density around TTP A 1001:

$2mF_o$-DF_c (at 0.7 rmsd) in gray
mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)



6.5 Other polymers (i)

There are no such residues in this entry.