



# Full wwPDB X-ray Structure Validation Report ⓘ

Mar 5, 2026 – 08:53 AM UTC

PDB ID : 1O7M / pdb\_00001o7m  
Title : NAPHTHALENE 1,2-DIOXYGENASE, BINARY COMPLEX WITH DIOXYGEN  
Authors : Karlsson, A.; Parales, J.V.; Parales, R.E.; Gibson, D.T.; Eklund, H.; Ramaswamy, S.  
Deposited on : 2002-11-11  
Resolution : 1.75 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4-5-2 with Phenix2.0
Mogul	:	2022.3.0, CSD as543be (2022)
Xtriage (Phenix)	:	2.0
EDS	:	3.0
Percentile statistics	:	20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4	:	9.0.010 (Gargrove)
Density-Fitness	:	1.0.12
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.49


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

*X-RAY DIFFRACTION*

The reported resolution of this entry is 1.75 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	180053	3183 (1.76-1.76)
Clashscore	190562	3299 (1.76-1.76)
Ramachandran outliers	187476	3274 (1.76-1.76)
Sidechain outliers	187428	3274 (1.76-1.76)
RSRZ outliers	180081	3183 (1.76-1.76)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	449	<div> <div>7%</div> <div>89%</div> <div>9%</div> <div>.</div> </div>
2	B	194	<div> <div>15%</div> <div>88%</div> <div>10%</div> <div>...</div> </div>

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
3	EDO	B	1696	-	-	X	-

## 2 Entry composition [i](#)

There are 8 unique types of molecules in this entry. The entry contains 5694 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.


- Molecule 1 is a protein called NAPHTHALENE 1,2-DIOXYGENASE ALPHA SUBUNIT.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	448	Total	C	N	O	S	0	2	1
			3487	2205	599	666	17			

- Molecule 2 is a protein called NAPHTHALENE 1,2-DIOXYGENASE BETA SUBUNIT.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
2	B	193	Total	C	N	O	S	0	0	0
			1608	1007	302	293	6			

- Molecule 3 is 1,2-ETHANEDIOL (CCD ID: EDO) (formula: C<sub>2</sub>H<sub>6</sub>O<sub>2</sub>).




Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
3	A	1	Total	C	O	0	0
			4	2	2		
3	A	1	Total	C	O	0	0
			4	2	2		

*Continued on next page...*

*Continued from previous page...*


Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
3	A	1	Total	C	O	0	0
			4	2	2		
3	B	1	Total	C	O	0	0
			4	2	2		
3	B	1	Total	C	O	0	0
			4	2	2		
3	B	1	Total	C	O	0	0
			4	2	2		

- Molecule 4 is OXYGEN MOLECULE (CCD ID: OXY) (formula: O<sub>2</sub>).




Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	A	1	Total	O	0	0
			2	2		

- Molecule 5 is SULFATE ION (CCD ID: SO4) (formula: O<sub>4</sub>S).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
5	A	1	Total	O	S	0	0
			5	4	1		
5	B	1	Total	O	S	0	0
			5	4	1		
5	B	1	Total	O	S	0	0
			5	4	1		

- Molecule 6 is FE2/S2 (INORGANIC) CLUSTER (CCD ID: FES) (formula:  $\text{Fe}_2\text{S}_2$ ).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
6	A	1	Total	Fe	S	0	0
			4	2	2		

- Molecule 7 is FE (III) ION (CCD ID: FE) (formula: Fe).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
7	A	1	Total 1	Fe 1	0	0


- Molecule 8 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
8	A	401	Total 401	O 401	0	0
8	B	152	Total 152	O 152	0	0


### 3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

#### • Molecule 1: NAPHTHALENE 1,2-DIOXYGENASE ALPHA SUBUNIT



#### • Molecule 2: NAPHTHALENE 1,2-DIOXYGENASE BETA SUBUNIT





## 4 Data and refinement statistics

Property	Value	Source
Space group	H 3 2	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	140.28Å 140.28Å 208.75Å 90.00° 90.00° 120.00°	Depositor
Resolution (Å)	30.00 – 1.75 30.00 – 1.75	Depositor EDS
% Data completeness (in resolution range)	98.9 (30.00-1.75) 98.9 (30.00-1.75)	Depositor EDS
$R_{merge}$	0.06	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	1.53 (at 1.57Å)	Xtriage
Refinement program	REFMAC 5.0.36	Depositor
R, $R_{free}$	0.178 , 0.187 0.190 , 0.194	Depositor DCC
$R_{free}$ test set	927 reflections (0.99%)	wwPDB-VP
Wilson B-factor (Å <sup>2</sup> )	16.5	Xtriage
Anisotropy	0.124	Xtriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.38 , 41.7	EDS
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.51$ , $\langle L^2 \rangle = 0.34$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
$F_o, F_c$ correlation	0.95	EDS
Total number of atoms	5694	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	16.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 4.03% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

## 5 Model quality [i](#)

### 5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: FES, SO4, EDO, OXY, FE

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z  > 5$	RMSZ	$\# Z  > 5$
1	A	0.80	1/3589 (0.0%)	1.03	6/4862 (0.1%)
2	B	0.78	0/1638	1.03	3/2209 (0.1%)
All	All	0.79	1/5227 (0.0%)	1.03	9/7071 (0.1%)

All (1) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	447	THR	C-N	-6.17	1.24	1.33

All (9) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	B	548	GLN	OE1-CD-NE2	-10.05	112.55	122.60
2	B	548	GLN	CG-CD-NE2	6.67	126.40	116.40
1	A	219	SER	CA-C-N	6.64	128.49	120.13
1	A	219	SER	C-N-CA	6.64	128.49	120.13
2	B	634	LEU	N-CA-C	6.32	119.02	108.73
1	A	220	GLY	N-CA-C	6.11	121.40	113.27
1	A	234	PRO	CB-CA-C	5.86	118.07	110.92
1	A	35	PHE	N-CA-C	5.28	117.12	111.36
1	A	222	SER	N-CA-C	5.08	117.61	110.14

There are no chirality outliers.

There are no planarity outliers.

### 5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen

atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	3487	0	3323	38	0
2	B	1608	0	1583	15	0
3	A	12	0	18	0	0
3	B	12	0	18	4	0
4	A	2	0	0	0	0
5	A	5	0	0	1	0
5	B	10	0	0	0	0
6	A	4	0	0	0	0
7	A	1	0	0	0	0
8	A	401	0	0	4	0
8	B	152	0	0	0	0
All	All	5694	0	4942	53	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 5.

All (53) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:240:LEU:HD13	1:A:252:VAL:HB	1.40	1.04
1:A:131:LYS:HD3	5:A:1452:SO4:O2	1.68	0.93
1:A:342:ARG:HH11	1:A:342:ARG:HG2	1.34	0.89
1:A:227:LEU:CD2	1:A:234:PRO:HD3	2.14	0.77
1:A:342:ARG:HG2	1:A:342:ARG:NH1	1.99	0.77
2:B:581:LEU:HD13	3:B:1696:EDO:H21	1.66	0.75
1:A:144:PHE:HD1	1:A:160:MET:HE1	1.52	0.73
1:A:144:PHE:CD1	1:A:160:MET:HE1	2.25	0.71
1:A:362:ASP:HB3	1:A:366[B]:MET:HE2	1.75	0.66
1:A:47:LEU:HD22	1:A:182:LEU:HD23	1.80	0.62
1:A:240:LEU:CD1	1:A:252:VAL:HB	2.24	0.62
2:B:525:CYS:SG	2:B:525:CYS:O	2.59	0.61
2:B:581:LEU:HD13	3:B:1696:EDO:C2	2.31	0.60
1:A:227:LEU:HD21	1:A:234:PRO:HD3	1.84	0.59
1:A:235:PRO:HB2	1:A:236:GLU:OE1	2.03	0.59
1:A:227:LEU:HD22	1:A:234:PRO:HD3	1.84	0.59
1:A:220:GLY:HA3	8:A:2228:HOH:O	2.03	0.59
2:B:581:LEU:CD1	3:B:1696:EDO:H21	2.33	0.58
2:B:634:LEU:HD11	2:B:662:LYS:HB3	1.87	0.56
2:B:631:ASP:OD1	2:B:633:GLU:HB2	2.07	0.55

*Continued on next page...*

*Continued from previous page...*

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:240:LEU:HD13	1:A:252:VAL:CB	2.28	0.54
1:A:220:GLY:O	1:A:221:GLU:HB2	2.09	0.53
1:A:160:MET:HE3	8:A:2076:HOH:O	2.09	0.53
1:A:266:VAL:HB	1:A:267:PRO:HD3	1.91	0.52
1:A:3:TYR:CE2	1:A:26:LEU:HD13	2.45	0.52
2:B:665:ARG:HG2	2:B:671:ARG:HG2	1.92	0.50
1:A:231:ALA:HB2	8:A:2230:HOH:O	2.12	0.50
2:B:581:LEU:O	2:B:582:ASN:C	2.56	0.49
1:A:204:GLY:HA3	1:A:412:SER:HB2	1.96	0.48
1:A:160:MET:HA	1:A:160:MET:HE2	1.95	0.48
1:A:2:ASN:OD1	1:A:4:ASN:HB2	2.12	0.48
2:B:665:ARG:HA	2:B:671:ARG:HA	1.96	0.47
1:A:236:GLU:HB2	1:A:237:GLY:H	1.36	0.47
1:A:44:HIS:HB2	1:A:182:LEU:HD22	1.97	0.47
2:B:581:LEU:HD13	3:B:1696:EDO:C1	2.45	0.46
2:B:634:LEU:HD21	2:B:662:LYS:HD3	1.96	0.46
2:B:631:ASP:C	2:B:633:GLU:H	2.23	0.46
1:A:366[A]:MET:HE1	8:A:2217:HOH:O	2.16	0.46
1:A:198:PRO:HB2	1:A:316:TRP:CE2	2.52	0.45
1:A:234:PRO:HA	1:A:235:PRO:HD2	1.79	0.45
1:A:160:MET:HE2	1:A:160:MET:CA	2.47	0.45
1:A:261:HIS:NE2	1:A:344:LEU:HD11	2.32	0.45
2:B:530:LEU:HD21	2:B:635:LEU:HD13	2.01	0.43
1:A:311:GLY:HA2	1:A:331:ILE:HG13	2.00	0.43
1:A:236:GLU:H	1:A:236:GLU:HG3	1.46	0.43
2:B:502:MET:HE3	2:B:502:MET:HB2	1.86	0.43
2:B:631:ASP:C	2:B:633:GLU:N	2.75	0.42
1:A:19:LEU:C	1:A:19:LEU:HD12	2.45	0.42
1:A:78:LEU:HD22	1:A:136:LYS:HD3	2.02	0.42
1:A:276:LYS:HD2	1:A:292:TYR:CZ	2.55	0.41
1:A:261:HIS:NE2	1:A:344:LEU:CD1	2.84	0.41
1:A:396:ASP:HB3	1:A:399:TYR:O	2.21	0.40
1:A:223:ILE:HD12	1:A:223:ILE:HA	1.89	0.40

There are no symmetry-related clashes.

## 5.3 Torsion angles

### 5.3.1 Protein backbone

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	448/449 (100%)	431 (96%)	15 (3%)	2 (0%)	30	15
2	B	191/194 (98%)	184 (96%)	6 (3%)	1 (0%)	24	11
All	All	639/643 (99%)	615 (96%)	21 (3%)	3 (0%)	24	11

All (3) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	236	GLU
2	B	525	CYS
1	A	235	PRO

### 5.3.2 Protein sidechains

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	369/369 (100%)	360 (98%)	9 (2%)	43	23
2	B	172/173 (99%)	160 (93%)	12 (7%)	14	2
All	All	541/542 (100%)	520 (96%)	21 (4%)	28	9

All (21) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	47	LEU
1	A	78	LEU
1	A	121	LYS

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type
1	A	131	LYS
1	A	201	ASN
1	A	219	SER
1	A	236	GLU
1	A	240	LEU
1	A	253	LEU
2	B	503	ILE
2	B	506	GLN
2	B	508	ASP
2	B	521	ARG
2	B	524	ASN
2	B	525	CYS
2	B	532	GLN
2	B	554	LEU
2	B	571	LEU
2	B	581	LEU
2	B	593	GLN
2	B	690	LEU

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (7) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	4	ASN
1	A	28	GLN
1	A	115	GLN
1	A	201	ASN
2	B	506	GLN
2	B	514	HIS
2	B	526	HIS

### 5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

## 5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

## 5.6 Ligand geometry [i](#)

Of 12 ligands modelled in this entry, 1 is monoatomic - leaving 11 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z  > 2$	Counts	RMSZ	$\# Z  > 2$
4	OXY	A	1451	7	1,1,1	1.02	0	-		
5	SO4	B	1697	-	4,4,4	0.30	0	6,6,6	0.06	0
3	EDO	A	1450	-	3,3,3	0.36	0	2,2,2	0.39	0
3	EDO	B	1696	-	3,3,3	0.38	0	2,2,2	0.40	0
3	EDO	A	1448	-	3,3,3	0.44	0	2,2,2	0.24	0
6	FES	A	1453	1	0,4,4	-	-	-		
5	SO4	B	1698	-	4,4,4	0.19	0	6,6,6	0.23	0
3	EDO	B	1695	-	3,3,3	0.30	0	2,2,2	0.44	0
3	EDO	B	1699	-	3,3,3	0.49	0	2,2,2	0.42	0
3	EDO	A	1449	-	3,3,3	0.45	0	2,2,2	0.60	0
5	SO4	A	1452	-	4,4,4	0.21	0	6,6,6	0.36	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	EDO	A	1450	-	-	0/1/1/1	-
3	EDO	B	1696	-	-	1/1/1/1	-
3	EDO	A	1448	-	-	0/1/1/1	-
6	FES	A	1453	1	-	-	0/1/1/1
3	EDO	B	1695	-	-	0/1/1/1	-
3	EDO	B	1699	-	-	0/1/1/1	-
3	EDO	A	1449	-	-	0/1/1/1	-

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

All (1) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
3	B	1696	EDO	O1-C1-C2-O2

There are no ring outliers.

2 monomers are involved in 5 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
3	B	1696	EDO	4	0
5	A	1452	SO4	1	0

## 5.7 Other polymers [i](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.



## 6 Fit of model and data ⓘ

### 6.1 Protein, DNA and RNA chains ⓘ

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	448/449 (99%)	0.18	32 (7%) 22 25	7, 13, 30, 44	2 (0%)
2	B	193/194 (99%)	0.52	30 (15%) 5 5	8, 14, 38, 46	0
All	All	641/643 (99%)	0.28	62 (9%) 13 15	7, 13, 33, 46	2 (0%)

All (62) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	447	THR	9.5
2	B	525	CYS	7.7
1	A	224	PHE	7.1
1	A	235	PRO	6.7
1	A	234	PRO	6.4
2	B	502	MET	6.1
2	B	523	PHE	6.1
2	B	629	VAL	6.0
1	A	264	ASP	6.0
1	A	262	SER	5.5
2	B	667	GLU	5.4
1	A	127	SER	5.3
1	A	236	GLU	5.2
1	A	446	THR	5.1
1	A	448	ASP	5.1
1	A	232	ALA	5.1
1	A	237	GLY	4.9
2	B	526	HIS	4.7
2	B	631	ASP	4.6
2	B	633	GLU	4.6
1	A	233	LEU	4.4
2	B	519	ILE	4.4
1	A	223	ILE	4.3
2	B	503	ILE	4.1

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type	RSRZ
1	A	225	SER	4.0
2	B	632	LYS	3.9
2	B	630	ASN	3.8
1	A	263	ALA	3.8
2	B	577	ARG	3.7
2	B	520	LEU	3.6
1	A	221	GLU	3.6
1	A	3	TYR	3.5
1	A	445	LYS	3.4
2	B	634	LEU	3.4
1	A	220	GLY	3.3
2	B	524	ASN	3.2
2	B	527	ASP	3.2
1	A	121	LYS	3.0
1	A	128	LEU	3.0
1	A	132	CYS	2.9
1	A	226	SER	2.8
2	B	666	GLY	2.8
2	B	528	SER	2.7
1	A	282	LYS	2.7
2	B	580	LYS	2.7
1	A	231	ALA	2.7
1	A	222	SER	2.6
2	B	635	LEU	2.6
2	B	532	GLN	2.5
2	B	670	VAL	2.5
1	A	129	ASN	2.4
2	B	578	ARG	2.4
1	A	227	LEU	2.2
1	A	124	TYR	2.2
2	B	575	SER	2.1
2	B	506	GLN	2.1
2	B	530	LEU	2.1
1	A	228	ALA	2.1
2	B	581	LEU	2.1
1	A	97	LYS	2.1
2	B	504	ASN	2.0
2	B	628	ASP	2.0

## 6.2 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

### 6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

### 6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors( $\text{\AA}^2$ )	Q<0.9
5	SO4	B	1697	5/5	0.65	0.20	45,47,49,50	0
3	EDO	A	1450	4/4	0.66	0.25	41,41,41,43	0
5	SO4	A	1452	5/5	0.73	0.18	51,52,53,53	0
4	OXY	A	1451	2/2	0.87	0.12	36,36,36,38	0
5	SO4	B	1698	5/5	0.91	0.12	36,36,37,37	0
3	EDO	B	1696	4/4	0.92	0.10	24,26,27,28	0
3	EDO	B	1699	4/4	0.93	0.16	12,22,23,26	0
3	EDO	A	1449	4/4	0.95	0.09	15,21,22,22	0
3	EDO	A	1448	4/4	0.98	0.04	8,10,11,11	0
3	EDO	B	1695	4/4	0.98	0.04	11,12,13,13	0
6	FES	A	1453	4/4	0.99	0.02	11,12,13,14	0
7	FE	A	1454	1/1	0.99	0.11	22,22,22,22	0

### 6.5 Other polymers [i](#)

There are no such residues in this entry.