



# wwPDB X-ray Structure Validation Summary Report ⓘ

Mar 12, 2026 – 03:33 PM UTC

PDB ID : 2O7D / pdb\_00002o7d  
Title : Tyrosine ammonia-lyase from Rhodobacter sphaeroides, complexed with caffeine  
Authors : Louie, G.V.; Bowman, M.E.; Moffitt, M.C.; Baiga, T.J.; Moore, B.S.; Noel, J.P.  
Deposited on : 2006-12-10  
Resolution : 1.90 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4-5-2 with Phenix2.0  
Mogul : 2022.3.0, CSD as543be (2022)  
Xtriage (Phenix) : 2.0  
EDS : 3.0  
Percentile statistics : 20250101.v01 (using entries in the PDB archive January 1st 2025)  
CCP4 : 9.0.010 (Gargrove)  
Density-Fitness : 1.0.12  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : 2.49


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

*X-RAY DIFFRACTION*

The reported resolution of this entry is 1.90 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	180053	7789 (1.90-1.90)
Clashscore	190562	8410 (1.90-1.90)
Ramachandran outliers	187476	8333 (1.90-1.90)
Sidechain outliers	187428	8333 (1.90-1.90)
RSRZ outliers	180081	7790 (1.90-1.90)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	521	 5% 88% 10% ..
1	B	521	 2% 84% 14% ..
1	C	521	 2% 87% 11% ..
1	D	521	 5% 85% 13% ..
1	E	521	 4% 89% 9% ..

*Continued on next page...*

Continued from previous page...

Mol	Chain	Length	Quality of chain
1	F	521	<div><div>%</div><div><div></div><div>88%</div><div>10% ..</div></div></div>
1	G	521	<div><div>2%</div><div><div></div><div>88%</div><div>10% .</div></div></div>
1	H	521	<div><div>6%</div><div><div></div><div>86%</div><div>12% ..</div></div></div>

## 2 Entry composition

There are 3 unique types of molecules in this entry. The entry contains 33789 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Putative histidine ammonia-lyase.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	515	Total	C	N	O	S	0	0	0
			3813	2363	731	705	14			
1	B	515	Total	C	N	O	S	0	0	0
			3813	2363	731	705	14			
1	C	514	Total	C	N	O	S	0	0	0
			3806	2358	730	704	14			
1	D	514	Total	C	N	O	S	0	0	0
			3806	2358	730	704	14			
1	E	514	Total	C	N	O	S	0	0	0
			3806	2358	730	704	14			
1	F	514	Total	C	N	O	S	0	0	0
			3806	2358	730	704	14			
1	G	514	Total	C	N	O	S	0	0	0
			3806	2358	730	704	14			
1	H	514	Total	C	N	O	S	0	0	0
			3806	2358	730	704	14			

There are 24 discrepancies between the modelled and reference sequences:


Chain	Residue	Modelled	Actual	Comment	Reference
A	149	MDO	ALA	SEE REMARK 999	UNP Q3IWB0
A	149	MDO	SER	SEE REMARK 999	UNP Q3IWB0
A	149	MDO	GLY	SEE REMARK 999	UNP Q3IWB0
B	149	MDO	ALA	SEE REMARK 999	UNP Q3IWB0
B	149	MDO	SER	SEE REMARK 999	UNP Q3IWB0
B	149	MDO	GLY	SEE REMARK 999	UNP Q3IWB0
C	149	MDO	ALA	SEE REMARK 999	UNP Q3IWB0
C	149	MDO	SER	SEE REMARK 999	UNP Q3IWB0
C	149	MDO	GLY	SEE REMARK 999	UNP Q3IWB0
D	149	MDO	ALA	SEE REMARK 999	UNP Q3IWB0
D	149	MDO	SER	SEE REMARK 999	UNP Q3IWB0
D	149	MDO	GLY	SEE REMARK 999	UNP Q3IWB0
E	149	MDO	ALA	SEE REMARK 999	UNP Q3IWB0

*Continued on next page...*

Continued from previous page...

Chain	Residue	Modelled	Actual	Comment	Reference
E	149	MDO	SER	SEE REMARK 999	UNP Q3IWB0
E	149	MDO	GLY	SEE REMARK 999	UNP Q3IWB0
F	149	MDO	ALA	SEE REMARK 999	UNP Q3IWB0
F	149	MDO	SER	SEE REMARK 999	UNP Q3IWB0
F	149	MDO	GLY	SEE REMARK 999	UNP Q3IWB0
G	149	MDO	ALA	SEE REMARK 999	UNP Q3IWB0
G	149	MDO	SER	SEE REMARK 999	UNP Q3IWB0
G	149	MDO	GLY	SEE REMARK 999	UNP Q3IWB0
H	149	MDO	ALA	SEE REMARK 999	UNP Q3IWB0
H	149	MDO	SER	SEE REMARK 999	UNP Q3IWB0
H	149	MDO	GLY	SEE REMARK 999	UNP Q3IWB0

- Molecule 2 is CAFFEIC ACID (CCD ID: DHC) (formula:  $C_9H_8O_4$ ).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
2	A	1	Total	C	O	0	0
			13	9	4		
2	B	1	Total	C	O	0	0
			13	9	4		
2	C	1	Total	C	O	0	0
			13	9	4		
2	D	1	Total	C	O	0	0
			13	9	4		
2	E	1	Total	C	O	0	0
			13	9	4		
2	F	1	Total	C	O	0	0
			13	9	4		

Continued on next page...

*Continued from previous page...*

Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
2	G	1	Total	C	O	0	0
			13	9	4		
2	H	1	Total	C	O	0	0
			13	9	4		


- Molecule 3 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	A	360	Total	O	0	0
			360	360		
3	B	369	Total	O	0	0
			369	369		
3	C	407	Total	O	0	0
			407	407		
3	D	404	Total	O	0	0
			404	404		
3	E	382	Total	O	0	0
			382	382		
3	F	491	Total	O	0	0
			491	491		
3	G	431	Total	O	0	0
			431	431		
3	H	379	Total	O	0	0
			379	379		


### 3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: Putative histidine ammonia-lyase




- Molecule 1: Putative histidine ammonia-lyase





- Molecule 1: Putative histidine ammonia-lyase









• Molecule 1: Putative histidine ammonia-lyase




• Molecule 1: Putative histidine ammonia-lyase




• Molecule 1: Putative histidine ammonia-lyase





• Molecule 1: Putative histidine ammonia-lyase







● Molecule 1: Putative histidine ammonia-lyase



## 4 Data and refinement statistics

Property	Value	Source
Space group	P 1 21 1	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	87.61Å 154.90Å 164.16Å 90.00° 94.06° 90.00°	Depositor
Resolution (Å)	500.00 – 1.90 163.75 – 1.90	Depositor EDS
% Data completeness (in resolution range)	98.9 (500.00-1.90) 98.9 (163.75-1.90)	Depositor EDS
$R_{merge}$	0.11	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	1.79 (at 1.85Å)	Xtriage
Refinement program	CNS 1.1	Depositor
R, $R_{free}$	0.175 , 0.198 0.175 , 0.197	Depositor DCC
$R_{free}$ test set	17030 reflections (4.59%)	wwPDB-VP
Wilson B-factor (Å <sup>2</sup> )	18.2	Xtriage
Anisotropy	0.233	Xtriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.35 , 43.5	EDS
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.50$ , $\langle L^2 \rangle = 0.33$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
$F_o, F_c$ correlation	0.96	EDS
Total number of atoms	33789	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	21.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 3.33% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

## 5 Model quality [i](#)

### 5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: MDO, DHC

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z  > 5$	RMSZ	$\# Z  > 5$
1	A	0.36	0/3862	0.86	6/5253 (0.1%)
1	B	0.36	0/3862	0.86	7/5253 (0.1%)
1	C	0.38	0/3854	0.88	11/5242 (0.2%)
1	D	0.37	0/3854	0.87	10/5242 (0.2%)
1	E	0.37	0/3854	0.86	6/5242 (0.1%)
1	F	0.38	0/3854	0.88	8/5242 (0.2%)
1	G	0.38	0/3854	0.87	8/5242 (0.2%)
1	H	0.36	0/3854	0.84	7/5242 (0.1%)
All	All	0.37	0/30848	0.86	63/41958 (0.2%)

There are no bond length outliers.

The worst 5 of 63 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	F	252	ARG	CA-C-N	7.22	128.87	119.84
1	F	252	ARG	C-N-CA	7.22	128.87	119.84
1	C	497	PHE	CA-C-N	7.13	124.86	119.66
1	C	497	PHE	C-N-CA	7.13	124.86	119.66
1	E	252	ARG	CA-C-N	6.97	128.55	119.84

There are no chirality outliers.

There are no planarity outliers.

### 5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	3813	0	3857	35	0
1	B	3813	0	3857	52	0
1	C	3806	0	3849	43	0
1	D	3806	0	3849	46	0
1	E	3806	0	3849	33	0
1	F	3806	0	3849	38	0
1	G	3806	0	3849	32	0
1	H	3806	0	3849	54	0
2	A	13	0	7	0	0
2	B	13	0	7	1	0
2	C	13	0	6	1	0
2	D	13	0	5	1	0
2	E	13	0	6	1	0
2	F	13	0	6	1	0
2	G	13	0	6	1	0
2	H	13	0	6	1	0
3	A	360	0	0	3	0
3	B	369	0	0	5	0
3	C	407	0	0	5	0
3	D	404	0	0	6	0
3	E	382	0	0	5	0
3	F	491	0	0	4	0
3	G	431	0	0	2	0
3	H	379	0	0	5	0
All	All	33789	0	30857	303	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 5.

The worst 5 of 303 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:F:153:LEU:HD22	1:F:202:VAL:HG12	1.63	0.80
1:D:153:LEU:HD22	1:D:202:VAL:HG12	1.64	0.79
1:E:487:LYS:O	1:E:491:GLN:HG3	1.83	0.78
1:B:153:LEU:HD22	1:B:202:VAL:HG12	1.70	0.74
1:B:487:LYS:O	1:B:491:GLN:HG3	1.87	0.74

There are no symmetry-related clashes.

## 5.3 Torsion angles [i](#)

### 5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	510/521 (98%)	499 (98%)	9 (2%)	2 (0%)	30	22
1	B	510/521 (98%)	503 (99%)	6 (1%)	1 (0%)	43	36
1	C	509/521 (98%)	503 (99%)	5 (1%)	1 (0%)	43	36
1	D	509/521 (98%)	499 (98%)	9 (2%)	1 (0%)	43	36
1	E	509/521 (98%)	501 (98%)	7 (1%)	1 (0%)	43	36
1	F	509/521 (98%)	502 (99%)	6 (1%)	1 (0%)	43	36
1	G	509/521 (98%)	503 (99%)	5 (1%)	1 (0%)	43	36
1	H	509/521 (98%)	500 (98%)	8 (2%)	1 (0%)	43	36
All	All	4074/4168 (98%)	4010 (98%)	55 (1%)	9 (0%)	43	36

5 of 9 Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	E	253	PRO
1	F	253	PRO
1	A	8	LYS
1	C	253	PRO
1	D	253	PRO

### 5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	384/389 (99%)	383 (100%)	1 (0%)	86	88

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	B	384/389 (99%)	383 (100%)	1 (0%)	86	88
1	C	383/389 (98%)	382 (100%)	1 (0%)	86	88
1	D	383/389 (98%)	383 (100%)	0	100	100
1	E	383/389 (98%)	382 (100%)	1 (0%)	86	88
1	F	383/389 (98%)	382 (100%)	1 (0%)	86	88
1	G	383/389 (98%)	382 (100%)	1 (0%)	86	88
1	H	383/389 (98%)	382 (100%)	1 (0%)	86	88
All	All	3066/3112 (98%)	3059 (100%)	7 (0%)	87	90

5 of 7 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	E	443	THR
1	F	172	ARG
1	H	54	ARG
1	G	443	THR
1	C	80	ARG

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 41 such sidechains are listed below:

Mol	Chain	Res	Type
1	F	436	GLN
1	H	78	ASN
1	F	491	GLN
1	G	435	ASN
1	H	195	HIS

### 5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

8 non-standard protein/DNA/RNA residues are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The

Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z  > 2	Counts	RMSZ	# Z  > 2
1	MDO	D	149	1	11,13,14	2.47	4 (36%)	15,18,20	2.64	5 (33%)
1	MDO	B	149	1	11,13,14	2.47	4 (36%)	15,18,20	2.70	5 (33%)
1	MDO	F	149	1	11,13,14	2.44	4 (36%)	15,18,20	2.68	5 (33%)
1	MDO	G	149	1	11,13,14	2.40	4 (36%)	15,18,20	2.75	5 (33%)
1	MDO	H	149	1	11,13,14	2.44	5 (45%)	15,18,20	2.65	5 (33%)
1	MDO	C	149	1	11,13,14	2.46	4 (36%)	15,18,20	2.59	5 (33%)
1	MDO	A	149	1	11,13,14	2.42	4 (36%)	15,18,20	2.73	5 (33%)
1	MDO	E	149	1	11,13,14	2.40	4 (36%)	15,18,20	2.67	5 (33%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
1	MDO	D	149	1	-	0/4/23/24	0/1/1/1
1	MDO	B	149	1	-	0/4/23/24	0/1/1/1
1	MDO	F	149	1	-	0/4/23/24	0/1/1/1
1	MDO	G	149	1	-	0/4/23/24	0/1/1/1
1	MDO	H	149	1	-	0/4/23/24	0/1/1/1
1	MDO	C	149	1	-	0/4/23/24	0/1/1/1
1	MDO	A	149	1	-	0/4/23/24	0/1/1/1
1	MDO	E	149	1	-	0/4/23/24	0/1/1/1

The worst 5 of 33 bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	H	149	MDO	O2-C2	5.43	1.34	1.23
1	B	149	MDO	O2-C2	5.35	1.34	1.23
1	D	149	MDO	O2-C2	5.31	1.33	1.23
1	C	149	MDO	O2-C2	5.12	1.33	1.23
1	F	149	MDO	O2-C2	5.12	1.33	1.23

The worst 5 of 40 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	149	MDO	CA2-C2-N3	7.13	109.49	103.50
1	H	149	MDO	CA2-C2-N3	7.01	109.39	103.50
1	G	149	MDO	CA2-C2-N3	6.95	109.34	103.50
1	E	149	MDO	CA2-C2-N3	6.94	109.33	103.50
1	F	149	MDO	CA2-C2-N3	6.87	109.27	103.50

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

## 5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

## 5.6 Ligand geometry [i](#)

8 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z  > 2$	Counts	RMSZ	$\# Z  > 2$
2	DHC	H	701	-	13,13,13	2.51	5 (38%)	17,17,17	0.83	0
2	DHC	B	701	-	13,13,13	2.53	5 (38%)	17,17,17	0.86	0
2	DHC	A	701	-	13,13,13	2.45	6 (46%)	17,17,17	0.82	0
2	DHC	D	701	-	13,13,13	2.55	6 (46%)	17,17,17	0.83	0
2	DHC	E	701	-	13,13,13	2.49	6 (46%)	17,17,17	0.82	0
2	DHC	G	701	-	13,13,13	2.44	6 (46%)	17,17,17	0.83	0
2	DHC	F	701	-	13,13,13	2.53	7 (53%)	17,17,17	0.93	0
2	DHC	C	701	-	13,13,13	2.49	7 (53%)	17,17,17	0.90	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral



centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	DHC	H	701	-	-	2/5/5/5	0/1/1/1
2	DHC	B	701	-	-	2/5/5/5	0/1/1/1
2	DHC	A	701	-	-	2/5/5/5	0/1/1/1
2	DHC	D	701	-	-	4/5/5/5	0/1/1/1
2	DHC	E	701	-	-	2/5/5/5	0/1/1/1
2	DHC	G	701	-	-	2/5/5/5	0/1/1/1
2	DHC	F	701	-	-	2/5/5/5	0/1/1/1
2	DHC	C	701	-	-	2/5/5/5	0/1/1/1

The worst 5 of 48 bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	D	701	DHC	C2'-C3'	5.15	1.46	1.38
2	E	701	DHC	C2'-C3'	4.79	1.45	1.38
2	B	701	DHC	C6'-C5'	4.77	1.46	1.38
2	F	701	DHC	C2'-C3'	4.70	1.45	1.38
2	C	701	DHC	C6'-C5'	4.66	1.46	1.38

There are no bond angle outliers.

There are no chirality outliers.

5 of 18 torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
2	F	701	DHC	O2-C1-C2-C3
2	F	701	DHC	O1-C1-C2-C3
2	G	701	DHC	O1-C1-C2-C3
2	E	701	DHC	O1-C1-C2-C3
2	E	701	DHC	O2-C1-C2-C3

There are no ring outliers.

7 monomers are involved in 7 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	H	701	DHC	1	0
2	B	701	DHC	1	0
2	D	701	DHC	1	0

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	E	701	DHC	1	0
2	G	701	DHC	1	0
2	F	701	DHC	1	0
2	C	701	DHC	1	0

## 5.7 Other polymers [i](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

## 6 Fit of model and data [i](#)

### 6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	514/521 (98%)	0.09	24 (4%) 36 39	10, 20, 41, 61	0
1	B	514/521 (98%)	0.01	8 (1%) 70 74	10, 20, 38, 53	0
1	C	513/521 (98%)	-0.11	12 (2%) 61 65	9, 18, 35, 48	0
1	D	513/521 (98%)	0.05	27 (5%) 32 34	8, 18, 41, 55	0
1	E	513/521 (98%)	-0.04	19 (3%) 45 48	9, 18, 37, 66	0
1	F	513/521 (98%)	-0.26	6 (1%) 76 79	8, 15, 32, 46	0
1	G	513/521 (98%)	-0.16	9 (1%) 67 71	7, 17, 33, 47	0
1	H	513/521 (98%)	0.17	32 (6%) 26 28	9, 20, 43, 59	0
All	All	4106/4168 (98%)	-0.03	137 (3%) 49 52	7, 18, 38, 66	0

The worst 5 of 137 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	D	74	ILE	8.5
1	A	296	GLY	7.8
1	H	74	ILE	7.3
1	A	289	ILE	6.3
1	D	70	ALA	6.0

### 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled ‘Q< 0.9’ lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å <sup>2</sup> )	Q<0.9
1	MDO	H	149	13/14	0.82	0.12	19,23,29,30	0

*Continued on next page...*

*Continued from previous page...*

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors( $\text{\AA}^2$ )	Q<0.9
1	MDO	B	149	13/14	0.85	0.11	19,22,27,28	0
1	MDO	F	149	13/14	0.89	0.10	15,19,23,27	0
1	MDO	E	149	13/14	0.89	0.09	13,17,22,23	0
1	MDO	A	149	13/14	0.90	0.09	14,19,21,24	0
1	MDO	C	149	13/14	0.90	0.09	13,15,20,25	0
1	MDO	G	149	13/14	0.90	0.09	14,17,21,22	0
1	MDO	D	149	13/14	0.90	0.10	18,21,29,29	0

### 6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

### 6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors( $\text{\AA}^2$ )	Q<0.9
2	DHC	D	701	13/13	0.77	0.16	34,39,44,47	0
2	DHC	A	701	13/13	0.85	0.16	34,42,46,48	0
2	DHC	H	701	13/13	0.85	0.14	33,40,49,53	0
2	DHC	B	701	13/13	0.86	0.11	27,31,37,40	0
2	DHC	E	701	13/13	0.87	0.10	24,30,34,34	0
2	DHC	G	701	13/13	0.88	0.11	24,28,36,39	0
2	DHC	C	701	13/13	0.88	0.10	21,26,33,36	0
2	DHC	F	701	13/13	0.90	0.09	24,28,32,36	0

### 6.5 Other polymers [i](#)

There are no such residues in this entry.