



wwPDB X-ray Structure Validation Summary Report ⓘ

Mar 5, 2026 – 06:20 PM UTC

PDB ID : 2OKX / pdb_00002okx
Title : Crystal structure of GH78 family rhamnosidase of Bacillus SP. GL1 AT 1.9 Å
Authors : Cui, Z.; Mikami, B.; Hashimoto, W.; Murata, K.
Deposited on : 2007-01-17
Resolution : 1.90 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4-5-2 with Phenix2.0
Mogul	:	2022.3.0, CSD as543be (2022)
Xtriage (Phenix)	:	2.0
EDS	:	3.0
Percentile statistics	:	20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4	:	9.0.010 (Gargrove)
Density-Fitness	:	1.0.12
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 1.90 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	7789 (1.90-1.90)
Clashscore	190562	8410 (1.90-1.90)
Ramachandran outliers	187476	8333 (1.90-1.90)
Sidechain outliers	187428	8333 (1.90-1.90)
RSRZ outliers	180081	7790 (1.90-1.90)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	956	% 77% 21% .
1	B	956	% 78% 19% .

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
3	GOL	B	3037	-	-	X	-

2 Entry composition [i](#)

There are 4 unique types of molecules in this entry. The entry contains 17068 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.


- Molecule 1 is a protein called Rhamnosidase B.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	954	Total	C	N	O	S	0	11	0
			7531	4773	1309	1420	29			
1	B	954	Total	C	N	O	S	0	9	0
			7520	4767	1304	1420	29			

- Molecule 2 is CALCIUM ION (CCD ID: CA) (formula: Ca).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
2	A	2	Total	Ca	0	0
			2	2		
2	B	2	Total	Ca	0	0
			2	2		

- Molecule 3 is GLYCEROL (CCD ID: GOL) (formula: C₃H₈O₃).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
3	A	1	Total	C	O	0	0
			6	3	3		
3	A	1	Total	C	O	0	0
			6	3	3		
3	A	1	Total	C	O	0	0
			6	3	3		
3	A	1	Total	C	O	0	0
			6	3	3		
3	A	1	Total	C	O	0	0
			6	3	3		
3	A	1	Total	C	O	0	0
			6	3	3		
3	A	1	Total	C	O	0	0
			6	3	3		
3	A	1	Total	C	O	0	0
			6	3	3		
3	A	1	Total	C	O	0	0
			6	3	3		
3	A	1	Total	C	O	0	0
			6	3	3		
3	A	1	Total	C	O	0	0
			6	3	3		
3	A	1	Total	C	O	0	0
			6	3	3		
3	A	1	Total	C	O	0	0
			6	3	3		
3	A	1	Total	C	O	0	0
			6	3	3		
3	A	1	Total	C	O	0	0
			6	3	3		
3	B	1	Total	C	O	0	0
			6	3	3		


Continued on next page...

Continued from previous page...

Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
3	B	1	Total	C	O	0	0
			6	3	3		
3	B	1	Total	C	O	0	0
			6	3	3		
3	B	1	Total	C	O	0	0
			6	3	3		
3	B	1	Total	C	O	0	0
			6	3	3		
3	B	1	Total	C	O	0	0
			6	3	3		
3	B	1	Total	C	O	0	0
			6	3	3		
3	B	1	Total	C	O	0	0
			6	3	3		
3	B	1	Total	C	O	0	0
			6	3	3		
3	B	1	Total	C	O	0	0
			6	3	3		
3	B	1	Total	C	O	0	0
			6	3	3		
3	B	1	Total	C	O	0	0
			6	3	3		
3	B	1	Total	C	O	0	0
			6	3	3		
3	B	1	Total	C	O	0	0
			6	3	3		
3	B	1	Total	C	O	0	0
			6	3	3		
3	B	1	Total	C	O	0	0
			6	3	3		
3	B	1	Total	C	O	0	0
			6	3	3		
3	B	1	Total	C	O	0	0
			6	3	3		

- Molecule 4 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	A	880	Total 880	O 880	0	0
4	B	875	Total 875	O 875	0	0



4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, α , β , γ	97.74Å 119.98Å 207.35Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	15.00 – 1.90 15.00 – 1.90	Depositor EDS
% Data completeness (in resolution range)	98.5 (15.00-1.90) 98.4 (15.00-1.90)	Depositor EDS
R_{merge}	0.07	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	2.29 (at 1.91Å)	Xtriage
Refinement program	CNS 1.1	Depositor
R, R_{free}	0.182 , 0.214 0.184 , 0.215	Depositor DCC
R_{free} test set	18716 reflections (9.91%)	wwPDB-VP
Wilson B-factor (Å ²)	24.2	Xtriage
Anisotropy	0.439	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.39 , 55.5	EDS
L-test for twinning ²	$\langle L \rangle = 0.50$, $\langle L^2 \rangle = 0.33$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.96	EDS
Total number of atoms	17068	wwPDB-VP
Average B, all atoms (Å ²)	29.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 3.14% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: CA, GOL

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.34	0/7782	0.90	29/10613 (0.3%)
1	B	0.35	0/7763	0.91	29/10588 (0.3%)
All	All	0.35	0/15545	0.91	58/21201 (0.3%)

There are no bond length outliers.

The worst 5 of 58 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	B	67	TRP	N-CA-C	-11.92	94.82	109.72
1	A	67	TRP	N-CA-C	-11.29	95.61	109.72
1	A	491	ARG	N-CA-C	-7.81	98.26	110.36
1	B	491[A]	ARG	N-CA-C	-7.33	99.00	110.36
1	B	491[B]	ARG	N-CA-C	-7.33	99.00	110.36

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	7531	0	7202	175	0
1	B	7520	0	7187	155	0
2	A	2	0	0	0	0
2	B	2	0	0	0	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
3	A	126	0	143	12	0
3	B	132	0	151	14	0
4	A	880	0	0	17	0
4	B	875	0	0	17	0
All	All	17068	0	14683	329	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 11.

The worst 5 of 329 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:765:GLN:H	1:B:765:GLN:HE21	0.99	0.98
1:B:746:ARG:HH11	1:B:746:ARG:HB3	1.32	0.94
1:A:765:GLN:H	1:A:765:GLN:HE21	1.13	0.90
1:A:410:ARG:HH22	3:A:3033:GOL:H32	1.38	0.87
1:A:236:HIS:HD2	1:A:238:ASN:H	1.23	0.87

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	963/956 (101%)	929 (96%)	30 (3%)	4 (0%)	30	22
1	B	961/956 (100%)	922 (96%)	37 (4%)	2 (0%)	43	36
All	All	1924/1912 (101%)	1851 (96%)	67 (4%)	6 (0%)	36	29

5 of 6 Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	692	GLU
1	B	614	TYR
1	A	614	TYR
1	A	37	GLU
1	A	686	PRO

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	787/777 (101%)	770 (98%)	17 (2%)	45	42
1	B	785/777 (101%)	764 (97%)	21 (3%)	39	34
All	All	1572/1554 (101%)	1534 (98%)	38 (2%)	44	38

5 of 38 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	B	587	ASN
1	B	765	GLN
1	B	617	GLN
1	B	743[A]	GLU
1	B	862	HIS

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 59 such sidechains are listed below:

Mol	Chain	Res	Type
1	A	894	GLN
1	B	849	ASN
1	B	227	ASN
1	B	817	GLN
1	B	680	ASN

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

Of 47 ligands modelled in this entry, 4 are monoatomic - leaving 43 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
3	GOL	A	3003	-	5,5,5	1.22	1 (20%)	5,5,5	0.37	0
3	GOL	A	3043	-	5,5,5	1.21	1 (20%)	5,5,5	0.37	0
3	GOL	B	3016	-	5,5,5	1.19	1 (20%)	5,5,5	0.28	0
3	GOL	B	3029	-	5,5,5	1.21	1 (20%)	5,5,5	0.30	0
3	GOL	B	3041	-	5,5,5	1.32	1 (20%)	5,5,5	0.58	0
3	GOL	A	3008	-	5,5,5	1.19	1 (20%)	5,5,5	0.29	0
3	GOL	A	3006	-	5,5,5	1.21	1 (20%)	5,5,5	0.36	0
3	GOL	A	3030	-	5,5,5	1.19	1 (20%)	5,5,5	0.28	0
3	GOL	A	3026	-	5,5,5	1.17	1 (20%)	5,5,5	0.30	0
3	GOL	A	3007	-	5,5,5	1.17	1 (20%)	5,5,5	0.37	0
3	GOL	A	3033	-	5,5,5	1.22	1 (20%)	5,5,5	0.34	0
3	GOL	A	3004	-	5,5,5	1.19	1 (20%)	5,5,5	0.29	0
3	GOL	B	3025	-	5,5,5	1.19	1 (20%)	5,5,5	0.27	0
3	GOL	B	3037	-	5,5,5	1.21	1 (20%)	5,5,5	0.33	0
3	GOL	A	3040	-	5,5,5	1.19	1 (20%)	5,5,5	0.37	0
3	GOL	B	3018	-	5,5,5	1.19	1 (20%)	5,5,5	0.30	0
3	GOL	A	3017	-	5,5,5	1.19	1 (20%)	5,5,5	0.30	0
3	GOL	B	3028	-	5,5,5	1.20	1 (20%)	5,5,5	0.32	0

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
3	GOL	B	3036	-	5,5,5	1.17	1 (20%)	5,5,5	0.27	0
3	GOL	A	3022	-	5,5,5	1.25	1 (20%)	5,5,5	0.39	0
3	GOL	A	3039	-	5,5,5	1.39	1 (20%)	5,5,5	0.56	0
3	GOL	B	3024	-	5,5,5	1.22	1 (20%)	5,5,5	0.34	0
3	GOL	A	3027	-	5,5,5	1.20	1 (20%)	5,5,5	0.31	0
3	GOL	B	3032	-	5,5,5	1.19	1 (20%)	5,5,5	0.28	0
3	GOL	B	3021	-	5,5,5	1.27	1 (20%)	5,5,5	0.40	0
3	GOL	B	3038	-	5,5,5	1.19	1 (20%)	5,5,5	0.31	0
3	GOL	A	3002	-	5,5,5	1.22	1 (20%)	5,5,5	0.31	0
3	GOL	A	3019	-	5,5,5	1.20	1 (20%)	5,5,5	0.29	0
3	GOL	B	3034	-	5,5,5	1.18	1 (20%)	5,5,5	0.35	0
3	GOL	A	3001	-	5,5,5	1.19	1 (20%)	5,5,5	0.32	0
3	GOL	B	3020	-	5,5,5	1.19	1 (20%)	5,5,5	0.28	0
3	GOL	B	3013	-	5,5,5	1.19	1 (20%)	5,5,5	0.29	0
3	GOL	A	3023	-	5,5,5	1.20	1 (20%)	5,5,5	0.32	0
3	GOL	B	3010	-	5,5,5	1.20	1 (20%)	5,5,5	0.30	0
3	GOL	A	3015	-	5,5,5	1.20	1 (20%)	5,5,5	0.31	0
3	GOL	B	3011	-	5,5,5	1.17	1 (20%)	5,5,5	0.29	0
3	GOL	B	3009	-	5,5,5	1.14	1 (20%)	5,5,5	0.29	0
3	GOL	A	3035	-	5,5,5	1.22	1 (20%)	5,5,5	0.31	0
3	GOL	A	3031	-	5,5,5	1.19	1 (20%)	5,5,5	0.31	0
3	GOL	B	3005	-	5,5,5	1.20	1 (20%)	5,5,5	0.30	0
3	GOL	B	3012	-	5,5,5	1.23	1 (20%)	5,5,5	0.35	0
3	GOL	B	3014	-	5,5,5	1.14	1 (20%)	5,5,5	0.31	0
3	GOL	B	3042	-	5,5,5	1.21	1 (20%)	5,5,5	0.36	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	GOL	A	3003	-	-	0/4/4/4	-
3	GOL	A	3043	-	-	0/4/4/4	-
3	GOL	B	3016	-	-	0/4/4/4	-
3	GOL	B	3029	-	-	0/4/4/4	-
3	GOL	B	3041	-	-	4/4/4/4	-

Continued on next page...

Continued from previous page...

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	GOL	A	3008	-	-	0/4/4/4	-
3	GOL	A	3006	-	-	0/4/4/4	-
3	GOL	A	3030	-	-	0/4/4/4	-
3	GOL	A	3026	-	-	0/4/4/4	-
3	GOL	A	3007	-	-	0/4/4/4	-
3	GOL	A	3033	-	-	0/4/4/4	-
3	GOL	A	3004	-	-	0/4/4/4	-
3	GOL	B	3025	-	-	0/4/4/4	-
3	GOL	B	3037	-	-	0/4/4/4	-
3	GOL	A	3040	-	-	0/4/4/4	-
3	GOL	B	3018	-	-	0/4/4/4	-
3	GOL	A	3017	-	-	0/4/4/4	-
3	GOL	B	3028	-	-	0/4/4/4	-
3	GOL	B	3036	-	-	0/4/4/4	-
3	GOL	A	3022	-	-	0/4/4/4	-
3	GOL	A	3039	-	-	1/4/4/4	-
3	GOL	B	3024	-	-	0/4/4/4	-
3	GOL	A	3027	-	-	0/4/4/4	-
3	GOL	B	3032	-	-	0/4/4/4	-
3	GOL	B	3021	-	-	0/4/4/4	-
3	GOL	B	3038	-	-	0/4/4/4	-
3	GOL	A	3002	-	-	0/4/4/4	-
3	GOL	A	3019	-	-	0/4/4/4	-
3	GOL	B	3034	-	-	0/4/4/4	-
3	GOL	A	3001	-	-	0/4/4/4	-
3	GOL	B	3020	-	-	0/4/4/4	-
3	GOL	B	3013	-	-	0/4/4/4	-
3	GOL	A	3023	-	-	0/4/4/4	-
3	GOL	B	3010	-	-	0/4/4/4	-
3	GOL	A	3015	-	-	0/4/4/4	-
3	GOL	B	3011	-	-	0/4/4/4	-
3	GOL	B	3009	-	-	0/4/4/4	-
3	GOL	A	3035	-	-	0/4/4/4	-
3	GOL	A	3031	-	-	0/4/4/4	-
3	GOL	B	3005	-	-	0/4/4/4	-
3	GOL	B	3012	-	-	0/4/4/4	-
3	GOL	B	3014	-	-	0/4/4/4	-
3	GOL	B	3042	-	-	2/4/4/4	-

The worst 5 of 43 bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
3	A	3039	GOL	O1-C1	-3.05	1.29	1.42
3	B	3041	GOL	O1-C1	-2.91	1.30	1.42
3	B	3012	GOL	O1-C1	-2.69	1.31	1.42
3	A	3003	GOL	O1-C1	-2.69	1.31	1.42
3	A	3002	GOL	O1-C1	-2.67	1.31	1.42

There are no bond angle outliers.

There are no chirality outliers.

5 of 7 torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
3	B	3041	GOL	O1-C1-C2-C3
3	B	3042	GOL	C1-C2-C3-O3
3	B	3042	GOL	O2-C2-C3-O3
3	B	3041	GOL	O1-C1-C2-O2
3	B	3041	GOL	C1-C2-C3-O3

There are no ring outliers.

17 monomers are involved in 26 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
3	A	3003	GOL	1	0
3	A	3043	GOL	1	0
3	B	3029	GOL	3	0
3	B	3041	GOL	1	0
3	A	3006	GOL	1	0
3	A	3030	GOL	2	0
3	A	3007	GOL	3	0
3	A	3033	GOL	1	0
3	B	3037	GOL	4	0
3	B	3018	GOL	1	0
3	A	3022	GOL	1	0
3	B	3024	GOL	1	0
3	A	3027	GOL	1	0
3	B	3021	GOL	1	0
3	B	3038	GOL	1	0
3	B	3034	GOL	2	0
3	A	3023	GOL	1	0

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	954/956 (99%)	-0.04	12 (1%) 75 78	10, 26, 46, 61	11 (1%)
1	B	954/956 (99%)	-0.13	10 (1%) 79 82	11, 25, 49, 64	9 (0%)
All	All	1908/1912 (99%)	-0.08	22 (1%) 76 79	10, 26, 47, 64	20 (1%)

The worst 5 of 22 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	367	LEU	3.9
1	B	118	ASP	3.7
1	B	367	LEU	3.6
1	A	7	ASN	3.1
1	B	36	VAL	2.9

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled ‘Q< 0.9’ lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors($\text{\AA}^2$)	Q<0.9
3	GOL	B	3038	6/6	0.56	0.21	64,65,66,67	0
3	GOL	B	3032	6/6	0.62	0.23	74,74,75,75	0
3	GOL	A	3031	6/6	0.63	0.19	84,85,85,86	0
3	GOL	B	3028	6/6	0.63	0.20	73,73,74,74	0
3	GOL	A	3008	6/6	0.67	0.17	54,58,59,59	0
3	GOL	B	3029	6/6	0.67	0.21	72,72,73,73	0
3	GOL	A	3027	6/6	0.69	0.22	72,73,73,75	0
3	GOL	A	3023	6/6	0.69	0.17	68,70,71,71	0
3	GOL	B	3037	6/6	0.70	0.25	67,68,69,71	0
3	GOL	A	3019	6/6	0.70	0.18	67,67,67,68	0
3	GOL	B	3024	6/6	0.72	0.19	67,68,68,69	0
3	GOL	B	3036	6/6	0.73	0.17	55,62,63,63	0
3	GOL	A	3030	6/6	0.73	0.15	68,68,68,69	0
3	GOL	A	3043	6/6	0.73	0.24	47,51,56,59	0
3	GOL	B	3042	6/6	0.73	0.21	47,49,51,54	0
3	GOL	A	3039	6/6	0.74	0.20	38,42,45,50	0
3	GOL	B	3041	6/6	0.74	0.23	58,60,62,64	0
3	GOL	A	3033	6/6	0.74	0.15	43,47,51,55	0
3	GOL	B	3018	6/6	0.75	0.20	74,75,75,75	0
3	GOL	B	3034	6/6	0.77	0.16	36,41,45,51	0
3	GOL	A	3017	6/6	0.77	0.17	66,67,67,67	0
3	GOL	A	3004	6/6	0.78	0.13	42,43,44,44	0
3	GOL	B	3012	6/6	0.78	0.14	42,44,45,50	0
3	GOL	B	3025	6/6	0.78	0.16	47,51,53,54	0
3	GOL	B	3020	6/6	0.79	0.18	41,48,48,49	0
3	GOL	A	3026	6/6	0.79	0.14	41,46,47,50	0
3	GOL	A	3035	6/6	0.80	0.16	51,54,55,56	0
3	GOL	A	3003	6/6	0.81	0.14	47,48,51,51	0
3	GOL	A	3002	6/6	0.83	0.12	43,45,47,47	0
3	GOL	B	3013	6/6	0.83	0.12	42,44,45,46	0
3	GOL	B	3016	6/6	0.83	0.14	50,55,56,57	0
3	GOL	B	3011	6/6	0.83	0.13	44,47,48,48	0
3	GOL	A	3007	6/6	0.84	0.11	25,29,30,31	0
3	GOL	A	3015	6/6	0.85	0.11	50,53,53,55	0
3	GOL	A	3022	6/6	0.85	0.15	34,35,36,37	0
3	GOL	B	3021	6/6	0.86	0.15	33,34,36,36	0
3	GOL	A	3006	6/6	0.86	0.13	34,37,39,39	0
3	GOL	A	3001	6/6	0.88	0.10	33,35,38,43	0
3	GOL	A	3040	6/6	0.89	0.12	48,51,51,52	0
3	GOL	B	3005	6/6	0.89	0.10	39,42,43,46	0
3	GOL	B	3010	6/6	0.89	0.10	45,46,46,47	0
3	GOL	B	3014	6/6	0.92	0.08	22,26,26,26	0
3	GOL	B	3009	6/6	0.92	0.11	27,34,34,35	0

Continued on next page...

Continued from previous page...

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors($\text{\AA}^2$)	Q<0.9
2	CA	B	4003	1/1	0.99	0.10	30,30,30,30	0
2	CA	A	4001	1/1	0.99	0.05	29,29,29,29	0
2	CA	A	4004	1/1	0.99	0.06	28,28,28,28	0
2	CA	B	4002	1/1	0.99	0.08	31,31,31,31	0

6.5 Other polymers [i](#)

There are no such residues in this entry.