



# wwPDB X-ray Structure Validation Summary Report ⓘ

Mar 8, 2026 – 04:53 PM UTC

PDB ID : 2OS7 / pdb\_00002os7  
Title : Caf1M periplasmic chaperone tetramer  
Authors : Knight, S.D.; Zavialov, A.Z.  
Deposited on : 2007-02-05  
Resolution : 2.90 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4-5-2 with Phenix2.0  
Xtrriage (Phenix) : 2.0  
EDS : 3.0  
Percentile statistics : 20250101.v01 (using entries in the PDB archive January 1st 2025)  
CCP4 : 9.0.010 (Gargrove)  
Density-Fitness : 1.0.12  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : 2.49


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

*X-RAY DIFFRACTION*

The reported resolution of this entry is 2.90 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.




Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	180053	2481 (2.90-2.90)
Clashscore	190562	2690 (2.90-2.90)
Ramachandran outliers	187476	2623 (2.90-2.90)
Sidechain outliers	187428	2625 (2.90-2.90)
RSRZ outliers	180081	2481 (2.90-2.90)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	235	
1	B	235	
1	C	235	
1	D	235	
1	E	235	

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Length	Quality of chain
1	F	235	 <p>A horizontal bar chart representing the quality of chain. The bar is divided into four segments: a small red segment at the beginning labeled '2%', followed by a large green segment labeled '51%', a yellow segment labeled '26%', and a grey segment at the end labeled '19%'. A small black dot is visible on the yellow segment.</p>

## 2 Entry composition

There is only 1 type of molecule in this entry. The entry contains 9426 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.


- Molecule 1 is a protein called Chaperone protein caf1M.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	206	Total 1628	C 1053	N 274	O 297	S 4	0	0	0
1	B	196	Total 1550	C 1006	N 263	O 277	S 4	0	0	0
1	C	205	Total 1612	C 1041	N 271	O 296	S 4	0	0	0
1	D	195	Total 1523	C 991	N 256	O 272	S 4	0	0	0
1	E	202	Total 1603	C 1035	N 271	O 293	S 4	0	0	0
1	F	190	Total 1510	C 982	N 256	O 268	S 4	0	0	0


### 3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: Chaperone protein caf1M




- Molecule 1: Chaperone protein caf1M




- Molecule 1: Chaperone protein caf1M







● Molecule 1: Chaperone protein caf1M



● Molecule 1: Chaperone protein caf1M



● Molecule 1: Chaperone protein caf1M



## 4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 2	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	141.09Å 174.27Å 60.58Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	30.00 – 2.90 30.00 – 2.90	Depositor EDS
% Data completeness (in resolution range)	99.8 (30.00-2.90) 99.8 (30.00-2.90)	Depositor EDS
$R_{merge}$	0.17	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	2.15 (at 2.90Å)	Xtrriage
Refinement program	REFMAC 5.2.0005	Depositor
R, $R_{free}$	0.249 , 0.298 0.247 , 0.297	Depositor DCC
$R_{free}$ test set	1720 reflections (5.06%)	wwPDB-VP
Wilson B-factor (Å <sup>2</sup> )	56.5	Xtrriage
Anisotropy	0.625	Xtrriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.30 , 28.7	EDS
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.44$ , $\langle L^2 \rangle = 0.27$	Xtrriage
Estimated twinning fraction	No twinning to report.	Xtrriage
$F_o, F_c$ correlation	0.89	EDS
Total number of atoms	9426	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	45.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 11.63% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

## 5 Model quality i

### 5.1 Standard geometry i

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# $ Z  > 5$	RMSZ	# $ Z  > 5$
1	A	1.03	3/1668 (0.2%)	1.12	6/2259 (0.3%)
1	B	0.91	0/1588	1.05	7/2148 (0.3%)
1	C	0.95	0/1651	1.06	8/2237 (0.4%)
1	D	0.90	0/1561	1.05	4/2114 (0.2%)
1	E	0.93	0/1642	1.02	3/2224 (0.1%)
1	F	0.91	0/1546	1.04	3/2091 (0.1%)
All	All	0.94	3/9656 (0.0%)	1.06	31/13073 (0.2%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	C	0	1
1	D	0	1
1	E	0	1
1	F	0	1
All	All	0	4

All (3) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	206	LYS	N-CA	5.57	1.53	1.46
1	A	44	ILE	CA-CB	5.29	1.60	1.54
1	A	10	LYS	N-CA	5.13	1.52	1.46

The worst 5 of 31 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	103	PRO	CA-C-N	-8.36	111.28	120.14
1	A	103	PRO	C-N-CA	-8.36	111.28	120.14
1	A	27	ALA	N-CA-C	-7.36	99.02	110.42

*Continued on next page...*

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	B	87	PHE	CA-C-N	-7.09	112.67	119.76
1	B	87	PHE	C-N-CA	-7.09	112.67	119.76

There are no chirality outliers.

All (4) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	C	10	LYS	Peptide
1	D	146	GLU	Peptide
1	E	26	ASP	Peptide
1	F	162	VAL	Peptide

## 5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1628	0	1644	37	0
1	B	1550	0	1561	45	0
1	C	1612	0	1617	42	0
1	D	1523	0	1522	49	0
1	E	1603	0	1612	39	0
1	F	1510	0	1532	52	0
All	All	9426	0	9488	226	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 12.

The worst 5 of 226 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:C:126:VAL:HG22	1:F:137:CYS:SG	2.10	0.91
1:C:7:PHE:HB2	1:F:130:VAL:HG22	1.54	0.90
1:C:148:LYS:NZ	1:C:148:LYS:H	1.81	0.78
1:E:225:LEU:HB2	1:F:31:MET:HB2	1.66	0.77
1:D:152:ILE:HD11	1:D:226:ASP:HB2	1.66	0.77

There are no symmetry-related clashes.

## 5.3 Torsion angles [i](#)

### 5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	200/235 (85%)	184 (92%)	13 (6%)	3 (2%)	8	28
1	B	188/235 (80%)	172 (92%)	14 (7%)	2 (1%)	11	36
1	C	199/235 (85%)	180 (90%)	16 (8%)	3 (2%)	8	28
1	D	187/235 (80%)	169 (90%)	17 (9%)	1 (0%)	24	54
1	E	196/235 (83%)	180 (92%)	13 (7%)	3 (2%)	8	28
1	F	180/235 (77%)	166 (92%)	13 (7%)	1 (1%)	21	51
All	All	1150/1410 (82%)	1051 (91%)	86 (8%)	13 (1%)	11	36

5 of 13 Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	205	PRO
1	E	146	GLU
1	B	164	GLY
1	C	146	GLU
1	D	86	VAL

### 5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	180/204 (88%)	159 (88%)	21 (12%)	5	17

*Continued on next page...*

Continued from previous page...

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	B	170/204 (83%)	147 (86%)	23 (14%)	4	12
1	C	177/204 (87%)	154 (87%)	23 (13%)	4	13
1	D	165/204 (81%)	140 (85%)	25 (15%)	3	9
1	E	177/204 (87%)	156 (88%)	21 (12%)	5	16
1	F	167/204 (82%)	148 (89%)	19 (11%)	5	18
All	All	1036/1224 (85%)	904 (87%)	132 (13%)	4	14

5 of 132 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	F	22	ILE
1	F	64	THR
1	F	179	ILE
1	C	34	VAL
1	C	15	THR

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 14 such sidechains are listed below:

Mol	Chain	Res	Type
1	D	82	GLN
1	D	121	ASN
1	F	222	GLN
1	D	222	GLN
1	F	145	ASN

### 5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

## 5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

## 5.6 Ligand geometry [i](#)

There are no ligands in this entry.

## 5.7 Other polymers [i](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

## 6 Fit of model and data [i](#)

### 6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	206/235 (87%)	0.25	6 (2%) 53 45	35, 44, 52, 60	0
1	B	196/235 (83%)	0.19	6 (3%) 51 42	34, 45, 53, 64	0
1	C	205/235 (87%)	0.24	3 (1%) 72 64	35, 44, 52, 67	0
1	D	195/235 (82%)	0.30	1 (0%) 87 83	40, 46, 58, 68	0
1	E	202/235 (85%)	0.36	1 (0%) 87 83	34, 44, 51, 65	0
1	F	190/235 (80%)	0.26	4 (2%) 63 54	40, 45, 52, 56	0
All	All	1194/1410 (84%)	0.27	21 (1%) 67 59	34, 45, 53, 68	0

The worst 5 of 21 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	F	204	LEU	2.9
1	A	129	PHE	2.8
1	B	26	ASP	2.8
1	F	60	PRO	2.8
1	E	101	GLY	2.7

### 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

### 6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

### 6.4 Ligands [i](#)

There are no ligands in this entry.

## 6.5 Other polymers [i](#)

There are no such residues in this entry.